

ERRATUM

Open Access



# Erratum to: Comparison of a drug-eluting balloon first and then bare metal stent with a drug-eluting stent for treatment of de novo lesions: study protocol of a randomized controlled trial

Sang-Don Park<sup>†</sup>, Chang-Hwan Yoon<sup>†</sup>, Il-Young Oh, Jung-Won Suh, Young-Suk Cho, Tae-Jin Youn, Dong-Ju Choi and In-Ho Chae<sup>\*</sup>

Unfortunately, the original version of this article [1] contained an error. The correct paragraph can be found below:

The calculation of number of patients was not correct. It was wrote the non-inferiority margin as 0.1 mm. But it was 0.19 mm in an original document. Non-inferiority margin in previous coronary intervention studies has been set between 0.1 ~ 0.2 mm. The 0.19 mm is not irrelevant in view of the current trial standard.

Received: 11 January 2017 Accepted: 11 January 2017  
Published online: 23 May 2017

## Reference

1. Park S, Yoon C, Oh I, Suh J, Cho Y, Youn T, Choi D, Chae I. Comparison of a drug-eluting balloon first and then bare metal stent with a drug-eluting stent for treatment of de novo lesions: study protocol of a randomized controlled trial. *Trials*. 2013;14:38. doi:10.1186/1745-6215-14-38.

\* Correspondence: ihchae@snu.ac.kr

<sup>†</sup>Equal contributors

Division of Cardiology, Department of Internal Medicine, Seoul National University Bundang Hospital, 82 Gumi-ro, 173 Bein-gil, Bundang-gu, Seongnam-si, Gyeonggi-do 463-707, Republic of Korea

