



저작자표시-비영리-변경금지 2.0 대한민국

이용자는 아래의 조건을 따르는 경우에 한하여 자유롭게

- 이 저작물을 복제, 배포, 전송, 전시, 공연 및 방송할 수 있습니다.

다음과 같은 조건을 따라야 합니다:



저작자표시. 귀하는 원저작자를 표시하여야 합니다.



비영리. 귀하는 이 저작물을 영리 목적으로 이용할 수 없습니다.



변경금지. 귀하는 이 저작물을 개작, 변형 또는 가공할 수 없습니다.

- 귀하는, 이 저작물의 재이용이나 배포의 경우, 이 저작물에 적용된 이용허락조건을 명확하게 나타내어야 합니다.
- 저작권자로부터 별도의 허가를 받으면 이러한 조건들은 적용되지 않습니다.

저작권법에 따른 이용자의 권리는 위의 내용에 의하여 영향을 받지 않습니다.

이것은 [이용허락규약\(Legal Code\)](#)을 이해하기 쉽게 요약한 것입니다.

[Disclaimer](#)

Thesis for the Degree of Master of Landscape Architecture

**Urban heat mitigation by roof surface
materials during the East Asian
summer monsoon**

동아시아 여름 몬순 기간 동안 지붕 표면 재료에
의한 도시 열 완화

February, 2016

**Graduate School of
Seoul National University
Department of Landscape Architecture and Rural Systems
Engineering, Landscape Architecture Major
Lee, Seungjoon**

Urban heat mitigation by roof surface
materials during the East Asian summer
monsoon

지도교수 류 영 렬

이 논문을 조경학 석사 학위논문으로 제출함

2015 년 10 월

서울대학교 대학원

조경지역시스템공학부 생태조경학 전공

이 승 준

이 승 준 의 석사학위논문을 인준함

2015 년 12 월

위 원 장 _____ (인)

부위원장 _____ (인)

위 원 _____ (인)

Abstract

Roof surface materials, such as green and white roofs, have attracted attention in their role in urban heat mitigation, and various studies have assessed the cooling performance of roof surface materials during hot and sunny summer seasons. However, summers in the East Asian monsoon climate region are characterized by significant fluctuations in weather events, such as dry periods, heatwaves, and rainy and cloudy days. This study investigated the efficacy of different roof surface materials for heat mitigation, considering the temperatures both at and beneath the surface of the roof covering materials during a summer monsoon in Seoul, Korea. We performed continuous observations of temperature at and beneath the surface of the roof covering materials, and manual observation of albedo and the normalized difference vegetation index (NDVI) for a white roof, two green roofs (grass [*Poa pratensis*] and sedum [*Sedum sarmentosum*]), and a reference surface. Overall, the surface temperature of the white roof was significantly lower than that of the grass and sedum roofs (1.1 and 1.3°C), whereas the temperature beneath the surface of the white roof did not differ significantly from that of the grass and sedum roofs during the summer. The degree of cloudiness significantly modified the surface temperature of the white roof compared with that of the grass and sedum roofs, which depended on plant metabolisms. It was difficult for the grass to maintain its cooling ability without adequate watering management. After considering the cooling performance and maintenance efforts for different environmental conditions, we concluded that white roof performed better in urban heat mitigation

than grass and sedum during the East Asian summer monsoon. Our findings will be useful in urban heat mitigation in the region.

Keywords : White roof, green roof, urban heat island and East Asian summer monsoon, albedo, NDVI

Student Number : 2014-20048

CONTENTS

| | |
|--|------------|
| ABSTRACT | i |
| CONTENTS | iii |
| LIST OF TABLES..... | v |
| LIST OF FIGURES | v |
| | |
| 1. INTRODCUTION..... | 1 |
| 2. METERIALS AND METHODS..... | 4 |
| 2.1 EXPERIMETNAL SITE AND PERIOD..... | 4 |
| 2.2 ROOF SURFACE MATERIALS..... | 4 |
| 2.3 DATA COLLECTION AND PROCESSING | 5 |
| 2.4 STATISTICAL ANALYSES..... | 7 |
| 3. RESULTS | 9 |
| 3.1 ALBEDOS OF THE ROOF SURFACE MATERIALS..... | 9 |
| 3.2 IMPACT OF CLOUDINESS ON SURFACE TEMPERATURE..... | 10 |
| 3.3 IMPACTS OF DRY PERIOD ON SURFACE TEMPERATURE AND NDVI | 11 |
| 3.4 TEMPERATURE AT AND BENEATH THE SURFACE OF THE ROOF SURFACE MATERIALS DURING SUMMER..... | 13 |
| 4. DISCUSSION..... | 16 |

| | |
|---|----|
| 4.1 HOW DOES CLOUDINESS AFFECT SURFACE TEMPERATURE IN THE ROOF SURFACE MATERIALS?..... | 16 |
| 4.2. WHICH ROOF SURFACE MATERIAL MINIMIZES MAINTENANCE EFFORTS TO SUSTAIN COOLING PERFORMANCE THROUGHOUT SUMMER?..... | 18 |
| 4.3 WHICH ROOF SURFACE MATERIAL IS THE MOST EFFECTIVE IN REDUCING THE TEMPERATURE AT AND BENEATH THE SURFACE? | |

20

| | |
|----------------------------|-----------|
| 5. Conclusion | 23 |
| 6. References | 24 |
| 국문 초록 | 28 |

LIST OF TABLES

Table 1. Albedos of four roof surface materials. Albedos of white, grass, sedum, and reference roofs (mean \pm 95%).

Table 2. The normalized difference vegetation index (NDVI) of grass and sedum roofs. NDVI of grass and sedum for irrigated grass roof (DOY 140), before a heatwave (DOY 181), after a heatwave (DOY 223). Error bars indicate 95% CI.

Table 3. Mean and 95% CI of the temperatures at and beneath the surface of white, grass, and sedum roofs during daytime, nighttime and all day (mean \pm 95% CI).

LIST OF FIGURES

Figure 1. Distribution of daily sky clearness index from June to August (DOY 152–243) between 1973 and 2015. Mean and standard deviation of the sky clearness index for 1973–2015 and 2015.

Figure 2. Diurnal patterns of surface temperature on clear and cloudy days. Diurnal patterns of (a) surface temperature on a clear day (DOY 157), and (b) surface temperature on a cloudy day (DOY 156), for a white, grass, and sedum roofs.

Figure 3. Relationships between sky clearness index and daytime surface temperature (T) differences between a reference and white, grass, and

sedum roofs.

Figure 4. Mean daytime surface temperature of four roof surface materials. Mean daytime surface temperature of four roof surface materials for different periods in summer: (a) DOY 146–152 for irrigated grass roof, (b) DOY 182–188 before a heatwave without irrigation, (c) DOY 221–227 period after a heat wave without irrigation. Error bars indicate 95% CI, letters indicate significant difference among mean daytime surface temperature of the roof surface materials (Tukey test, $p < 0.05$).

Figure 5. Diurnal patterns of the temperature at and beneath the surface of three roof materials. (a) Surface temperature of white roof, grass roof, sedum roof, and reference roof. (b) Temperature beneath the surface of the white roof, grass roof, and sedum roof. Surface temperature of reference roof is include

1. Introduction

The urban heat island effect exists in urban regions worldwide that are home to more than half of the global population (Kalnay and Cai, 2003; Karl *et al.*, 1988; Oke, 1973; Grimm *et al.*, 2008). Given the space restrictions that characterize urban environments, roofs provide optimal spaces that can be secured easily and account for a large proportion of urban surface that can be used to mitigate urban heat through the installation of white roofs (cool or reflective roofs) or green roofs such as grass and sedum that can endure hot and dry conditions in summer (Ban-Weiss *et al.*, 2015). Despite active research in recent years on roof surface materials and heat mitigation (Santamouris, 2014), it remains unclear which roof surface types perform best in heat mitigation under frequently cloudy conditions, dry periods, and heatwaves that characterize summer monsoon climates (Hoag, 2015).

To assess the heat mitigation effects by roof surface materials, it is essential to consider the temperatures both at the surface and beneath the roof covering materials as the former influences air temperature and urban microclimate and the latter the temperature inside the building (Virk *et al.*, 2015). A white-painted thin coating (*i.e.*, a white roof) can be applied to the roof surface. The difference between the temperature at the surface and underneath the coating on white roofs is expected to be marginal. In contrast, grass and sedum roofs form thick layers

composed of planter material, vegetation, and soil layers, which can insulate the transfer of heat from the vegetation surface to the roof surface. Thus, there could be significant differences in the temperatures at the surface and underneath the layers on grass and sedum roofs (Jaffal *et al.*, 2012).

Most previous studies on roof surface materials have focused on either the surface temperature or the temperature beneath the surface, and have not considered both temperatures simultaneously. In general, studies at the building scale have focused on temperature reduction beneath the surface to cool the building interior in summer (Sailor *et al.*, 2011; Kolokotsa *et al.*, 2013). Most studies at the city scale have focused on the surface temperature in roof surface materials that directly influence air temperature and the urban heat island (Li *et al.*, 2014; Susca *et al.*, 2011).

The performance of roof surface materials in heat mitigation under cloudy summer conditions remains unknown. A number of studies on roof surfaces and urban heat mitigation have been carried out in Mediterranean climate regions in the dry season (Akbari *et al.*, 2001; Zinzi and Agnoli, 2012). Most experimental studies have focused on specific short periods during the hottest sunny days in summer (Coutts *et al.*, 2013; Takebayashi and Moriyama, 2007). A modeling study by Georgescu *et al.* (2014) showed that the effectiveness of green and white roofs in urban heat mitigation decreased in humid regions such as Florida in summer. Cloudy and rainy days account for a significant proportion of summer in the East Asian Monsoon climate region and can therefore influence the heat mitigation

effects of roof surface materials.

Minimizing the maintenance efforts of roof surface materials is essential for sustainable roof surface management. Vegetation requires water; thus, previous studies assessed the effects of irrigation and dry periods on the cooling performance of green roofs (Sun *et al.*, 2014; Nagase and Dunnett, 2010). East Asian monsoon summers typically have both heatwaves as well as rainfall, and it is unclear whether vegetation, such as grass and sedum, can survive on roofs and provide cooling effects without watering during summer.

The goal of this study was to investigate which roof surface materials provide the most effective cooling by considering the temperatures at and beneath the surface of the roof covering during the summer monsoon in Seoul. We took field measurements of the temperatures at and beneath the surface for white, sedum, grass, and reference roof surface materials during three months in the summer monsoon. The questions we wished to address included: 1) how does cloudiness affect surface temperature for the three roof surface materials, and 2) which roof surface material minimizes maintenance efforts to sustain the cooling performance throughout summer?

2. Materials and Methods

2.1. Experimental site and period

The experimental site was located on the rooftop of the College of Agriculture and Life Sciences building of Seoul National University, Seoul, South Korea (37.457297°N, 126.948263°E). Data were collected from 20 May to 14 September (Day of Year (DOY): 140–257) in 2015. Summer is defined as the period from June to August (DOY 152–243).

2.2. Roof surface materials

We investigated three roof surface materials; Kentucky bluegrass (*Poa pratensis*; hereinafter called grass), *Sedum sarmentosum* (hereinafter called sedum), and a white-painted surface. The reference roof surface was an existing rooftop of the building that was painted green. On the rooftop, we measured the surface temperature of the reference roof in a 0.5 × 0.5-m plot. The other materials were laid horizontally on the reference roof. We planted Kentucky bluegrass (*Poa pratensis*) in a 1.8 × 1.8 × 0.4-m planter that had been filled with a 0.15-m layer of organic soil on top of a 0.1-m layer of perlite (artificial soil). This soil mix is commonly used in roof gardens (Sun *et al.*, 2014). We used the succulent sedum plant (*Sedum sarmentosum*) in a 1.0 × 1.0 × 0.1-m planter, filled with a 0.05-m layer of organic soil on top of a 0.05-m layer of perlite. High-albedo paint (ESU–

100, NOROO Paint and Coatings Co.) was painted onto the surface of a $1.8 \times 1.2 \times 0.01$ -m aluminum panel.

2.3. Data collection and processing

A hyperspectroradiometer (FieldSpec 4 Wide-Res, ASD, Inc.) was used to measure albedo for four roof surface materials (white, grass, sedum, and reference) on DOY 173 and 257. The same instrument was used to measure the normalized difference vegetation index (NDVI) for grass and sedum on DOY 140, 181, and 223. For each of the four roof surface materials, spectrum measurements were performed through a reference panel (Spectralon, Labsphere) above the target surface and then again immediately after removing the panel; the two spectra were compared to calculate the albedo and reflectance of the surface. NDVI was calculated from the red reflectance (630–690 nm) and near-infrared reflectance (750–800 nm) (Tucker, 1979). Type E thermocouples (CS220, Campbell Scientific) were attached underneath the surfaces of the white, grass, and sedum materials to measure the temperature beneath the surface of each roof surface material during DOY 174–243. Three infrared radiometers (SI-121, Apogee Instrument, Inc.) were installed at 0.8 m above the sedum roof, 0.9 m above the white roof, and 0.5 m above the grass roof to measure surface temperature during DOY 140–243.

There were heatwave alerts (days when a peak daily temperature exceeding 33°C occurred on more than two consecutive days) in Seoul on DOY 218–220 in 2015.

To compare the degree of required management for the different roof surface materials, we specifically analyzed data for three periods that included an irrigated period before the heatwave (DOY 146–152), a non-irrigated period before the heatwave (DOY 182–188), and a non-irrigated period after the heatwave (DOY 221–227). We watered the grass with 30 L per day during the irrigated period.

The daily air temperature, relative humidity, precipitation (for 1953–2015), and solar radiation (from 1973 to August 2015) were obtained from an observatory of the Korea Meteorological Administration (Seoul 37.571406°N, 126.965708°E). To obtain the sky clearness index (the ratio of actual daily total solar radiation at the land surface, R_{sur} , to daily total potential solar radiation at the top of the atmosphere, R_{TOA}), R_{TOA} was calculated as follows:

$$R_{TOA} = \sum_{i=1}^{48} 60 \times 30 \times R_{TOAi} \quad (1)$$

where i is the half hour of the day. R_{TOAi} was calculated as follows (Liu and Jordan, 1960); (Ryu *et al.*, 2012):

$$R_{TOAi} = S_{sc} \times \left[1 + 0.033 \cos \left(\frac{2\pi t_d}{365} \right) \right] \times \cos \beta \quad (2)$$

where S_{sc} is the solar constant (1360 Wm^{-2} ; (Kopp and Lean, 2011), t_d is the day of the year, and β is the solar zenith angle calculated from the latitude, longitude, and local time (Michalsky, 1988). We calculated the daily sky clearness index by dividing R_{sur} recorded at the weather station by R_{TOA} (Eq. 3):

$$\text{Sky clearness index} = \frac{R_{sur}}{R_{TOA}} \quad (3)$$

The distribution of the sky clearness indices for 43 years of summers (1973–2015) is shown in Figure 1.




Figure 1 Distribution of daily sky clearness index from June to August (DOY 152–243) between 1973 and 2015. Mean and standard deviation of the sky clearness index for 1973–2015 and 2015.

2.4. Statistical analyses

Statistical analyses were performed using SigmaPlot 12.0 software (Systat Software Inc., Chicago, IL, USA). We utilized one-way analysis of variance (ANOVA) followed by Tukey’s post-hoc test to compare the mean daytime surface

temperature of each of the four roof surface materials for the three periods. All results are presented as mean \pm 95% CI unless otherwise specified. A *p*-value of < 0.05 was considered to be statistically significant.

3. Results

3.1. Albedos of the roof surface materials

The albedos of the roof surface materials varied significantly (Table 1). The white roof had the highest mean albedo, followed by the reference, grass, and sedum roofs. The albedo of the white roof decreased significantly between DOY 173 (0.819) and 257 (0.748) ($p < 0.05$). The grass and sedum roofs had similar albedos (~ 0.18) on DOY 173, but the albedo of the sedum roof decreased significantly on DOY 257 ($p < 0.05$). The reference surface maintained an albedo (~ 0.3) higher than that of the grass and sedum roofs from DOY 173–257.

Table 1. Albedos of four roof surface materials. Albedos of white, grass, sedum, and reference roofs (mean \pm 95%)

| | White roof | Grass roof | Sedum roof | Reference |
|---------------------------|-------------------|-------------------|-------------------|-------------------|
| DOY 173 (Early Summer) | 0.819 \pm 0.016 | 0.180 \pm 0.016 | 0.174 \pm 0.002 | 0.309 \pm 0.002 |
| DOY 257 (After summer) | 0.748 \pm 0.011 | 0.186 \pm 0.009 | 0.141 \pm 0.005 | 0.315 \pm 0.008 |
| Mean | 0.782 \pm 0.018 | 0.182 \pm 0.010 | 0.160 \pm 0.007 | 0.312 \pm 0.004 |

3.2. Impact of cloudiness on surface temperature

Clouds strongly influenced the diurnal patterns of surface temperature among the roof surface materials (Figure 2). The diurnal patterns of surface temperature of the white, grass, and sedum roofs differed between clear and cloudy days. The peak surface temperature of the white roof was significantly lower than that of the grass (by up to 5°C) and sedum (by up to 13°C) roofs on clear days (Figure 2a). The surface temperature of the white, grass, and sedum roof materials had similar diurnal patterns and values on cloudy days (Figure 2b).




Figure 2. Diurnal patterns of surface temperature on clear and cloudy days. Diurnal patterns of (a) surface temperature on a clear day (DOY 157), and (b) surface temperature on a cloudy day (DOY 156), for a white, grass, and sedum roofs.

The relationships between the sky clearness index and the temperature difference between the roof surface materials and the reference surface during daytime throughout summer differed between the white roof and grass/sedum roofs (Figure 3). The sky clearness index and surface temperature difference of the white roof showed a marked negative correlation ($R = -0.71$). Compared with the white roof, the surface temperature difference between the green roofs and the reference showed weaker correlations with the sky clearness index ($R = -0.30$ for grass, and $R = -0.28$ for sedum).




Figure 3. Relationships between sky clearness index and daytime surface temperature (T) differences between a reference and white, grass, and sedum roofs.

3.3. Impacts of dry period on surface temperature and NDVI

The variations in mean daytime surface temperature and NDVI of the roof surface materials differed for the three summer periods. The mean daytime surface

temperature of the white roof was lowest for all three periods regardless of watering (Figure 4a) and heatwave occurrence (before heat wave: Figure 4b, after heat wave: Figure 4c) ($p < 0.05$). There was no significant difference in the mean daytime surface temperature between the white and irrigated grass roofs ($p > 0.05$; Figure 4a). While the NDVI of the sedum roof was similar in the three periods, the NDVI of the grass roof decreased significantly from 0.643 to 0.554 after the end of watering, and from 0.554 to 0.371 after the heatwave (Table 2). However, there were no significant differences in mean daytime surface temperature between the grass and the sedum roofs in the three periods ($p > 0.05$).




Figure 4. Mean daytime surface temperature of four roof surface materials. Mean daytime surface temperature of four roof surface materials for different periods in summer: (a) DOY 146–152 for irrigated grass roof, (b) DOY 182–188 before a heatwave without irrigation, (c) DOY 221–227 period after a heat wave without irrigation. Error bars indicate 95% CI, letters indicate significant difference among mean daytime surface temperature of the roof surface materials (Tukey test, $p < 0.05$).

Table 2. The normalized difference vegetation index (NDVI) of grass and sedum roofs. NDVI of grass and sedum for irrigated grass roof (DOY 140), before a heatwave (DOY 181), after a heatwave (DOY 223). Error bars indicate 95% CI.

| NDVI | DOY 140 | DOY 181 | DOY 223 |
|-------|---------------|---------------|---------------|
| Grass | 0.643 ± 0.001 | 0.554 ± 0.003 | 0.371 ± 0.055 |
| sedum | 0.686 ± 0.006 | 0.671 ± 0.003 | 0.678 ± 0.007 |

3.4. Temperature at and beneath the surface of the roof surface materials during summer

The mean of the white roof surface temperature was significantly lower than that of the other roofs (1.1°C) over the entire summer period (Table 3). The white, grass, and sedum roofs had lower surface temperatures than the reference during both the daytime and nighttime. The white roof had the lowest peak daytime surface temperature (Figure 5; $p < 0.05$). The mean differences in the surface temperature between the white and green roofs at nighttime (0.1 and 0.8°C for the grass and sedum roofs, respectively) were significantly smaller than the differences during daytime (1.9 and 1.7°C for the grass and sedum roofs, respectively; Table 3). The grass and sedum roofs had similar diurnal patterns in surface temperature.




Figure 5. Diurnal patterns of the temperature at and beneath the surface of three roof materials. (a) Surface temperature of white roof, grass roof, sedum roof, and reference roof. (b) Temperature beneath the surface of the white roof, grass roof, and sedum roof. Surface temperature of reference roof is included.

The diurnal pattern and mean of the temperatures beneath the surface differed from those at the surface among the white, grass, and sedum roofs. The mean daytime temperature beneath the surface of the sedum roof was 0.26°C and 0.8°C lower than that of the white and grass roofs, respectively (Table 3). The temperature beneath the surface of the white roof had a larger peak than that of the green roofs during daytime, but decreased rapidly at night (Figure 5b). The mean temperature beneath the surface of the white roof was lowest at night, whereas that of the sedum roof was lowest during the day ($p < 0.05$). The timings of the peaks of the diurnal patterns of the temperature beneath the surface for the grass and sedum

roofs were delayed compared with the white roof. The grass and sedum roofs had higher temperatures beneath the surface from 00:00 to 06:00.

Table 3. Mean and 95% CI of the temperatures at and beneath the surface of white, grass, and sedum roofs during daytime, nighttime and all day (mean \pm 95% CI).

| Surface temperature | White roof | Grass roof | Sedum roof | Reference | Temperature beneath surface | White roof | Grass roof | Sedum roof |
|----------------------------|----------------|----------------|----------------|----------------|------------------------------------|----------------|----------------|----------------|
| | Daytime | 28.8 \pm 0.3 | 30.7 \pm 0.3 | 30.5 \pm 0.3 | 34.0 \pm 0.3 | Daytime | 28.3 \pm 0.2 | 28.0 \pm 0.1 |
| Nighttime | 25.2 \pm 0.3 | 25.3 \pm 0.3 | 26.0 \pm 0.3 | 27.7 \pm 0.2 | Nighttime | 26.1 \pm 0.2 | 27.8 \pm 0.1 | 27.3 \pm 0.1 |
| All day | 27.3 \pm 0.2 | 28.4 \pm 0.3 | 28.6 \pm 0.3 | 31.4 \pm 0.3 | All day | 27.4 \pm 0.1 | 27.9 \pm 0.1 | 27.1 \pm 0.1 |

4. Discussion

4.1. How does cloudiness affect surface temperature in the roof surface materials?

Our results indicate that the degree of cloudiness can significantly affect the surface temperature of white roofs during the East Asian summer monsoon. Daytime surface temperatures among the three roof surface materials differed significantly between clear and cloudy days. Under clear sky conditions, surface temperature in white roof was lower than that in grass and sedum up to 10°C (e.g. Figure 2a), which is comparable to previous studies (Takebayashi and Moriyama, 2007; Zinzi and Agnoli, 2012). The lowest surface temperature on clear days appeared in the white roof mainly resulted from its high albedo, which reflected the greatest amount of solar radiation (Figure 2a). It has been widely reported that white roofs of high albedo maintain a lower daytime surface temperature on clear-sky days (Akbari *et al.*, 1997; Synnefa *et al.*, 2012; Takebayashi and Moriyama, 2007).

Under cloudy days, we observed that the differences in daytime surface temperature among the three roof surface materials were lower (Figure 2b). This tendency is in agreement with a previous modeling study for London, which showed that the difference in the daytime air temperature between a white and

green roof decreased on cloudy days (Virk *et al.*, 2015). The modeling study of Georgescu *et al.* (2014) showed that the difference in the air temperature between white and green roofs in California will be greater than the difference in the air temperature among these roof materials in Florida and the Mid-Atlantic in 2100. Yang *et al.*, (2015) reported that cooling performance of white roof depends on meteorological and geographical conditions. Although the surface cooling performance by white roof was largely controlled by cloudiness, it showed the lowest mean surface temperature among the four surface materials over the whole summer period (Table 3).

The difference in the surface temperature between the white roof and the reference surface showed a stronger negative correlation ($R = - 0.71$) than that between the grass ($R = - 0.30$) and sedum ($R = - 0.28$) roofs and the reference surface. This result indicates that the surface temperature of green roofs was controlled by not only cloudiness but also physiological mechanisms such as stomata closure and transpiration (Figure 3). Thus, the cooling performance of grass roofs for diverse sky conditions is highly variable given the extreme weather events such as dry period, heat spell and rainfall.

The white roofs showed consistently lower surface temperatures than the reference for the range of sky clearness index of the 40-year mean ± 1 S.D. (0.172–0.508; Figure 3a). Thus, we expect that the installed white roof could be cooler than the reference during summer in Seoul for most sky conditions. The annual

mean surface solar radiation in Korea increased over the 15-year period from 1990 to 2005 (Wild *et al.*, 2009), and the sky clearness index in Seoul has also significantly increased over the 25 years from 1990 to 2014 (0.002 year^{-1} , $p < 0.05$ from Seoul weather station). Thus, the cooling effects by white roof in this region are likely to increase in future.

4.2. Which roof surface material minimizes maintenance efforts to sustain cooling performance throughout summer?

The white roof required minimal maintenance to sustain cooling performance. The daytime surface temperature among the three roof surface materials differed for the three periods (watering, and before and after the heatwave). There were larger variations in surface temperature for the same sky clearness index for the green roofs (Figure 3). This indicates that the green roofs were influenced by biological mechanisms, such as stomata conductance and transpiration. While there was no significant difference in the surface temperature between the transpiring grass roof and the white roof during the irrigated period (Figure 4a), during the non-irrigated periods, the surface temperature of the grass roof was significantly higher than that of the white roof. The effect of the green roofs in mitigating heat is small without a thick soil layer and watering (Sun *et al.*, 2014; Coutts *et al.*, 2013; Li *et al.*, 2014; Thuring *et al.*, 2010), because grass has a shallow rooting depth; thus, its metabolic

activity is highly sensitive to the level of soil moisture (Ryu *et al.*, 2008; Ryu *et al.*, 2010). Although there was little change in the NDVI of the sedum roof over the three periods, and a significant decrease in the NDVI of the grass roof between DOY 181 and 223, there was no significant difference in the surface temperature between the sedum and grass roofs on these dates (Figure 4 and Table 2). This indicates that the sedum tolerated the hot summer conditions well, which was reflected in the stable NDVI, but it did not decrease the daytime surface temperature effectively (Nagase and Dunnett, 2010). A previous study showed that a sedum roof that received sufficient water could significantly reduce the sensible heat flux (Coutts *et al.*, 2013) thus surface temperature, which differed from the findings in our study. The 2015 summer monsoon brought 395.5 mm of rainfall between June and August (Seoul weather station, Korea Meteorological Administration), which was the lowest recorded summer rainfall since 1954. We found that this amount of rainfall was insufficient to support a covering of healthy grass, which rapidly reached senescence in early August, as shown by the decrease in the NDVI (Table 2). Furthermore, the vapor pressure deficit, an indicator for plant water stress, in Seoul has increased over the last 61 years (1954–2014; 56.7 Pa decade⁻¹, $p < 0.05$ from Seoul weather station). Thus, we expect the management of green roofs in this region to become more difficult in the future.

The accumulated dust on the white roof surface slightly decreased the albedo. A previous study reported that accumulated dust on a white roof surface reduced the surface albedo, which resulted in the decrease in the surface temperature (Gaffin *et*

al., 2012). We measured the albedo of the white roof on DOY 173 and 257 and found a significant decrease (from 0.819 to 0.748) during this period (Table 1). To maintain consistent cooling performance of white roofs in the megacity Seoul, which experiences significant air pollution, periodic cleaning of the white roof surface would be required, which is likely easier and more cost-effective than the watering required by green roofs.

4.3. Which roof surface material is the most effective in reducing the temperature at and beneath the surface?

The white roof showed the most effective surface cooling for both daytime and nighttime throughout the summer. Our results suggest that the performance of the roof surface materials depends on the characteristics of the weather. Although the difference in the surface temperature between the green roofs and the white roof was small on cloudy days, the surface temperature of the white roof was clearly lowest on clear days (Figure 2). The white roof had the lowest daytime surface temperature, which is consistent with previous studies (Takebayashi and Moriyama, 2007; Coutts *et al.*, 2013; Georgescu *et al.*, 2014). The white roof exhibited the lowest mean surface temperature among the roof surfaces throughout the entire summer period (Table 3).

The peak daytime temperature beneath the surface among the three roof surface materials differed from the peak daytime surface temperature (Figure 5). The white

roof had the highest daytime temperature beneath the surface (Figure 5 and Table 3). The heat capacity of the surface materials plays an important role in the variations in the temperature beneath the surface. Because the white roof consisted of a single aluminum panel that had been painted, the temperature difference between the surface and beneath the surface was smaller ($0.7 \pm 0.5^{\circ}\text{C}$) than that of the green roofs ($2.7 \pm 0.8^{\circ}\text{C}$ and $3.5 \pm 1^{\circ}\text{C}$ for the grass and sedum roofs, respectively), which were composed of soil layers. The soil layers created an insulating effect during the daytime and released their stored heat at night (Jim, 2015). This caused the temperature beneath the surface of the green roofs to be higher than that of the white and reference roofs at night. These diurnal patterns in the temperature beneath the surface were also observed for white and sedum roofs in a previous study (Coutts, Daly et al. 2013).

Overall, white roof is the most effective strategy for urban heat mitigation during the East Asian monsoon summer. Although the daytime temperature beneath the surface of the white roof was higher than that of the sedum roof (Table 3), the mean difference in the surface temperature between the white and sedum roofs ($1.3 \pm 0.4^{\circ}\text{C}$) was greater than the mean difference in the temperature beneath the surface between the white and sedum roofs ($0.3 \pm 0.3^{\circ}\text{C}$) on all days. Previous studies in southern Europe where low-rise buildings dominate tended to focus on reducing the temperature beneath the roof surface to decrease the energy use of the top floor of the building (Synnefa *et al.*, 2012; Romeo and Zinzi, 2013). In the case of Seoul,

a large proportion of the city is composed of high-rise buildings and apartments. Thus, reducing the roof surface temperature is likely more effective for urban heat mitigation than reducing the heat transfer from beneath the roof surface to the top floor of the building.

5. Conclusion

As the size and number of cities continue to increase, the urban heat island effect will increase accordingly, and the demand for roofs with cooling properties is likely to rise. Although the East Asian region accounts for approximately 20 percent of the world's population, there have been few studies assessing roof types that mitigate urban heat in a monsoon climate region. We found that white roof showed the lowest mean surface temperature (27.3°C) over the whole summer compared to grass (28.4°C), sedum (28.6°C) and reference (31.4°C) roofs although cloudiness strongly regulated surface temperature in white roof. White roof also required less effort to maintain its cooling performance than the green roofs, which required adequate watering. Thus, we suggest that white roof is more effective in heat mitigation than grass, and sedum during summer in Seoul. The findings of this study could contribute to the development of effective strategies for mitigating urban heat and provide information for city planners and policy makers in selecting more effective roof surfaces for cooling.

6. References

- Akbari H, Bretz S, Kurn D M and Hanford J 1997 Peak power and cooling energy savings of high-albedo roofs *Energy and Buildings* **25** 117-26
- Akbari H, Pomerantz M and Taha H 2001 Cool surfaces and shade trees to reduce energy use and improve air quality in urban areas *Solar Energy* **70** 295-310
- Ban-Weiss G A, Woods J, Millstein D and Levinson R 2015 Using remote sensing to quantify albedo of roofs in seven California cities, Part 2: Results and application to climate modeling *Solar Energy* **115** 791-805
- Coutts A M, Daly E, Beringer J and Tapper N J 2013 Assessing practical measures to reduce urban heat: Green and cool roofs *Building and Environment* **70** 266-76
- Gaffin S, Imhoff M, Rosenzweig C, Khanbilvardi R, Pasqualini A, Kong A Y, Grillo D, Freed A, Hillel D and Hartung E 2012 Bright is the new black—multi-year performance of high-albedo roofs in an urban climate *Environmental Research Letters* **7** doi:10.1088/1748-9326/7/1/014029
- Georgescu M, Morefield P E, Bierwagen B G and Weaver C P 2014 Urban adaptation can roll back warming of emerging megapolitan regions *Proceedings of the National Academy of Sciences of the United States of America* **111** 2909-14
- Grimm N B, Faeth S H, Golubiewski N E, Redman C L, Wu J, Bai X and Briggs J M 2008 Global change and the ecology of cities *science* **319** 756-60
- Hoag H 2015 How cities can beat the heat *Nature* **524** 402-4
- Jaffal I, Ouldboukhitine S E and Belarbi R 2012 A comprehensive study of the impact of green roofs on building energy performance *Renewable Energy* **43** 157-64
- Jim C Y 2015 Diurnal and partitioned heat-flux patterns of coupled green-building

- roof systems *Renewable Energy* **81** 262-74
- Kalnay E and Cai M 2003 Impact of urbanization and land-use change on climate *Nature* **423** 528-31
- Karl T R, Diaz H F and Kukla G 1988 Urbanization: Its Detection and Effect in the United States Climate Record. In: *Journal of Climate*, pp 1099-123
- Kolokotsa D, Santamouris M and Zerefos S C 2013 Green and cool roofs' urban heat island mitigation potential in European climates for office buildings under free floating conditions *Solar Energy* **95** 118-30
- Kopp G and Lean J L 2011 A new, lower value of total solar irradiance: Evidence and climate significance *Geophysical Research Letters* **38**, L01706 doi:10.1029/2010GL045777
- Li D, Bou-Zeid E and Oppenheimer M 2014 The effectiveness of cool and green roofs as urban heat island mitigation strategies *Environmental Research Letters* **9** doi:10.1088/1748-9326/9/5/055002
- Liu B Y and Jordan R C 1960 The interrelationship and characteristic distribution of direct, diffuse and total solar radiation *Solar energy* **4** 1-19
- Michalsky J J 1988 The astronomical almanac's algorithm for approximate solar position (1950–2050) *Solar Energy* **40** 227-35
- Nagase A and Dunnett N 2010 Drought tolerance in different vegetation types for extensive green roofs: Effects of watering and diversity *Landscape and Urban Planning* **97** 318-27
- Oke T R 1973 City size and the urban heat island. In: *Atmospheric Environment (1967)*, pp 769-79
- Romeo C and Zinzi M 2013 Impact of a cool roof application on the energy and comfort performance in an existing non-residential building. A Sicilian case study *Energy and Buildings* **67** 647-57
- Ryu Y, Baldocchi D D, Black T A, Detto M, Law B E, Leuning R, Miyata A, Reichstein M, Vargas R and Ammann C 2012 On the temporal upscaling of evapotranspiration from instantaneous remote sensing measurements to 8-

- day mean daily-sums *Agricultural and Forest Meteorology* **152** 212-22
- Ryu Y, Baldocchi D D, Ma S and Hehn T 2008 Interannual variability of evapotranspiration and energy exchange over an annual grassland in California *Journal of Geophysical Research* **113**, **D09104**, doi:10.1029/2007JD009263
- Ryu Y, Baldocchi D D, Verfaillie J, Ma S, Falk M, Ruiz-Mercado I, Hehn T and Sonnentag O 2010 Testing the performance of a novel spectral reflectance sensor, built with light emitting diodes (LEDs), to monitor ecosystem metabolism, structure and function *Agricultural and Forest Meteorology* **150** 1597-606
- Sailor D J, Elley T B and Gibson M 2011 Exploring the building energy impacts of green roof design decisions—a modeling study of buildings in four distinct climates *Journal of Building Physics* **0(0)** doi: 10.1177/1744259111420076
- Santamouris M 2014 Cooling the cities – A review of reflective and green roof mitigation technologies to fight heat island and improve comfort in urban environments *Solar Energy* **103** 682-703
- Sun T, Bou-Zeid E and Ni G-H 2014 To irrigate or not to irrigate: Analysis of green roof performance via a vertically-resolved hygrothermal model *Building and Environment* **73** 127-37
- Susca T, Gaffin S R and Dell’Osso G R 2011 Positive effects of vegetation: Urban heat island and green roofs *Environmental Pollution* **159** 2119-26
- Synnefa A, Saliari M and Santamouris M 2012 Experimental and numerical assessment of the impact of increased roof reflectance on a school building in Athens *Energy and Buildings* **55** 7-15
- Takebayashi H and Moriyama M 2007 Surface heat budget on green roof and high reflection roof for mitigation of urban heat island *Building and Environment* **42** 2971-9
- Thuring C E, Berghage R D and Beattie D J 2010 Green roof plant responses to different substrate types and depths under various drought conditions

HortTechnology **20** 395-401

- Tucker C J 1979 Red and photographic infrared linear combinations for monitoring vegetation *Remote Sensing of Environment* **8** 127-50
- Virk G, Jansz A, Mavrogianni A, Mylona A, Stocker J and Davies M 2015 Microclimatic effects of green and cool roofs in London and their impacts on energy use for a typical office building *Energy and Buildings* **88** 214-28
- Wild M, Trüssel B, Ohmura A, Long C N, König-Langlo G, Dutton E G and Tsvetkov A 2009 Global dimming and brightening: An update beyond 2000 *Journal of Geophysical Research* **114**, **D00D13**, doi:10.1029/2008JD011382
- Yang J, Wang Z-H and Kaloush K E 2015 Environmental impacts of reflective materials: Is high albedo a 'silver bullet' for mitigating urban heat island? *Renewable and Sustainable Energy Reviews* **47** 830-43
- Zinzi M and Agnoli S 2012 Cool and green roofs. An energy and comfort comparison between passive cooling and mitigation urban heat island techniques for residential buildings in the Mediterranean region *Energy and Buildings* **55** 66-76

국문 초록

화이트 지붕과 옥상 녹화와 같은 지붕 표면 재료들은 도시 열 저감 목적으로 큰 관심을 받아 왔고 다양한 선행 연구들은 주로 매우 고온 건조한 여름 기간 동안 지붕 표면 재료들의 쿨링 성능에 대해 평가했다. 그러나 동아시아 몬순 기후 지역의 여름은 가뭄, 폭염, 장마와 흐린 날과 같이 다양한 날씨가 혼재 되어 있다. 본 연구는 서울에서 여름 동안 여러 가지 지붕 표면 재료들에 대해 밀면과 표면 온도를 모두 고려 하여 열 저감 효과 차이를 비교 하였다. 연구에서는 화이트 지붕, 두 가지 옥상 녹화 (잔디 (*Poa pratensis*) 와 세덤 (*Sedum sarmentosum*)) 표면과 기존 옥상 표면에 대해 밀면 온도와 표면 온도를 연속으로 관측하고, 알베도 (albedo)와 정규식생지수(normalized difference vegetation index, NDVI)를 수동으로 관측 하였다. 종합적으로 여름 동안, 화이트 지붕의 표면 온도는 잔디와 세덤 표면 온도 보다 통계적으로 낮았지만 (1.1°C and 1.3°C), 화이트 지붕의 밀면 온도는 잔디와 세덤 밀면 온도와 통계적인 차이를 보이지 않았다. 식물 대사에 달려있는 잔디와 세덤 표면과 비교할 때, 운량의 정도는 화이트 지붕의 표면 온도를 통계적으로 바꾸었다. 잔디는 충분한 관수 관리 없이 쿨링 능력을 유지 하기 어려웠

다. 여러 다른 환경적 상황에서 쿨링 성능과 관리 노력을 모두 고려 하였을 때, 화이트 지붕은 동아시아 여름 몬순 기간 동안 잔디와 세덤 표면 보다 도시 열 저감에 더 효과적이라는 결론을 내렸다. 연구 결과는 동아시아 몬순 지역에서 열 저감에 도움을 줄 것 이다.

주요어 : 화이트 지붕, 옥상녹화, 도시열섬, 동아시아 여름 몬순, 알베도, 정규식생지수

학번 : 2014-20048