


RECEIVED: July 13, 2023

ACCEPTED: November 13, 2023

PUBLISHED: November 24, 2023

Search for Z' bosons decaying to pairs of heavy Majorana neutrinos in proton-proton collisions at $\sqrt{s} = 13 \text{ TeV}$



The CMS collaboration

E-mail: cms-publication-committee-chair@cern.ch

ABSTRACT: A search for the production of pairs of heavy Majorana neutrinos (N_ℓ) from the decays of Z' bosons is performed using the CMS detector at the LHC. The data were collected in proton-proton collisions at a center-of-mass energy of $\sqrt{s} = 13 \text{ TeV}$, with an integrated luminosity of 138 fb^{-1} . The signature for the search is an excess in the invariant mass distribution of the final-state objects, two same-flavor leptons (e or μ) and at least two jets. No significant excess of events beyond the expected background is observed. Upper limits at 95% confidence level are set on the product of the Z' production cross section and its branching fraction to a pair of N_ℓ , as functions of N_ℓ and Z' boson masses (m_{N_ℓ} and $m_{Z'}$, respectively) for $m_{Z'}$ from 0.4 to 4.6 TeV and m_{N_ℓ} from 0.1 TeV to $m_{Z'}/2$. In the theoretical framework of a left-right symmetric model, exclusion bounds in the m_{N_ℓ} - $m_{Z'}$ plane are presented in both the electron and muon channels. The observed upper limit on $m_{Z'}$ reaches up to 4.42 TeV. These are the most restrictive limits to date on the mass of N_ℓ as a function of the Z' boson mass.

KEYWORDS: Beyond Standard Model, Hadron-Hadron Scattering

ARXIV EPRINT: [2307.06959](https://arxiv.org/abs/2307.06959)

Contents

1	Introduction	1
2	The CMS detector	3
3	Simulated samples	3
4	Event reconstruction and object identification	4
4.1	Lepton selection	6
4.2	Jet selection	6
5	Event selections	7
6	Background estimation	8
7	Systematic uncertainties and fitting procedure	9
8	Results and discussion	13
9	Summary	16
	The CMS collaboration	23

1 Introduction

The observation of neutrino oscillations [1–4] established that at least two of the standard model (SM) neutrinos have mass. The nonzero masses of the neutrinos are clear evidence of physics beyond the SM, in which neutrinos are massless. Upper limits on the neutrino masses have been obtained from cosmological observations [5]. A direct measurement from tritium beta decays [6] indicates that the electron-type neutrino mass is less than 0.8 eV. The fact that neutrino masses are so much smaller than the other fermion masses and that right-handed neutrinos are not observed suggest that neutrino masses may have an origin other than a Yukawa coupling to the Higgs field.

In addition to its inability to explain neutrino mass, the SM does not provide any clear answer as to what is the source of the parity violation in the weak sector or of the matter dominance of the universe. One of the leading theoretical solutions is to introduce a left-right symmetry model (LRSM) [7]. The LRSM is based on a gauge group of $SU(3)_C \otimes SU(2)_L \otimes SU(2)_R \otimes U(1)_{B-L}$, established by a symmetry between the left- and right-handed $SU(2)$ groups. In this model, there are three additional gauge bosons, W_R^\pm and Z' , as well as three right-handed neutrinos (N_e , N_μ , and N_τ). The spontaneous symmetry breaking of the LRSM, at some energy scale, can generate the gauge group of




Figure 1. Feynman diagram representing the pair production of N_ℓ via Z' boson exchange, where $\ell = e$ or μ . Each N_ℓ decays into a lepton and two quarks, and we assume that the W_R^\pm is heavier than the N_ℓ . Both opposite- and same-sign dileptons from the decays of the heavy neutrino pair are allowed, because of the Majorana nature of N_ℓ .

the SM in a way that naturally includes the observed parity violation in the weak sector, and also provides an additional source of CP violation that can explain the asymmetry between matter and antimatter in the universe. Heavy right-handed neutrinos and light left-handed neutrinos are also naturally generated by the symmetry breaking, through a process referred to as the see-saw mechanism [8–12].

As the LRSM provides explanations for these basic observations of beyond the SM physics, it is compelling to probe signatures of this model. Although there have been previous searches for both new heavy gauge bosons and heavy right-handed neutrinos [13–17], the unique signature of the LRSM is the presence of events that have an extra gauge boson as well as heavy right-handed neutrinos. In this analysis, we search for the pair production of heavy right-handed neutrinos through an extra neutral gauge boson (Z'). The resonance production of a Z' boson and its decay into heavy neutrinos are shown in figure 1. In the benchmark model used in this study, the heavy right-handed neutrino is a Majorana particle and decays into a lepton plus two quarks. We therefore select events with two electrons or two muons, and at least two reconstructed jets, to search for an excess in the invariant mass distribution of them. We use opposite- and same-sign leptons inclusively, and mixing between heavy neutrinos of electron type and muon type is not considered. Rich scalar sectors are predicted by LRSM models in general. In this search, results are presented in terms of the product of the cross section and branching fraction of the process, thus no assumptions are made concerning the nature of the scalar sector. Detailed underlying assumptions for this LRSM will be discussed in section 3.

Searches for a W_R^\pm boson and its decay to a heavy Majorana neutrino (N_ℓ) have been performed in the framework of the LRSM by the ATLAS and the CMS Collaborations at the LHC [18–28]. The most stringent lower limit on the W_R^\pm mass from the ATLAS Col-

laboration is about 6.4 TeV [28], which was obtained using a data set of proton-proton (pp) collisions at a center-of-mass energy of 13 TeV, corresponding to an integrated luminosity of 139 fb^{-1} . A search for an extra gauge boson Z' and its decay to a pair of N_ℓ has also been performed by the ATLAS Collaboration, which obtained a lower limit on the Z' mass ($m_{Z'}$) of 2.2 TeV [21] using a data sample corresponding to 20.3 fb^{-1} at $\sqrt{s} = 8 \text{ TeV}$.

All these results are limited by a reduced efficiency for the signals in the kinematic region where the heavy neutrino mass (m_{N_ℓ}) is much smaller than the mass of the extra gauge boson. A lepton from the decay of a highly Lorentz-boosted heavy neutrino overlaps with jets from the same decay, and lepton identification requires the use of isolation criteria. As a result, the signal selection efficiency is reduced in this kinematic region. It is important to recover the efficiency in this kinematic region, because the dominant decay of extra gauge bosons is to heavy neutrinos. Here we present a method that is sensitive to a broad range of N_ℓ masses, including the region where m_{N_ℓ} is much smaller than $m_{Z'}$. The analysis presented here uses a pp collision data set at $\sqrt{s} = 13 \text{ TeV}$ collected with the CMS detector, corresponding to an integrated luminosity of 138 fb^{-1} .

Tabulated results are provided in the HEPData record for this analysis [29].

2 The CMS detector

The central feature of the CMS apparatus is a superconducting solenoid of 6 m internal diameter, providing a magnetic field of 3.8 T. Within the solenoid volume are a silicon pixel and strip tracker, a lead tungstate crystal electromagnetic calorimeter (ECAL), and a brass and scintillator hadron calorimeter (HCAL), each composed of a barrel and two endcap sections. Forward calorimeters extend the pseudorapidity (η) coverage provided by the barrel and endcap detectors. Muons are detected in gas-ionization detectors embedded in the steel flux-return yoke outside the solenoid. A more detailed description of the CMS detector, together with a definition of the coordinate system used and the relevant kinematic variables, can be found in ref. [30].

3 Simulated samples

Signal and background events for each data-taking year from 2016 to 2018 are produced using Monte Carlo (MC) event generators and a detailed simulation of the CMS detector. The expected background processes include Drell-Yan (DY) dilepton plus jets production, and production of $t\bar{t}$, dibosons, $W + \text{jets}$, single top quarks, and QCD multijet events. The DY events are simulated with `MADGRAPH5_aMC@NLO 2.2.2` [31] with up to four jets at leading order (LO) in quantum chromodynamics (QCD) for 2016. For 2017 and 2018, `MADGRAPH5_aMC@NLO 2.4.2` is used. The W events associated with jets are obtained with the `MADGRAPH5_aMC@NLO 2.3.3` [31] generator with up to four jets at LO. The MLM scheme [32] is used to match partons from matrix element calculations and parton showers for these two samples. The $t\bar{t}$ production is simulated with the `POWHEG v2` [33–36] generator at next-to-LO (NLO). Diboson (WW , WZ , and ZZ) events are simulated with the `PYTHIA 8.212` [37] generator at LO. The t -channel and tW single top quark

events are produced with the POWHEG v1 [33–35, 38, 39] generator. Single top quark production in the s -channel is simulated with the MADGRAPH5_aMC@NLO generator at NLO precision. The $t\bar{t}$ production in association with a W (Z) boson is simulated using MADGRAPH5_aMC@NLO at NLO (LO) precision. QCD multijet events are produced using MADGRAPH5_aMC@NLO at LO precision.

The NNPDF3.0 [40] parton distribution function (PDF) set is used to simulate background events corresponding to the data recorded in 2016. For the 2017 and 2018 backgrounds and all signal events, the NNPDF3.1 [41] PDF set is used. Parton showering, including photon radiation and hadronization processes, with the choice of NNPDF3.0 (or 3.1) are modeled using PYTHIA 8.226 (8.230) [37] with the CUETP8M1 (CP5) [42, 43] tune. Finally, the response of the CMS detector is modeled using GEANT4 [44] for all simulated samples.

The signal events for the process shown in figure 1 are simulated using MADGRAPH5_aMC@NLO 2.6.0 at NLO precision with the LRSM model cards [45]. Heavy neutrinos with electron and muon flavors are considered, and no mixing between the two flavors is assumed. Both opposite- and same-sign configurations of dileptons are taken into account in the simulation. The signal samples are simulated for various $m_{Z'}$ and m_{N_ℓ} hypotheses, with $m_{Z'}$ between 400 and 5000 GeV, and m_{N_ℓ} between 100 GeV and $m_{Z'}/2$. In the simulation, the W_R mass is assumed to be 5 TeV, so that the heavy neutrinos are always lighter than the W_R boson. The product of the NLO cross section and the branching fraction (BF) is the same for both dilepton channels and is 27.77 pb for $m_{Z'} = 400$ GeV, and 8.60×10^{-5} pb for $m_{Z'} = 5000$ GeV with $m_{N_\ell} = 100$ GeV. With $m_{N_\ell} = 2400$ GeV, it is 4.94×10^{-7} pb for $m_{Z'} = 5000$ GeV. A product of cross section and branching fraction of 1 fb corresponds typically to the Z' boson masses in the range 3 to 4 TeV, depending slightly on m_{N_ℓ} .

Minimum bias events generated with PYTHIA are superimposed on each simulated hard scattering event to reproduce the effect of extra pp interactions within the same or neighboring bunch crossings (pileup). The simulated events are weighted such that the distribution of the number of additional pileup interactions, estimated from the measured instantaneous luminosity for each bunch crossing, matches that observed in data sets individually for each year of data-taking in 2016–2018. The simulated events are processed with the same reconstruction software used for pp collision data.

4 Event reconstruction and object identification

Candidate events from decays of heavy neutrino pairs are selected using a two-tiered trigger system [46]. The first level (L1), composed of custom hardware processors, uses information from the calorimeters and muon detectors to select events at a rate of around 100 kHz within a fixed time interval of less than $4\mu\text{s}$. The second level, known as the high-level trigger (HLT), consists of a farm of processors running a version of the full event reconstruction software optimized for fast processing, and reduces the event rate to around 1 kHz before data storage.

For dielectron channel events, a diphoton trigger is used, which requires two electron or photon (e/γ) objects detected in the ECAL with a minimum transverse energy (E_T) of 60 GeV for 2016 data or 70 GeV for 2017 and 2018 data. Dimuon channel events must pass a single-muon trigger that requires the presence of a muon with a minimum p_T of 50 GeV.

In the dielectron channel, the trigger efficiency of signal events is about 85%, dropping to about 10% in the kinematic region where m_{N_ℓ} is much smaller than $m_{Z'}$. The reduction of efficiency is mainly caused by the requirement on H/E, the ratio between energies deposited in HCAL and ECAL, being smaller than 0.15 (barrel) or 0.10 (endcap). Since the requirement on H/E is essential for distinguishing electrons and photons from hadrons, there are inevitable signal losses in the dielectron channel for highly Lorentz-boosted events. This analysis uses a double-photon trigger, since it has the loosest H/E requirement among all e/γ HLT paths.

The trigger efficiency for signal events in the dimuon channel depends on m_{N_ℓ} and $m_{Z'}$, and ranges between 80 and 95%, except at the lightest Z' boson mass points, where it is somewhat lower because of the high momentum threshold of the muon trigger. For example an efficiency of about 60% is observed for the mass points at the 400 GeV and 600 GeV. The cross sections for these mass points are higher than those for heavier Z' bosons. This allows good sensitivity to be maintained while keeping the same trigger for all masses.

The primary vertex (PV) is taken to be the vertex corresponding to the hardest scattering in the event, evaluated using tracking information alone, as described in section 9.4.1 of ref. [47].

Particles in selected events are reconstructed and identified using the global event reconstruction (also called particle-flow event reconstruction [48]) with an optimized combination of all subdetector information. The identification of the particle type (photon, electron, muon, charged hadron, neutral hadron) plays an important role in the determination of the particle direction and energy. Photons (e.g., coming from π^0 decays or from electron bremsstrahlung) are identified as ECAL energy clusters not linked to the extrapolation of any charged particle trajectory to the ECAL. Electrons (e.g., from Z boson leptonic decays) are identified as a primary charged particle track and potentially many ECAL energy clusters corresponding to this track extrapolation to the ECAL and to possible bremsstrahlung photons emitted while traversing the tracker material. Muons (e.g., from W boson leptonic decays) are identified as tracks in the central tracker matching with either a track or several hits in the muon system, and associated with calorimeter deposits compatible with the muon hypothesis. Charged hadrons are identified as charged particle tracks neither identified as electrons, nor as muons. Finally, neutral hadrons are identified as HCAL energy clusters not linked to any charged hadron trajectory, or as a combined ECAL and HCAL energy excess with respect to the expected charged hadron energy deposit.

The energy of photons is obtained from the ECAL measurement. The energy of electrons is determined from a combination of the track momentum at the PV, the corresponding ECAL cluster energy, and the energy sum of all bremsstrahlung photons associated with the track. The energy of muons is obtained from the corresponding track momentum. The energy of charged hadrons is determined from a combination of the track momentum and

the corresponding ECAL and HCAL energies, corrected for the response function of the calorimeters to hadronic showers. Finally, the energy of neutral hadrons is obtained from the corresponding corrected ECAL and HCAL energies.

4.1 Lepton selection

Electrons are selected in the region of $|\eta_C| < 2.5$, where η_C is the pseudorapidity of the ECAL cluster with respect to the nominal center of the CMS detector. The transition region between ECAL barrel and endcaps, $1.44 < |\eta_C| < 1.57$, is excluded. The electron E_T is required to be greater than 65 (75) GeV for the 2016 (2017 and 2018) data. In this analysis, two electron identification criteria are used: “loose” and “tight”. A “loose” electron is an object passing the “HEEP” selection or passing the selection of a loose cut-based electron with no isolation criteria [49, 50]. The “HEEP” criteria have been developed for the analysis of high-energy electrons. A “tight” electron is an object passing the “HEEP” requirement.

In addition, a “veto” electron is defined using the same criteria as the loose electron but lowering the E_T requirement to 10 GeV.

Muons are selected in the region $|\eta| < 2.4$ with $p_T > 65$ (75) GeV for 2016 (2017 and 2018). Higher p_T requirements for muons compared to the trigger thresholds are applied to reduce QCD background. A dedicated muon identification algorithm, HighPtID [51], adapted from searches for resonances using high-momentum leptons [52] is used for this analysis. Two muon identification criteria are used, “loose” and “tight”. A “loose” muon is required to satisfy the HighPtID requirements. A “tight” muon must pass both the HighPtID and isolation requirements. The isolation is defined as the p_T sum of tracks within a cone of radius $\Delta R = \sqrt{(\Delta\eta)^2 + (\Delta\phi)^2} = 0.3$ around the muon candidate direction, where ϕ is its azimuthal angle in radians. The momentum of the muon candidate is excluded from the sum. If the sum is less than 10% of the muon candidate’s p_T , it passes the isolation requirement. In addition, a “veto” muon is defined using the particle-flow muons with $p_T > 10$ GeV.

4.2 Jet selection

For each event, hadronic jets are reconstructed by clustering particle-flow objects using the anti- k_T algorithm [53, 54] with two distance parameters of 0.4 and 0.8, producing AK4 and AK8 jets, respectively. The jet momentum is determined as the vectorial sum of all particle momenta in the jet, and is found from the simulation to be within 5 to 10% of the true momentum, on average, over the entire p_T spectrum and detector acceptance. Pileup interactions can contribute extra tracks and calorimetric energy depositions to the jet momentum.

In order to mitigate the effect of pileup, the pileup-per-particle identification algorithm [55, 56] is used at the reconstructed-particle level, making use of local shape information, event pileup properties and tracking information. Charged particles identified to be originating from pileup vertices are discarded. For each neutral particle, a local shape variable is computed using the surrounding charged particles compatible with the PV within the tracker acceptance ($|\eta| < 2.5$), and using both charged and neutral particles in the region outside of the tracker coverage. The momenta of the neutral particles are then rescaled according to their probability to originate from the PV deduced from the

local shape variables, superseding the need for jet-based pileup corrections [57]. For this analysis, AK4 (AK8) jets are required to have $p_T > 40$ (300) GeV and $|\eta| < 2.7$.

AK8 jets are groomed using the soft-drop algorithm [58, 59]. In this algorithm, the constituents of the AK8 jets are reclustered using the Cambridge-Aachen algorithm [60, 61]. Soft radiation and wide-angle radiation from the jet are removed by setting the angular exponent β to zero, the soft cutoff threshold to 0.1, and the characteristic radius R_0 to 0.8 [62]. All AK8 jets are required to have mass greater than 40 GeV to reduce the number of AK8 jets that originate from a single parton.

In order to avoid double counting of AK4 jets with a lepton or an AK8 jet, any AK4 jet located within $\Delta R < 0.4$ of a loose lepton or within $\Delta R < 1.0$ of an AK8 jet is not used in this analysis.

5 Event selections

The signature targeted in this search is two same-flavor leptons and four jets from heavy neutrino decays, as shown in figure 1. The kinematic distributions of the final state are strongly dependent on the ratio of twice the heavy neutrino mass and the Z' boson mass ($2m_{N_\ell}/m_{Z'}$). If the ratio is much smaller than one, the lepton and two jets from a heavy neutrino decay are merged into one jet because of the high Lorentz boost provided by the heavy neutrino. As a consequence, leptons are mostly reconstructed as loose leptons since they seldom satisfy the isolation criteria. In contrast, leptons and jets are well separated if the ratio is close to unity. A dedicated strategy is developed to achieve a good signal selection efficiency in the wide kinematic region covered by this search.

The selected events are required to have exactly two same-flavor (ee or $\mu\mu$) loose leptons using both the opposite- and same-sign configurations. Events with additional veto leptons are removed. To suppress the background contribution from DY events, the mass of the dilepton pair ($m_{\ell\ell}$) is required to be greater than 150 GeV.

No b-tagging veto criteria are considered in the analysis, since high mass heavy neutrinos have a significant branching fraction into bottom quark jets, via $N_\ell \rightarrow \ell^- t \bar{b}$ and $N_\ell \rightarrow \ell^+ t \bar{b}$ decays. Furthermore, for lower heavy neutrino masses, final states typically contain jets with very high energies, for which b-tagging performance is degraded and, as a result, systematic uncertainties prevent an improvement in sensitivity.

We define three different signal regions (SRs), depending on the number of AK8 jets found in the event, as shown in table 1. The region SR1 contains no AK8 jets, SR2 contains exactly one AK8 jet, and SR3 contains at least two AK8 jets.

In SR1, we require two tight leptons and at least four AK4 jets, but no AK8 jets. To reconstruct two heavy neutrinos for an event, the two tight leptons and the leading four AK4 jets are used. Each heavy neutrino candidate is reconstructed using one lepton and two AK4 jets. From the 12 possible lepton plus two-jet combination combinations, we select the one that minimizes the difference between the masses of the two heavy neutrino candidates.

In SR2, we require exactly one AK8 jet, at least one tight lepton and at least two AK4 jets. In events with only one tight lepton, this lepton is paired with the two leading AK4 jets to reconstruct one heavy neutrino. The AK8 jet is assigned as a proxy to the other

SR	$N(\text{AK8 jet})$	$N(\text{tight leptons})$	$N(\text{AK4 jet})$
SR1	=0	=2	≥ 4
SR2	=1	≥ 1	≥ 2
SR3	≥ 2	—	—

Table 1. Multiplicity requirements for AK8 jets, tight leptons and AK4 jets in the different signal regions considered in the search.

heavy neutrino. If an event has two tight leptons, the AK8 jet is paired with the closer lepton, and the other lepton is paired with the two leading AK4 jets.

In SR3, we require at least two AK8 jets, and loose leptons are ignored. In cases where an event has no tight leptons, the two leading AK8 jets are assigned as proxies to the two heavy neutrinos. If there is only one tight lepton, the lepton is paired with the closer AK8 jet out of the two leading AK8 jets to reconstruct one heavy neutrino. The remaining AK8 jet is assigned as a proxy to the other heavy neutrino. If there are two tight leptons, the leading AK8 jet is paired to the closer lepton while the secondary AK8 jet and the other lepton are used to reconstruct the other heavy neutrino.

In all three SRs, the invariant mass of the Z' boson is obtained using the two reconstructed heavy neutrinos. If a tight lepton is included in the energy of the AK8 jet, the four-momentum of the AK8 jet is assigned to the four-momentum of the heavy neutrino. For $m_{N_\ell} = 100 \text{ GeV}$ and $m_{Z'} = 4 \text{ TeV}$, about 2% of the signal events that enter the SR3 are of this type. Once the Z' boson and heavy neutrinos are reconstructed, additional requirements (reconstructed $m_{N_\ell} > 80 \text{ GeV}$ and reconstructed $m_{Z'} > 300 \text{ GeV}$) are applied to reduce backgrounds, and we start search range from $m_{Z'} = 400 \text{ GeV}$ and $m_{N_\ell} = 100 \text{ GeV}$.

The products of acceptance and efficiency for the simulated signal samples are 20–25 (25–33)% and 1.2–4.7 (7–16)% in the dielectron (dimuon) channel for the most resolved and boosted mass points, respectively, depending on $m_{Z'}$.

6 Background estimation

The backgrounds are dominated by SM processes containing prompt leptons. These are estimated using the simulated samples described in section 3. The leading background contribution is from $t\bar{t}$ events. The simulated $t\bar{t}$ sample is normalized using the QCD next-to-NLO cross section, which includes resummation of next-to-next-to-leading logarithmic soft gluon terms for the top quark mass of 172.5 GeV [63]. A control region (CR), CR1, is defined to validate the $t\bar{t}$ background using an $e\mu$ sideband passing the same single-muon triggers used in the dimuon SR. The region CR1 has exactly the same event selection as the SR, except the flavors of the two leptons are required to be different. The events of CR1 are classified in three orthogonal regions according to the same criteria used for the SR, which are summarized in table 1. Since no mixing between generations of heavy neutrinos is assumed and the heavy neutrino always decays into a lepton and two jets in the signal process, the signal in CR1 is negligible. This expectation is supported by

the observed consistency between simulated background predictions in CR1 and the corresponding data-driven estimates. Additional scale factors are applied to the $t\bar{t}$ simulated samples, depending on the AK8 jet multiplicity. The scale factors are extracted from a simultaneous fit to all CRs and SRs and range from 0.83 to 1.06. The uncertainties in these scale factors, determined from the fit, are assumed to cover the theoretical uncertainties in the simulated $t\bar{t}$ distributions.

Because of the large production cross section, DY events constitute the second largest background, although most of these are removed by the requirement, $m_{\ell\ell} > 150 \text{ GeV}$. A second CR (CR2) is chosen to verify that the simulated sample of DY events is well modeled. The selection used to define CR2 is similar to that of the SR except the requirement on $m_{\ell\ell}$ is changed to $|m_{\ell\ell} - m_Z| < 10 \text{ GeV}$, where m_Z is the mass of Z boson. The events of CR2 are classified in three orthogonal regions according to the same criteria used for the SR, which are summarized in table 1. To correct for the mismodeling of the dilepton p_T in simulated events, a dedicated correction estimated using inclusive dimuon events is applied as a function of generator level dilepton p_T and mass [64]. The overall normalization factors as functions of the AK8 jet multiplicity are fit using all the CRs and SRs. They range from 0.93 to 1.14 for events with fewer than two AK8 jets. For events with two or more AK8 jets, normalization factors have a wider range from 0.87 to 1.28 owing to the small numbers of events. Figure 3 shows the agreement between the observed and simulated distributions of the reconstructed Z' candidate mass in CR2.

Additional prompt backgrounds are considered from di- and triboson production, $t\bar{t}$ pair production in association with a vector boson, and single top production. The contribution of these backgrounds is found to be almost negligible. We assign a conservative 50% uncertainty in the estimated size of these backgrounds, to account for limited MC samples in the extreme phase space regions of interest. The background contribution with nonprompt leptons, which are leptons not originating from the hard scattering process, is estimated using the simulated samples. We assign an uncertainty of 100% in the estimated size of the nonprompt-lepton background since this depends on detector effects, which are often more difficult to estimate.

7 Systematic uncertainties and fitting procedure

Binned maximum likelihood fits using the distribution of reconstructed $m_{Z'}$ for background and signal as templates are performed to extract the signal contribution for each SR. The choice of bin size is based on the resolution for the reconstructed $m_{Z'}$, which ranges from 150 to 550 GeV. Where necessary, bins are then merged to ensure that the predicted number of MC events is nonzero in all bins of all SRs and CRs for each year. Various sources of uncertainty in both shape and normalization of reconstructed $m_{Z'}$ distributions are considered as nuisance parameters and profiled in the fit. Log-normal and Gaussian prior probability densities are used for normalization and shape uncertainties, respectively. The normalizations of $t\bar{t}$ and DY backgrounds are left freely floating in the fit. The pre-fit and post-fit predictions on the plots of figures 2–4 correspond to the results obtained




Figure 2. Reconstructed mass of the Z' candidate in CR1 ($e\mu$), which consists of flavor sidebands of SR1 (upper), SR2 (middle), and SR3 (lower) regions. Pre-fit (post-fit) results are shown on the left (right). The fitting procedure is described in section 7. The last bin of each plot includes the overflow events. The contribution from DY events is too small to be visible on these plots. The lower panel of each plot shows the ratio of observed events to expected background.




Figure 3. Reconstructed mass of the Z' candidate in CR2 (ee and $\mu\mu$), which consists of the dilepton mass sidebands of SR1 (upper), SR2 (middle), and SR3 (lower) regions. Post-fit dielectron (dimuon) channel results are shown on the left (right). The fitting procedure is described in section 7. The last bin of each plot includes the overflow events. The lower panel of each plot shows the ratio of observed events to expected background.

before and after likelihood fitting, respectively. No nuisance parameter is pulled beyond one standard deviation from the pre-fit value.

For leptons, the efficiencies for reconstruction, identification, isolation, and trigger are estimated using observed and simulated $Z \rightarrow ee, \mu\mu$ events that are not included in the SRs. Scale factors (SFs) are applied to simulated events to correct for mismodeling in the simulation. Systematic uncertainties in momentum scale and resolution [49, 51] are estimated in simulations by varying the lepton momentum scales within their uncertainties.

The jet energy corrections and resolutions estimated using hadronic jets are used for both hadronic jets and AK8 jets with high p_T leptons inside. Such corrections were found to provide a good modeling of jets also in the case of non-resolved lepton(s) within the AK8 jet in CRs. The uncertainty in the jet energy scale and resolution are estimated using the same procedures as those used to determine systematic uncertainty in lepton momenta. An additional uncertainty coming from the soft-drop mass scale is found to be less than 1.3% [65].

The integrated luminosities of the 2016, 2017, and 2018 data-taking periods are individually known with uncertainties in the range 1.2–2.5% [66–68], while the total 2016–2018 integrated luminosity has an uncertainty of 1.6%, the improvement in precision reflecting the (uncorrelated) time evolution of some systematic effects.

The simulation of pileup events is based on a cross section of 69.2 mb for inelastic pp collisions with energy deposits inside the CMS acceptance. A systematic uncertainty for the pileup weight is assigned by varying the cross section up and down by 5%. An additional event weight that accounts for the L1 trigger inefficiency observed in the $|\eta| < 2.0$ region for the 2016 and 2017 data-taking period is also applied [69]. The systematic uncertainty for the L1 trigger inefficiency weight is assigned by varying the weight within its uncertainty.

The uncertainties related to the choice of PDF can affect both the cross section and kinematic distributions of the signal. The effect of the PDF uncertainty and the variation related to the strong coupling constant (α_S) are considered, following the method introduced by PDF4LHC15 [70]. The uncertainties in the renormalization and factorization scales (μ_R and μ_F , respectively) are determined by varying them by a factor of two on an event-by-event basis using combinations in which the two scales do not move in opposite directions at the same time. The effect of these variations on the shape of the $m_{Z'}$ distribution is determined while maintaining the signal normalization.

The normalization uncertainties for the rare SM processes and nonprompt-lepton backgrounds are also incorporated into the likelihood as nuisance parameters. We observe that the fit procedure returns an uncertainty of about 40% in the nuisance parameter of nonprompt-lepton background, which is initially constrained at the 100% level. Since this is consistent with the precision of independent background estimation methods using control samples in data-driven methods [15, 16, 71], we use the fit to constrain the normalization.

The uncertainty in the dilepton p_T shape corrections to the simulated DY events is estimated by varying correction factors within their uncertainties.

A complete list of these uncertainties in the dielectron and dimuon channels is shown in table 2. The lepton trigger scale factors, jet energy resolution, L1 trigger inefficiency, and nonprompt lepton background normalizations are assumed to be uncorrelated between the 2016, 2017, and 2018 data sets, since their corrections and corresponding uncertainties

Source	Bkgd./Signal process	Year-to-year treatment	ee bkgd. (%)	ee signal (%)	$\mu\mu$ bkgd. (%)	$\mu\mu$ signal (%)
Integrated luminosity	All bkgd./Signal	Correlated	1.6	1.6	1.6	1.6
Electron momentum scale	All bkgd./Signal	Correlated	0.1–0.9	0.0–0.5	<0.1	<0.1
Electron momentum smear	All bkgd./Signal	Correlated	<0.1	0.0–0.1	<0.1	<0.1
Muon momentum scale	All bkgd./Signal	Correlated	<0.1	0.0–0.1	0.0–5.0	0.0–1.1
Jet energy resolution	All bkgd./Signal	Uncorrelated	1.4–14.4	0.0–10.8	0.5–14.8	0.0–9.5
Jet energy scale	All bkgd./Signal	Correlated	5.4–15.3	0.0–11.4	3.6–16.7	0.0–9.2
Soft-drop mass scale	All bkgd./Signal	Correlated	0.0–0.5	0.0–1.3	0.0–0.5	0.0–1.2
Pileup weight	All bkgd./Signal	Correlated	0.0–2.0	0.0–2.9	0.0–1.7	0.0–1.9
L1 trigger inefficiency	All bkgd./Signal	Uncorrelated	0.0–1.8	0.0–2.3	0.0–1.1	0.0–1.0
Electron reconstruction SF	All bkgd./Signal	Correlated	1.3–2.2	1.1–2.3	<0.1	<0.1
Electron ident. SF	All bkgd./Signal	Correlated	1.9–4.0	0.3–5.3	<0.1	<0.1
Electron trigger SF	All bkgd./Signal	Uncorrelated	0.4–2.8	0.1–15.1	<0.1	<0.1
Muon reconstruction SF	All bkgd./Signal	Correlated	<0.1	<0.1	0.3–1.0	0.3–10.4
Muon ident. SF	All bkgd./Signal	Correlated	<0.1	<0.1	0.2–1.6	0.2–1.6
Muon isolation SF	All bkgd./Signal	Correlated	<0.1	<0.1	<0.1	<0.1
Muon trigger SF	All bkgd./Signal	Uncorrelated	<0.1	<0.1	0.1–0.2	0.1–1.3
Dilepton p_T shape correction	DY+jets	Correlated	0.0–19.2	—	0.0–18.8	—
PDF	Signal	Correlated	—	<0.1	—	<0.1
Scale (μ_R, μ_F)	Signal	Correlated	—	<0.1	—	<0.1
Nonprompt norm.	Nonprompt	Uncorrelated	100	—	100	—
Rare SM norm.	Others	Correlated	50	—	50	—

Table 2. Systematic uncertainties and their impacts on the total number of events in the signal regions before the binned maximum likelihood fit.

are affected by differences in operating conditions and detector performance between data-taking years. Other experimental and all theoretical uncertainties are taken as correlated.

8 Results and discussion

The background-only post-fit invariant mass distributions of Z' candidates in the different SRs are shown in figure 4. Representative signal distributions are also shown.

We do not observe any significant excess in data with respect to the SM prediction. The 95% confidence level (CL) upper limits on the product of signal cross section and BF are obtained using the distributions of the likelihood ratio calculated with the asymptotic approximation [72] and the CL_s criterion [73, 74].

The observed and expected exclusion limits on the signal cross section times BF are shown in figure 5. The maximum local significance is 3.32σ , observed in the dielectron channel for the $(m_{Z'}, m_{N_\ell}) = (4.6, 0.1)$ TeV. The probability to observe similar or larger excess in the dielectron channel across the full analysis mass range is estimated with pseudo experiments under background-only hypothesis. This probability is 1.12×10^{-2} , corresponding to a global significance of 2.28σ . For cases where $m_{N_\ell} = m_{Z'}/4$, the observed (expected) lower limits at 95% CL on the mass of the Z' boson are 3.59 (3.90) TeV in the dielectron channel and 4.10 (3.86) TeV in the dimuon channel. For the phase space with mostly boosted signals ($m_{N_\ell} = 100$ GeV), the observed (expected) lower limits are found




Figure 4. Distributions of the reconstructed Z' candidate mass in SR1 (upper row), SR2 (middle row) and SR3 (lower row). The left (right) column corresponds to the dielectron (dimuon) channel. Signal samples with $(m_{Z'}, m_{N_\ell}) = (1000, 100)$ GeV, $(1000, 300)$ GeV, $(4000, 200)$ GeV, and $(4000, 1200)$ GeV are shown together with a reference cross section times branching fraction of 1 fb. The last bin of each plot includes the overflow events. The lower panel of each plot shows the ratio of observed events to expected background.




Figure 5. The observed and expected 95% CL upper limits on the product of Z' boson cross section and branching fraction are shown for the case of $m_{N_\ell} = m_{Z'}/4$ (upper row) and $m_{N_\ell} = 100$ GeV (lower row) for dielectron (left column) and dimuon (right column) channels. The green and yellow bands indicate the 68% and 95% CL regions around the expected limit. The red lines represent values coming from the benchmark LRSM model [45]. Both opposite- and same-sign dileptons from decays of heavy neutrino pairs are considered.

to be $m_{Z'} = 2.79$ (3.12) TeV and 4.38 (4.22) TeV in the dielectron and dimuon channels, respectively. The use of a dedicated algorithm for boosted signals provides a significant improvement in sensitivity, in particular for the muon channel. We note that the sensitivity for Dirac-type heavy neutrinos is identical as long as the branching fraction of the Z' boson to a heavy neutrino pair stays the same.

One of the assumptions in this analysis is the presence of a nonnegligible Z' coupling to quarks. A comparison with the sensitivity of the search for high mass dijet resonances performed by CMS using the same data set [75] is therefore possible. For comparable Z'




Figure 6. Observed and expected exclusion regions at 95% CL in the 2D phase space of $m_{Z'}$ vs. m_{N_ℓ} for dielectron (left) and dimuon (right) channels. One standard deviation (s.d.) limits are also shown. Both opposite- and same-sign dileptons from decays of heavy neutrino pairs are considered.

branching fractions into quarks and heavy neutrinos, our search is found to be about one order of magnitude more sensitive.

The upper limits across the m_{N_ℓ} - $m_{Z'}$ plane are shown in figure 6. The SR1 is sensitive in the phase space regions $m_{Z'} = 1$ to 2 TeV and $m_{N_\ell} = 0.5$ to 1 TeV. The SR2 is particularly sensitive for higher $m_{Z'}$ with resolved heavy neutrinos. The SR3 is most important where heavy neutrinos are boosted. The dimuon channel shows better sensitivity where heavy neutrinos are mostly boosted because the dielectron channel suffers for the H/E requirements discussed in section 4. In the dielectron channel, for $m_{N_\ell} > 0.9$ TeV and $m_{Z'} > 2.85$ TeV, the observed limits obtained in the SR2 are better than those of the combined limits. This indicates that the slight excess observed for the combined limits in this region of phase space is driven by the SR3.

A priori, a further categorization separating same-sign and opposite-sign dilepton pair events might increase the sensitivity. However, with the current data sample size, the evaluation of same-sign dilepton backgrounds using control samples, particularly in a highly boosted regime for heavy neutrinos, is affected by large systematic uncertainties. Thus introducing sign categorization would not improve the present search.

9 Summary

A search for the pair production of heavy Majorana neutrinos (N_ℓ) via the decay of a Z' boson, in a final state with two same-flavor leptons and at least two reconstructed jets has been performed in proton-proton collisions at a center-of-mass energy of 13 TeV, using LHC 2016–2018 data corresponding to an integrated luminosity of 138 fb^{-1} . No significant excess of events beyond the expected background is observed. Upper limits on the product of signal cross section and branching fraction in the context of a left-right symmetry model scenario are set [45]. Exclusion regions in the dielectron and dimuon channels are set at 95% confidence level. These are the first results of a search for this process at 13 TeV and are the most restrictive limits to date on the mass of N_ℓ as a function of the Z' boson mass.

Acknowledgments

We congratulate our colleagues in the CERN accelerator departments for the excellent performance of the LHC and thank the technical and administrative staffs at CERN and at other CMS institutes for their contributions to the success of the CMS effort. In addition, we gratefully acknowledge the computing centers and personnel of the Worldwide LHC Computing Grid and other centers for delivering so effectively the computing infrastructure essential to our analyses. Finally, we acknowledge the enduring support for the construction and operation of the LHC, the CMS detector, and the supporting computing infrastructure provided by the following funding agencies: SC (Armenia), BMBWF and FWF (Austria); FNRS and FWO (Belgium); CNPq, CAPES, FAPERJ, FAPERGS, and FAPESP (Brazil); MES and BNSF (Bulgaria); CERN; CAS, MoST, and NSFC (China); MINCIENCIAS (Colombia); MSES and CSF (Croatia); RIF (Cyprus); SENESCYT (Ecuador); MoER, ERC PUT and ERDF (Estonia); Academy of Finland, MEC, and HIP (Finland); CEA and CNRS/IN2P3 (France); BMBF, DFG, and HGF (Germany); GSRI (Greece); NKFIH (Hungary); DAE and DST (India); IPM (Iran); SFI (Ireland); INFN (Italy); MSIP and NRF (Republic of Korea); MES (Latvia); LAS (Lithuania); MOE and UM (Malaysia); BUAP, CINVESTAV, CONACYT, LNS, SEP, and UASLP-FAI (Mexico); MOS (Montenegro); MBIE (New Zealand); PAEC (Pakistan); MES and NSC (Poland); FCT (Portugal); MESTD (Serbia); MCIN/AEI and PCTI (Spain); MOSTR (Sri Lanka); Swiss Funding Agencies (Switzerland); MST (Taipei); MHESI and NSTDA (Thailand); TUBITAK and TENMAK (Turkey); NASU (Ukraine); STFC (United Kingdom); DOE and NSF (U.S.A.).

Individuals have received support from the Marie-Curie program and the European Research Council and Horizon 2020 Grant, contract Nos. 675440, 724704, 752730, 758316, 765710, 824093, 884104, and COST Action CA16108 (European Union); the Leventis Foundation; the Alfred P. Sloan Foundation; the Alexander von Humboldt Foundation; the Science Committee, project no. 22rl-037 (Armenia); the Belgian Federal Science Policy Office; the Fonds pour la Formation à la Recherche dans l’Industrie et dans l’Agriculture (FRIA-Belgium); the Agentschap voor Innovatie door Wetenschap en Technologie (IWT-Belgium); the F.R.S.-FNRS and FWO (Belgium) under the “Excellence of Science — EOS” — be.h project n. 30820817; the Beijing Municipal Science & Technology Commission, No. Z191100007219010; the Ministry of Education, Youth and Sports (MEYS) of the Czech Republic; the Hellenic Foundation for Research and Innovation (HFRI), Project Number 2288 (Greece); the Deutsche Forschungsgemeinschaft (DFG), under Germany’s Excellence Strategy — EXC 2121 “Quantum Universe” — 390833306, and under project number 400140256 — GRK2497; the Hungarian Academy of Sciences, the New National Excellence Program — ÚNKP, the NKFIH research grants K 124845, K 124850, K 128713, K 128786, K 129058, K 131991, K 133046, K 138136, K 143460, K 143477, 2020-2.2.1-ED-2021-00181, and TKP2021-NKTA-64 (Hungary); the Council of Science and Industrial Research, India; the Latvian Council of Science; the Ministry of Education and Science, project no. 2022/WK/14, and the National Science Center, contracts Opus 2021/41/B/ST2/01369 and 2021/43/B/ST2/01552 (Poland); the Fundação para a Ciência e a Tecnologia, grant CEECIND/01334/2018 (Portugal); the National Priorities Research Program by Qatar

National Research Fund; MCIN/AEI/10.13039/501100011033, ERDF “a way of making Europe”, and the Programa Estatal de Fomento de la Investigación Científica y Técnica de Excelencia María de Maeztu, grant MDM-2017-0765 and Programa Severo Ochoa del Principado de Asturias (Spain); the Chulalongkorn Academic into Its 2nd Century Project Advancement Project, and the National Science, Research and Innovation Fund via the Program Management Unit for Human Resources & Institutional Development, Research and Innovation, grant B05F650021 (Thailand); the Kavli Foundation; the Nvidia Corporation; the SuperMicro Corporation; the Welch Foundation, contract C-1845; and the Weston Havens Foundation (U.S.A.).

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](#)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] PARTICLE DATA GROUP collaboration, *Review of particle physics*, *PTEP* **2022** (2022) 083C01 [[INSPIRE](#)].
- [2] SNO collaboration, *Direct evidence for neutrino flavor transformation from neutral current interactions in the Sudbury Neutrino Observatory*, *Phys. Rev. Lett.* **89** (2002) 011301 [[nucl-ex/0204008](#)] [[INSPIRE](#)].
- [3] SUPER-KAMIOKANDE collaboration, *Evidence for oscillation of atmospheric neutrinos*, *Phys. Rev. Lett.* **81** (1998) 1562 [[hep-ex/9807003](#)] [[INSPIRE](#)].
- [4] SUPER-KAMIOKANDE collaboration, *Evidence for an oscillatory signature in atmospheric neutrino oscillation*, *Phys. Rev. Lett.* **93** (2004) 101801 [[hep-ex/0404034](#)] [[INSPIRE](#)].
- [5] E. Di Valentino, S. Gariazzo and O. Mena, *Most constraining cosmological neutrino mass bounds*, *Phys. Rev. D* **104** (2021) 083504 [[arXiv:2106.15267](#)] [[INSPIRE](#)].
- [6] KATRIN collaboration, *Direct neutrino-mass measurement with sub-electronvolt sensitivity*, *Nature Phys.* **18** (2022) 160 [[arXiv:2105.08533](#)] [[INSPIRE](#)].
- [7] A. Maiezza, M. Nemevsek, F. Nesti and G. Senjanovic, *Left-right symmetry at LHC*, *Phys. Rev. D* **82** (2010) 055022 [[arXiv:1005.5160](#)] [[INSPIRE](#)].
- [8] P. Minkowski, $\mu \rightarrow e\gamma$ at a rate of one out of 10^9 muon decays?, *Phys. Lett. B* **67** (1977) 421 [[INSPIRE](#)].
- [9] M. Gell-Mann, P. Ramond and R. Slansky, *Complex spinors and unified theories*, in the proceedings of the *Proc. supergravity workshop at Stony Brook*, North-Holland, Amsterdam, The Netherlands (1979), p. 341 [[arXiv:1306.4669](#)] [[INSPIRE](#)].
- [10] T. Yanagida, *Horizontal gauge symmetry and masses of neutrinos*, in the proceedings of the *Proc. workshop on the unified theory and the baryon number in the universe*, (1979), p. 95 [[INSPIRE](#)].
- [11] R.N. Mohapatra and G. Senjanovic, *Neutrino mass and spontaneous parity nonconservation*, *Phys. Rev. Lett.* **44** (1980) 912 [[INSPIRE](#)].
- [12] M. Doi et al., *CP violation in Majorana neutrinos*, *Phys. Lett. B* **102** (1981) 323 [[INSPIRE](#)].

- [13] CMS collaboration, *Search for resonant and nonresonant new phenomena in high-mass dilepton final states at $\sqrt{s} = 13$ TeV*, *JHEP* **07** (2021) 208 [[arXiv:2103.02708](#)] [[INSPIRE](#)].
- [14] CMS collaboration, *Search for new physics in the lepton plus missing transverse momentum final state in proton-proton collisions at $\sqrt{s} = 13$ TeV*, *JHEP* **07** (2022) 067 [[arXiv:2202.06075](#)] [[INSPIRE](#)].
- [15] CMS collaboration, *Search for heavy Majorana neutrinos in same-sign dilepton channels in proton-proton collisions at $\sqrt{s} = 13$ TeV*, *JHEP* **01** (2019) 122 [[arXiv:1806.10905](#)] [[INSPIRE](#)].
- [16] CMS collaboration, *Search for heavy neutral leptons in events with three charged leptons in proton-proton collisions at $\sqrt{s} = 13$ TeV*, *Phys. Rev. Lett.* **120** (2018) 221801 [[arXiv:1802.02965](#)] [[INSPIRE](#)].
- [17] CMS collaboration, *Search for long-lived heavy neutral leptons with displaced vertices in proton-proton collisions at $\sqrt{s} = 13$ TeV*, *JHEP* **07** (2022) 081 [[arXiv:2201.05578](#)] [[INSPIRE](#)].
- [18] ATLAS collaboration, *Search for heavy neutrinos and right-handed W bosons in events with two leptons and jets in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector*, *Eur. Phys. J. C* **72** (2012) 2056 [[arXiv:1203.5420](#)] [[INSPIRE](#)].
- [19] CMS collaboration, *Search for heavy neutrinos and W_R bosons with right-handed couplings in a left-right symmetric model in pp collisions at $\sqrt{s} = 7$ TeV*, *Phys. Rev. Lett.* **109** (2012) 261802 [[arXiv:1210.2402](#)] [[INSPIRE](#)].
- [20] CMS collaboration, *Search for heavy neutrinos and W bosons with right-handed couplings in proton-proton collisions at $\sqrt{s} = 8$ TeV*, *Eur. Phys. J. C* **74** (2014) 3149 [[arXiv:1407.3683](#)] [[INSPIRE](#)].
- [21] ATLAS collaboration, *Search for heavy Majorana neutrinos with the ATLAS detector in pp collisions at $\sqrt{s} = 8$ TeV*, *JHEP* **07** (2015) 162 [[arXiv:1506.06020](#)] [[INSPIRE](#)].
- [22] CMS collaboration, *Search for heavy neutrinos or third-generation leptoquarks in final states with two hadronically decaying τ leptons and two jets in proton-proton collisions at $\sqrt{s} = 13$ TeV*, *JHEP* **03** (2017) 077 [[arXiv:1612.01190](#)] [[INSPIRE](#)].
- [23] CMS collaboration, *Search for third-generation scalar leptoquarks and heavy right-handed neutrinos in final states with two tau leptons and two jets in proton-proton collisions at $\sqrt{s} = 13$ TeV*, *JHEP* **07** (2017) 121 [[arXiv:1703.03995](#)] [[INSPIRE](#)].
- [24] CMS collaboration, *Search for a heavy right-handed W boson and a heavy neutrino in events with two same-flavor leptons and two jets at $\sqrt{s} = 13$ TeV*, *JHEP* **05** (2018) 148 [[arXiv:1803.11116](#)] [[INSPIRE](#)].
- [25] ATLAS collaboration, *Search for heavy Majorana or Dirac neutrinos and right-handed W gauge bosons in final states with two charged leptons and two jets at $\sqrt{s} = 13$ TeV with the ATLAS detector*, *JHEP* **01** (2019) 016 [[arXiv:1809.11105](#)] [[INSPIRE](#)].
- [26] ATLAS collaboration, *Search for a right-handed gauge boson decaying into a high-momentum heavy neutrino and a charged lepton in pp collisions with the ATLAS detector at $\sqrt{s} = 13$ TeV*, *Phys. Lett. B* **798** (2019) 134942 [[arXiv:1904.12679](#)] [[INSPIRE](#)].
- [27] CMS collaboration, *Search for a right-handed W boson and a heavy neutrino in proton-proton collisions at $\sqrt{s} = 13$ TeV*, *JHEP* **04** (2022) 047 [[arXiv:2112.03949](#)] [[INSPIRE](#)].

- [28] ATLAS collaboration, *Search for heavy Majorana or Dirac neutrinos and right-handed W gauge bosons in final states with charged leptons and jets in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*, [arXiv:2304.09553](https://arxiv.org/abs/2304.09553) [INSPIRE].
- [29] *HEPData record for this analysis*, CMS-EXO-20-006 (2023)
- [30] CMS collaboration, *The CMS experiment at the CERN LHC, 2008* JINST **3** S08004 [INSPIRE].
- [31] J. Alwall et al., *The automated computation of tree-level and next-to-leading order differential cross sections, and their matching to parton shower simulations*, JHEP **07** (2014) 079 [[arXiv:1405.0301](https://arxiv.org/abs/1405.0301)] [INSPIRE].
- [32] J. Alwall et al., *Comparative study of various algorithms for the merging of parton showers and matrix elements in hadronic collisions*, Eur. Phys. J. C **53** (2008) 473 [[arXiv:0706.2569](https://arxiv.org/abs/0706.2569)] [INSPIRE].
- [33] P. Nason, *A new method for combining NLO QCD with shower Monte Carlo algorithms*, JHEP **11** (2004) 040 [[hep-ph/0409146](https://arxiv.org/abs/hep-ph/0409146)] [INSPIRE].
- [34] S. Frixione, P. Nason and C. Oleari, *Matching NLO QCD computations with parton shower simulations: the POWHEG method*, JHEP **11** (2007) 070 [[arXiv:0709.2092](https://arxiv.org/abs/0709.2092)] [INSPIRE].
- [35] S. Alioli, P. Nason, C. Oleari and E. Re, *A general framework for implementing NLO calculations in shower Monte Carlo programs: the POWHEG BOX*, JHEP **06** (2010) 043 [[arXiv:1002.2581](https://arxiv.org/abs/1002.2581)] [INSPIRE].
- [36] S. Frixione, P. Nason and G. Ridolfi, *A positive-weight next-to-leading-order Monte Carlo for heavy flavour hadroproduction*, JHEP **09** (2007) 126 [[arXiv:0707.3088](https://arxiv.org/abs/0707.3088)] [INSPIRE].
- [37] T. Sjöstrand et al., *An introduction to PYTHIA 8.2*, Comput. Phys. Commun. **191** (2015) 159 [[arXiv:1410.3012](https://arxiv.org/abs/1410.3012)] [INSPIRE].
- [38] S. Alioli, P. Nason, C. Oleari and E. Re, *NLO single-top production matched with shower in POWHEG: s- and t-channel contributions*, JHEP **09** (2009) 111 [*Erratum ibid.* **02** (2010) 011] [[arXiv:0907.4076](https://arxiv.org/abs/0907.4076)] [INSPIRE].
- [39] E. Re, *Single-top Wt-channel production matched with parton showers using the POWHEG method*, Eur. Phys. J. C **71** (2011) 1547 [[arXiv:1009.2450](https://arxiv.org/abs/1009.2450)] [INSPIRE].
- [40] NNPDF collaboration, *Parton distributions for the LHC run II*, JHEP **04** (2015) 040 [[arXiv:1410.8849](https://arxiv.org/abs/1410.8849)] [INSPIRE].
- [41] NNPDF collaboration, *Parton distributions from high-precision collider data*, Eur. Phys. J. C **77** (2017) 663 [[arXiv:1706.00428](https://arxiv.org/abs/1706.00428)] [INSPIRE].
- [42] CMS collaboration, *Event generator tunes obtained from underlying event and multiparton scattering measurements*, Eur. Phys. J. C **76** (2016) 155 [[arXiv:1512.00815](https://arxiv.org/abs/1512.00815)] [INSPIRE].
- [43] CMS collaboration, *Extraction and validation of a new set of CMS PYTHIA8 tunes from underlying-event measurements*, Eur. Phys. J. C **80** (2020) 4 [[arXiv:1903.12179](https://arxiv.org/abs/1903.12179)] [INSPIRE].
- [44] GEANT4 collaboration, *GEANT4 — a simulation toolkit*, Nucl. Instrum. Meth. A **506** (2003) 250 [INSPIRE].
- [45] O. Mattelaer, M. Mitra and R. Ruiz, *Automated neutrino jet and top jet predictions at next-to-leading-order with parton shower matching in effective left-right symmetric models*, [arXiv:1610.08985](https://arxiv.org/abs/1610.08985) [INSPIRE].

- [46] CMS collaboration, *The CMS trigger system*, **2017 JINST** **12** P01020 [[arXiv:1609.02366](#)] [[INSPIRE](#)].
- [47] D. Contardo et al., *Technical proposal for the phase-II upgrade of the CMS detector*, CERN-LHCC-2015-010, CERN, Geneva, Switzerland (2015) [[DOI:10.17181/CERN.VU8I.D59J](#)].
- [48] CMS collaboration, *Particle-flow reconstruction and global event description with the CMS detector*, **2017 JINST** **12** P10003 [[arXiv:1706.04965](#)] [[INSPIRE](#)].
- [49] CMS collaboration, *Electron and photon reconstruction and identification with the CMS experiment at the CERN LHC*, **2021 JINST** **16** P05014 [[arXiv:2012.06888](#)] [[INSPIRE](#)].
- [50] CMS collaboration, *ECAL 2016 refined calibration and run2 summary plots*, CMS-DP-2020-021, CERN, Geneva, Switzerland (2020).
- [51] CMS collaboration, *Performance of the CMS muon detector and muon reconstruction with proton-proton collisions at $\sqrt{s} = 13$ TeV*, **2018 JINST** **13** P06015 [[arXiv:1804.04528](#)] [[INSPIRE](#)].
- [52] CMS collaboration, *Search for high-mass resonances in dilepton final states in proton-proton collisions at $\sqrt{s} = 13$ TeV*, **JHEP** **06** (2018) 120 [[arXiv:1803.06292](#)] [[INSPIRE](#)].
- [53] M. Cacciari, G.P. Salam and G. Soyez, *The anti- k_t jet clustering algorithm*, **JHEP** **04** (2008) 063 [[arXiv:0802.1189](#)] [[INSPIRE](#)].
- [54] M. Cacciari, G.P. Salam and G. Soyez, *FastJet user manual*, **Eur. Phys. J. C** **72** (2012) 1896 [[arXiv:1111.6097](#)] [[INSPIRE](#)].
- [55] D. Bertolini, P. Harris, M. Low and N. Tran, *Pileup per particle identification*, **JHEP** **10** (2014) 059 [[arXiv:1407.6013](#)] [[INSPIRE](#)].
- [56] CMS collaboration, *Pileup mitigation at CMS in 13 TeV data*, **2020 JINST** **15** P09018 [[arXiv:2003.00503](#)] [[INSPIRE](#)].
- [57] CMS collaboration, *Jet algorithms performance in 13 TeV data*, CMS-PAS-JME-16-003, CERN, Geneva, Switzerland (2017).
- [58] M. Dasgupta, A. Fregoso, S. Marzani and G.P. Salam, *Towards an understanding of jet substructure*, **JHEP** **09** (2013) 029 [[arXiv:1307.0007](#)] [[INSPIRE](#)].
- [59] J.M. Butterworth, A.R. Davison, M. Rubin and G.P. Salam, *Jet substructure as a new Higgs search channel at the LHC*, **Phys. Rev. Lett.** **100** (2008) 242001 [[arXiv:0802.2470](#)] [[INSPIRE](#)].
- [60] Y.L. Dokshitzer, G.D. Leder, S. Moretti and B.R. Webber, *Better jet clustering algorithms*, **JHEP** **08** (1997) 001 [[hep-ph/9707323](#)] [[INSPIRE](#)].
- [61] M. Wobisch and T. Wengler, *Hadronization corrections to jet cross-sections in deep inelastic scattering*, in the proceedings of the *Workshop on Monte Carlo generators for HERA physics (plenary starting meeting)*, (1998), p. 270 [[hep-ph/9907280](#)] [[INSPIRE](#)].
- [62] A.J. Larkoski, S. Marzani, G. Soyez and J. Thaler, *Soft drop*, **JHEP** **05** (2014) 146 [[arXiv:1402.2657](#)] [[INSPIRE](#)].
- [63] M. Czakon and A. Mitov, *Top++: a program for the calculation of the top-pair cross-section at hadron colliders*, **Comput. Phys. Commun.** **185** (2014) 2930 [[arXiv:1112.5675](#)] [[INSPIRE](#)].
- [64] CMS collaboration, *Measurements of differential Z boson production cross sections in proton-proton collisions at $\sqrt{s} = 13$ TeV*, **JHEP** **12** (2019) 061 [[arXiv:1909.04133](#)] [[INSPIRE](#)].

- [65] CMS collaboration, *Jet energy scale and resolution in the CMS experiment in pp collisions at 8 TeV*, *2017 JINST* **12** P02014 [[arXiv:1607.03663](#)] [[INSPIRE](#)].
- [66] CMS collaboration, *Precision luminosity measurement in proton-proton collisions at $\sqrt{s} = 13 \text{ TeV}$ in 2015 and 2016 at CMS*, *Eur. Phys. J. C* **81** (2021) 800 [[arXiv:2104.01927](#)] [[INSPIRE](#)].
- [67] CMS collaboration, *CMS luminosity measurement for the 2017 data-taking period at $\sqrt{s} = 13 \text{ TeV}$* , **CMS-PAS-LUM-17-004**, CERN, Geneva, Switzerland (2018).
- [68] CMS collaboration, *CMS luminosity measurement for the 2018 data-taking period at $\sqrt{s} = 13 \text{ TeV}$* , **CMS-PAS-LUM-18-002**, CERN, Geneva, Switzerland (2019).
- [69] CMS collaboration, *Performance of the CMS level-1 trigger in proton-proton collisions at $\sqrt{s} = 13 \text{ TeV}$* , *2020 JINST* **15** P10017 [[arXiv:2006.10165](#)] [[INSPIRE](#)].
- [70] J. Butterworth et al., *PDF4LHC recommendations for LHC run II*, *J. Phys. G* **43** (2016) 023001 [[arXiv:1510.03865](#)] [[INSPIRE](#)].
- [71] CMS collaboration, *Search for new physics in same-sign dilepton events in proton-proton collisions at $\sqrt{s} = 13 \text{ TeV}$* , *Eur. Phys. J. C* **76** (2016) 439 [[arXiv:1605.03171](#)] [[INSPIRE](#)].
- [72] G. Cowan, K. Cranmer, E. Gross and O. Vitells, *Asymptotic formulae for likelihood-based tests of new physics*, *Eur. Phys. J. C* **71** (2011) 1554 [Erratum *ibid.* **73** (2013) 2501] [[arXiv:1007.1727](#)] [[INSPIRE](#)].
- [73] T. Junk, *Confidence level computation for combining searches with small statistics*, *Nucl. Instrum. Meth. A* **434** (1999) 435 [[hep-ex/9902006](#)] [[INSPIRE](#)].
- [74] A.L. Read, *Presentation of search results: the CL_s technique*, *J. Phys. G* **28** (2002) 2693 [[INSPIRE](#)].
- [75] CMS collaboration, *Search for high mass dijet resonances with a new background prediction method in proton-proton collisions at $\sqrt{s} = 13 \text{ TeV}$* , *JHEP* **05** (2020) 033 [[arXiv:1911.03947](#)] [[INSPIRE](#)].

The CMS collaboration

Yerevan Physics Institute, Yerevan, Armenia

A. Tumasyan 

Institut für Hochenergiephysik, Vienna, Austria

W. Adam , J.W. Andrejkovic, T. Bergauer , S. Chatterjee , K. Damanakis ,
 M. Dragicevic , A. Escalante Del Valle , P.S. Hussain , M. Jeitler ¹, N. Krammer ,
 L. Lechner , D. Liko , I. Mikulec , P. Paulitsch, F.M. Pitters, J. Schieck ¹, R. Schöfbeck ,
 D. Schwarz , S. Templ , W. Waltenberger , C.-E. Wulz ¹

Universiteit Antwerpen, Antwerpen, Belgium

M.R. Darwish ², T. Janssen , T. Kello³, H. Rejeb Sfar, P. Van Mechelen 

Vrije Universiteit Brussel, Brussel, Belgium

E.S. Bols , J. D'Hondt , A. De Moor , M. Delcourt , H. El Faham , S. Lowette ,
 S. Moortgat , A. Morton , D. Müller , A.R. Sahasransu , S. Tavernier , W. Van Doninck,
 D. Vannerom 

Université Libre de Bruxelles, Bruxelles, Belgium

B. Clerbaux , G. De Lentdecker , L. Favart , D. Hohov , J. Jaramillo , K. Lee ,
 M. Mahdavikhorrami , I. Makarenko , A. Malara , S. Paredes , L. Pétré , N. Postiau,
 E. Starling , L. Thomas , M. Vanden Bemden , C. Vander Velde , P. Vanlaer 

Ghent University, Ghent, Belgium

D. Dobur , J. Knolle , L. Lambrecht , G. Mestdach, M. Niedziela , C. Rendón, C. Roskas ,
 A. Samalan, K. Skovpen , M. Tytgat , N. Van Den Bossche , B. Vermassen, L. Wezenbeek 

Université Catholique de Louvain, Louvain-la-Neuve, Belgium

A. Benecke , G. Bruno , F. Bury , C. Caputo , P. David , C. Delaere , I.S. Donertas ,
 A. Giannmanco , K. Jaffel , Sa. Jain , V. Lemaitre, K. Mondal , J. Prisciandaro,
 A. Taliercio , T.T. Tran , P. Vischia , S. Wertz 

Centro Brasileiro de Pesquisas Fisicas, Rio de Janeiro, Brazil

G.A. Alves , E. Coelho , C. Hensel , A. Moraes , P. Rebello Teles 

Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil

W.L. Aldá Júnior , M. Alves Gallo Pereira , M. Barroso Ferreira Filho ,
 H. Brandao Malbouisson , W. Carvalho , J. Chinellato⁴, E.M. Da Costa ,
 G.G. Da Silveira ⁵, D. De Jesus Damiao , V. Dos Santos Sousa , S. Fonseca De Souza ,
 J. Martins ⁶, C. Mora Herrera , K. Mota Amarilo , L. Mundim , H. Nogima ,
 A. Santoro , S.M. Silva Do Amaral , A. Sznajder , M. Thiel ,
 F. Torres Da Silva De Araujo ⁷, A. Vilela Pereira 

Universidade Estadual Paulista, Universidade Federal do ABC, São Paulo, Brazil

C.A. Bernardes⁵, L. Calligaris^{ID}, T.R. Fernandez Perez Tomei^{ID}, E.M. Gregores^{ID}, P.G. Mercadante^{ID}, S.F. Novaes^{ID}, Sandra S. Padula^{ID}

Institute for Nuclear Research and Nuclear Energy, Bulgarian Academy of Sciences, Sofia, Bulgaria

A. Aleksandrov^{ID}, G. Antchev^{ID}, R. Hadjiiska^{ID}, P. Iaydjiev^{ID}, M. Misheva^{ID}, M. Rodozov, M. Shopova^{ID}, G. Sultanov^{ID}

University of Sofia, Sofia, Bulgaria

A. Dimitrov^{ID}, T. Ivanov^{ID}, L. Litov^{ID}, B. Pavlov^{ID}, P. Petkov^{ID}, A. Petrov^{ID}, E. Shumka^{ID}

Instituto De Alta Investigación, Universidad de Tarapacá, Casilla 7 D, Arica, Chile

S. Thakur^{ID}

Beihang University, Beijing, China

T. Cheng^{ID}, T. Javaid^{ID}⁸, M. Mittal^{ID}, L. Yuan^{ID}

Department of Physics, Tsinghua University, Beijing, China

M. Ahmad^{ID}, G. Bauer⁹, Z. Hu^{ID}, S. Lezki^{ID}, K. Yi^{ID}^{9,10}

Institute of High Energy Physics, Beijing, China

G.M. Chen^{ID}⁸, H.S. Chen^{ID}⁸, M. Chen^{ID}⁸, F. Iemmi^{ID}, C.H. Jiang, A. Kapoor^{ID}, H. Liao^{ID}, Z.-A. Liu^{ID}¹¹, V. Milosevic^{ID}, F. Monti^{ID}, R. Sharma^{ID}, J. Tao^{ID}, J. Thomas-Wilske^{ID}, J. Wang^{ID}, H. Zhang^{ID}, J. Zhao^{ID}

State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing, China

A. Agapitos^{ID}, Y. An^{ID}, Y. Ban^{ID}, C. Chen, A. Levin^{ID}, C. Li^{ID}, Q. Li^{ID}, X. Lyu, Y. Mao, S.J. Qian^{ID}, X. Sun^{ID}, D. Wang^{ID}, J. Xiao^{ID}, H. Yang

Sun Yat-Sen University, Guangzhou, China

M. Lu^{ID}, Z. You^{ID}

Institute of Modern Physics and Key Laboratory of Nuclear Physics and Ion-beam Application (MOE) — Fudan University, Shanghai, China

X. Gao^{ID}³, D. Leggat, H. Okawa^{ID}, Y. Zhang^{ID}

Zhejiang University, Hangzhou, Zhejiang, China

Z. Lin^{ID}, C. Lu^{ID}, M. Xiao^{ID}

Universidad de Los Andes, Bogota, Colombia

C. Avila^{ID}, D.A. Barbosa Trujillo, A. Cabrera^{ID}, C. Florez^{ID}, J. Fraga^{ID}

Universidad de Antioquia, Medellin, ColombiaJ. Mejia Guisao , F. Ramirez , M. Rodriguez , J.D. Ruiz Alvarez **University of Split, Faculty of Electrical Engineering, Mechanical Engineering and Naval Architecture, Split, Croatia**D. Giljanovic , N. Godinovic , D. Lelas , I. Puljak **University of Split, Faculty of Science, Split, Croatia**Z. Antunovic, M. Kovac , T. Sculac **Institute Rudjer Boskovic, Zagreb, Croatia**V. Brigljevic , B.K. Chitroda , D. Ferencek , D. Majumder , M. Roguljic , A. Starodumov¹² , T. Susa **University of Cyprus, Nicosia, Cyprus**A. Attikis , K. Christoforou , G. Kole , M. Kolosova , S. Konstantinou , J. Mousa , C. Nicolaou, F. Ptochos , P.A. Razis , H. Rykaczewski, H. Saka **Charles University, Prague, Czech Republic**M. Finger , M. Finger Jr. , A. Kveton **Escuela Politecnica Nacional, Quito, Ecuador**E. Ayala **Universidad San Francisco de Quito, Quito, Ecuador**E. Carrera Jarrin **Academy of Scientific Research and Technology of the Arab Republic of Egypt, Egyptian Network of High Energy Physics, Cairo, Egypt**S. Elgammal¹³, A. Ellithi Kamel¹⁴**Center for High Energy Physics (CHEP-FU), Fayoum University, El-Fayoum, Egypt**A. Lotfy , M.A. Mahmoud **National Institute of Chemical Physics and Biophysics, Tallinn, Estonia**S. Bhowmik , R.K. Dewanjee , K. Ehataht , M. Kadastik, T. Lange , S. Nandan , C. Nielsen , J. Pata , M. Raidal , L. Tani , C. Veelken **Department of Physics, University of Helsinki, Helsinki, Finland**P. Eerola , H. Kirschenmann , K. Osterberg , M. Voutilainen **Helsinki Institute of Physics, Helsinki, Finland**S. Bharthuar , E. Brücken , F. Garcia , J. Havukainen , M.S. Kim , R. Kinnunen, T. Lampén , K. Lassila-Perini , S. Lehti , T. Lindén , M. Lotti, L. Martikainen , M. Myllymäki , J. Ott , M.m. Rantanen , H. Siikonen , E. Tuominen , J. Tuominiemi 

Lappeenranta-Lahti University of Technology, Lappeenranta, FinlandP. Luukka¹⁰, H. Petrow¹⁰, T. Tuuva**IRFU, CEA, Université Paris-Saclay, Gif-sur-Yvette, France**C. Amendola¹⁰, M. Besancon¹⁰, F. Couderc¹⁰, M. Dejardin¹⁰, D. Denegri, J.L. Faure, F. Ferri¹⁰, S. Ganjour¹⁰, P. Gras¹⁰, G. Hamel de Monchenault¹⁰, P. Jarry¹⁰, V. Lohezic¹⁰, J. Malcles¹⁰, J. Rander, A. Rosowsky¹⁰, M.Ö. Sahin¹⁰, A. Savoy-Navarro¹⁰¹⁵, P. Simkina¹⁰, M. Titov¹⁰**Laboratoire Leprince-Ringuet, CNRS/IN2P3, Ecole Polytechnique, Institut Polytechnique de Paris, Palaiseau, France**C. Baldenegro Barrera¹⁰, F. Beaudette¹⁰, A. Buchot Perraguin¹⁰, P. Busson¹⁰, A. Cappati¹⁰, C. Charlot¹⁰, F. Damas¹⁰, O. Davignon¹⁰, B. Diab¹⁰, G. Falmagne¹⁰, B.A. Fontana Santos Alves¹⁰, S. Ghosh¹⁰, R. Granier de Cassagnac¹⁰, A. Hakimi¹⁰, B. Harikrishnan¹⁰, G. Liu¹⁰, J. Motta¹⁰, M. Nguyen¹⁰, C. Ochando¹⁰, L. Portales¹⁰, R. Salerno¹⁰, U. Sarkar¹⁰, J.B. Sauvan¹⁰, Y. Sirois¹⁰, A. Tarabini¹⁰, E. Vernazza¹⁰, A. Zabi¹⁰, A. Zghiche¹⁰**Université de Strasbourg, CNRS, IPHC UMR 7178, Strasbourg, France**J.-L. Agram¹⁰¹⁶, J. Andrea¹⁰, D. Apparu¹⁰, D. Bloch¹⁰, G. Bourgatte¹⁰, J.-M. Brom¹⁰, E.C. Chabert¹⁰, C. Collard¹⁰, D. Darej, U. Goerlach¹⁰, C. Grimault, A.-C. Le Bihan¹⁰, P. Van Hove¹⁰**Institut de Physique des 2 Infinis de Lyon (IP2I), Villeurbanne, France**S. Beauceron¹⁰, C. Bernet¹⁰, B. Blançon¹⁰, G. Boudoul¹⁰, A. Carle, N. Chanon¹⁰, J. Choi¹⁰, D. Contardo¹⁰, P. Depasse¹⁰, C. Dozen¹⁰¹⁷, H. El Mamouni, J. Fay¹⁰, S. Gascon¹⁰, M. Gouzevitch¹⁰, G. Grenier¹⁰, B. Ille¹⁰, I.B. Laktineh, M. Lethuillier¹⁰, L. Mirabito, S. Perries, L. Torterotot¹⁰, M. Vander Donckt¹⁰, P. Verdier¹⁰, S. Viret**Georgian Technical University, Tbilisi, Georgia**D. Chokheli¹⁰, I. Lomidze¹⁰, Z. Tsamalaidze¹⁰¹²**RWTH Aachen University, I. Physikalisches Institut, Aachen, Germany**V. Botta¹⁰, L. Feld¹⁰, K. Klein¹⁰, M. Lipinski¹⁰, D. Meuser¹⁰, A. Pauls¹⁰, N. Röwert¹⁰, M. Teroerde¹⁰**RWTH Aachen University, III. Physikalisches Institut A, Aachen, Germany**S. Diekmann¹⁰, A. Dodonova¹⁰, N. Eich¹⁰, D. Eliseev¹⁰, M. Erdmann¹⁰, P. Fackeldey¹⁰, D. Fasanella¹⁰, B. Fischer¹⁰, T. Hebbeker¹⁰, K. Hoepfner¹⁰, F. Ivone¹⁰, M.y. Lee¹⁰, L. Mastrolorenzo, M. Merschmeyer¹⁰, A. Meyer¹⁰, S. Mondal¹⁰, S. Mukherjee¹⁰, D. Noll¹⁰, A. Novak¹⁰, F. Nowotny, A. Pozdnyakov¹⁰, Y. Rath, W. Redjeb¹⁰, H. Reithler¹⁰, A. Schmidt¹⁰, S.C. Schuler, A. Sharma¹⁰, L. Vigilante, S. Wiedenbeck¹⁰, S. Zaleski**RWTH Aachen University, III. Physikalisches Institut B, Aachen, Germany**C. Dziwok¹⁰, G. Flügge¹⁰, W. Haj Ahmad¹⁰¹⁸, O. Hlushchenko, T. Kress¹⁰, A. Nowack¹⁰, O. Pooth¹⁰, A. Stahl¹⁰, T. Ziemons¹⁰, A. Zott¹⁰

Deutsches Elektronen-Synchrotron, Hamburg, Germany

H. Aarup Petersen , M. Aldaya Martin , P. Asmuss, S. Baxter , M. Bayatmakou , O. Behnke , A. Bermúdez Martínez , S. Bhattacharya , A.A. Bin Anuar , F. Blekman ¹⁹, K. Borras ²⁰, D. Brunner , A. Campbell , A. Cardini , C. Cheng, F. Colombina , S. Consuegra Rodríguez , G. Correia Silva , M. De Silva , L. Didukh , G. Eckerlin, D. Eckstein , L.I. Estevez Banos , O. Filatov , E. Gallo ¹⁹, A. Geiser , A. Giraldi , G. Greau, A. Grohsjean , V. Guglielmi , M. Guthoff , A. Jafari ²¹, N.Z. Jomhari , B. Kaech , A. Kasem ²⁰, M. Kasemann , H. Kaveh , C. Kleinwort , R. Kogler , M. Komm , D. Krücker , W. Lange, D. Leyva Pernia , K. Lipka , W. Lohmann ²², R. Mankel , I.-A. Melzer-Pellmann , M. Mendizabal Morentin , J. Metwally, A.B. Meyer , G. Milella , M. Mormile , A. Mussgiller , A. Nürnberg , Y. Otarid, D. Pérez Adán , A. Raspereza , B. Ribeiro Lopes , J. Rübenach, A. Saggio , A. Saibel , M. Savitskyi , M. Scham ^{23,20}, V. Scheurer, S. Schnake ²⁰, P. Schütze , C. Schwanenberger ¹⁹, M. Shchedrolosiev , R.E. Sosa Ricardo , D. Stafford, N. Tonon †, M. Van De Klundert , F. Vazzoler , A. Ventura Barroso , R. Walsh , D. Walter , Q. Wang , Y. Wen , K. Wichmann, L. Wiens ²⁰, C. Wissing , S. Wuchterl , Y. Yang , A. Zimermann Castro Santos

University of Hamburg, Hamburg, Germany

A. Albrecht , S. Albrecht , M. Antonello , S. Bein , L. Benato , M. Bonanomi , P. Connor , K. De Leo , M. Eich, K. El Morabit , F. Feindt, A. Fröhlich, C. Garbers , E. Garutti , M. Hajheidari, J. Haller , A. Hinzmänn , H.R. Jabusch , G. Kasieczka , R. Klanner , W. Korcari , T. Kramer , V. Kutzner , J. Lange , A. Lobanov , C. Matthies , A. Mehta , L. Moureaux , M. Mrowietz, A. Nigamova , Y. Nissan, A. Paasch , K.J. Pena Rodriguez , M. Rieger , O. Rieger, P. Schleper , M. Schröder , J. Schwandt , H. Stadie , G. Steinbrück , A. Tews, M. Wolf

Karlsruhe Institut fuer Technologie, Karlsruhe, Germany

J. Bechtel , S. Brommer , M. Burkart, E. Butz , R. Caspart , T. Chwalek , A. Dierlamm , A. Droll, N. Faltermann , M. Giffels , J.O. Gosewisch, A. Gottmann , F. Hartmann ²⁴, M. Horzela , U. Husemann , P. Keicher, M. Klute , R. Koppenhöfer , S. Maier , S. Mitra , Th. Müller , M. Neukum, G. Quast , K. Rabbertz , J. Rauser, D. Savoiu , M. Schnepf, D. Seith, I. Shvetsov , H.J. Simonis , N. Trevisani , R. Ulrich , J. van der Linden , R.F. Von Cube , M. Wassmer , S. Wieland , R. Wolf , S. Wozniewski , S. Wunsch, X. Zuo

Institute of Nuclear and Particle Physics (INPP), NCSR Demokritos, Aghia Paraskevi, Greece

G. Anagnostou, P. Assiouras , G. Daskalakis , A. Kyriakis, A. Stakia 

National and Kapodistrian University of Athens, Athens, Greece

M. Diamantopoulou, D. Karasavvas, P. Kontaxakis , A. Manousakis-Katsikakis , A. Panagiotou, I. Papavergou , N. Saoulidou , K. Theofilatos , E. Tziaferi , K. Vellidis , E. Vourliotis , I. Zisopoulos 

National Technical University of Athens, Athens, Greece

G. Bakas^{ID}, T. Chatzistavrou, K. Kousouris^{ID}, I. Papakrivopoulos^{ID}, G. Tsipolitis,
A. Zacharopoulou

University of Ioánnina, Ioánnina, Greece

K. Adamidis, I. Bestintzanos, I. Evangelou^{ID}, C. Foudas, P. Gianneios^{ID}, C. Kamtsikis,
P. Katsoulis, P. Kokkas^{ID}, P.G. Kosmoglou Kiouseoglou^{ID}, N. Manthos^{ID}, I. Papadopoulos^{ID},
J. Strologas^{ID}

**MTA-ELTE Lendület CMS Particle and Nuclear Physics Group, Eötvös
Loránd University, Budapest, Hungary**

M. Csand^{ID}, K. Farkas^{ID}, M.M.A. Gadallah^{ID}²⁵, S. Lokos^{ID}²⁶, P. Major^{ID}, K. Mandal^{ID},
G. Pasztor^{ID}, A.J. Radl^{ID}²⁷, O. Suranyi^{ID}, G.I. Veres^{ID}

Wigner Research Centre for Physics, Budapest, Hungary

M. Bartok^{ID}²⁸, G. Bencze, C. Hajdu^{ID}, D. Horvath^{ID}^{29,30}, F. Sikler^{ID}, V. Veszpremi^{ID}

Institute of Nuclear Research ATOMKI, Debrecen, Hungary

N. Beni^{ID}, S. Czellar, J. Karancsi^{ID}²⁸, J. Molnar, Z. Szillasi, D. Teyssier^{ID}

Institute of Physics, University of Debrecen, Debrecen, Hungary

P. Raics, B. Ujvari^{ID}³¹

Karoly Robert Campus, MATE Institute of Technology, Gyongyos, Hungary

T. Csorgo^{ID}²⁷, F. Nemes^{ID}²⁷, T. Novak^{ID}

Panjab University, Chandigarh, India

J. Babbar^{ID}, S. Bansal^{ID}, S.B. Beri, V. Bhatnagar^{ID}, G. Chaudhary^{ID}, S. Chauhan^{ID},
N. Dhingra^{ID}³², R. Gupta, A. Kaur^{ID}, A. Kaur^{ID}, H. Kaur^{ID}, M. Kaur^{ID}, S. Kumar^{ID},
P. Kumari^{ID}, M. Meena^{ID}, K. Sandeep^{ID}, T. Sheokand, J.B. Singh^{ID}³³, A. Singla^{ID}, A. K. Virdi^{ID}

University of Delhi, Delhi, India

A. Ahmed^{ID}, A. Bhardwaj^{ID}, B.C. Choudhary^{ID}, M. Gola, A. Kumar^{ID}, M. Naimuddin^{ID},
P. Priyanka^{ID}, K. Ranjan^{ID}, S. Saumya^{ID}, A. Shah^{ID}

Saha Institute of Nuclear Physics, HBNI, Kolkata, India

S. Baradia^{ID}, S. Barman^{ID}³⁴, S. Bhattacharya^{ID}, D. Bhowmik, S. Dutta^{ID}, S. Dutta,
B. Gomber^{ID}³⁵, M. Maity³⁴, P. Palit^{ID}, P.K. Rout^{ID}, G. Saha^{ID}, B. Sahu^{ID}, S. Sarkar

Indian Institute of Technology Madras, Madras, India

P.K. Behera^{ID}, S.C. Behera^{ID}, P. Kalbhor^{ID}, J.R. Komaragiri^{ID}³⁶, D. Kumar^{ID}³⁶,
A. Muhammad^{ID}, L. Panwar^{ID}³⁶, R. Pradhan^{ID}, P.R. Pujahari^{ID}, A. Sharma^{ID}, A.K. Sikdar^{ID},
P.C. Tiwari^{ID}³⁶, S. Verma^{ID}

Bhabha Atomic Research Centre, Mumbai, India

K. Naskar^{ID}³⁷

Tata Institute of Fundamental Research-A, Mumbai, India

T. Aziz, I. Das , S. Dugad, M. Kumar , G.B. Mohanty , P. Suryadevara

Tata Institute of Fundamental Research-B, Mumbai, India

S. Banerjee , R. Chudasama , M. Guchait , S. Karmakar , S. Kumar , G. Majumder , K. Mazumdar , S. Mukherjee , A. Thachayath

National Institute of Science Education and Research, An OCC of Homi Bhabha National Institute, Bhubaneswar, Odisha, IndiaS. Bahinipati ³⁸, C. Kar , P. Mal , T. Mishra , V.K. Muraleedharan Nair Bindhu ³⁹, A. Nayak ³⁹, P. Saha , S.K. Swain , D. Vats **Indian Institute of Science Education and Research (IISER), Pune, India**

A. Alpana , S. Dube , B. Kansal , A. Laha , S. Pandey , A. Rastogi , S. Sharma

Isfahan University of Technology, Isfahan, IranH. Bakhshiansohi ^{40,41}, E. Khazaie ⁴¹, M. Zeinali ⁴²**Institute for Research in Fundamental Sciences (IPM), Tehran, Iran**S. Chenarani ⁴³, S.M. Etesami , M. Khakzad , M. Mohammadi Najafabadi **University College Dublin, Dublin, Ireland**

M. Grunewald

INFN Sezione di Bari^a, Università di Bari^b, Politecnico di Bari^c, Bari, Italy

M. Abbrescia , R. Aly , C. Aruta , A. Colaleo , D. Creanza , N. De Filippis , M. De Palma , A. Di Florio , W. Elmetenawee , F. Errico , L. Fiore , G. Iaselli , M. Ince , G. Maggi , M. Maggi , I. Margjeka , V. Mastrapasqua , S. My , S. Nuzzo , A. Pellecchia , A. Pompili , G. Pugliese , R. Radogna , D. Ramos , A. Ranieri , G. Selvaggi , L. Silvestris , F.M. Simone , Ü. Sözbilir , A. Stamerra , R. Venditti , P. Verwilligen

INFN Sezione di Bologna^a, Università di Bologna^b, Bologna, ItalyG. Abbiendi , C. Battilana , D. Bonacorsi , L. Borgonovi , L. Brigliadori , R. Campanini , P. Capiluppi , A. Castro , F.R. Cavallo , M. Cuffiani , G.M. Dallavalle , T. Diotallevi , F. Fabbri , A. Fanfani , P. Giacomelli , L. Giommi , C. Grandi , L. Guiducci , S. Lo Meo ,⁴⁴ L. Lunerti , S. Marcellini , G. Masetti , F.L. Navarria , A. Perrotta , F. Primavera , A.M. Rossi , T. Rovelli , G.P. Siroli **INFN Sezione di Catania^a, Università di Catania^b, Catania, Italy**S. Costa ,^{a,b,45} A. Di Mattia , R. Potenza , A. Tricomi ,^{a,b,45} C. Tuve **INFN Sezione di Firenze^a, Università di Firenze^b, Firenze, Italy**

G. Barbagli , B. Camaiani , A. Cassese , R. Ceccarelli , V. Ciulli , C. Civinini , R. D'Alessandro , E. Focardi , G. Latino , P. Lenzi , M. Lizzo , M. Meschini , S. Paoletti , R. Seidita , G. Sguazzoni , L. Viliani

INFN Laboratori Nazionali di Frascati, Frascati, ItalyL. Benussi , S. Bianco , S. Meola ²⁴, D. Piccolo **INFN Sezione di Genova^a, Università di Genova^b, Genova, Italy**M. Bozzo , P. Chatagnon ^a, F. Ferro ^a, R. Mulargia ^a, E. Robutti ^a, S. Tosi ^{a,b}**INFN Sezione di Milano-Bicocca^a, Università di Milano-Bicocca^b, Milano, Italy**A. Benaglia ^a, G. Boldrini ^a, F. Brivio ^{a,b}, F. Cetorelli ^{a,b}, F. De Guio ^{a,b}, M.E. Dinardo ^{a,b}, P. Dini ^a, S. Gennai ^a, A. Ghezzi ^{a,b}, P. Govoni ^{a,b}, L. Guzzi ^{a,b}, M.T. Lucchini ^{a,b}, M. Malberti ^a, S. Malvezzi ^a, A. Massironi ^a, D. Menasce ^a, L. Moroni ^a, M. Paganoni ^{a,b}, D. Pedrini ^a, B.S. Pinolini^a, S. Ragazzi ^{a,b}, N. Redaelli ^a, T. Tabarelli de Fatis ^{a,b}, D. Zuolo ^{a,b}**INFN Sezione di Napoli^a, Università di Napoli ‘Federico II’^b, Napoli, Italy; Università della Basilicata^c, Potenza, Italy; Università G. Marconi^d, Roma, Italy**S. Buontempo ^a, F. Carnevali^{a,b}, N. Cavallo ^{a,c}, A. De Iorio ^{a,b}, F. Fabozzi ^{a,c}, A.O.M. Iorio ^{a,b}, L. Lista ^{a,b,46}, P. Paolucci ^{a,24}, B. Rossi ^a, C. Sciacca ^{a,b}**INFN Sezione di Padova^a, Università di Padova^b, Padova, Italy; Università di Trento^c, Trento, Italy**P. Azzi ^a, N. Bacchetta ^{a,47}, D. Bisello ^{a,b}, P. Bortignon ^a, A. Bragagnolo ^{a,b}, P. Checchia ^a, T. Dorigo ^a, S. Fantinel ^a, F. Fanzago ^a, F. Gasparini ^{a,b}, U. Gasparini ^{a,b}, G. Grossi ^a, L. Layer^{a,48}, E. Lusiani ^a, M. Margoni ^{a,b}, A.T. Meneguzzo ^{a,b}, J. Pazzini ^{a,b}, P. Ronchese ^{a,b}, R. Rossin ^{a,b}, F. Simonetto ^{a,b}, G. Strong ^a, M. Tosi ^{a,b}, H. Yarai ^{a,b}, M. Zanetti ^{a,b}, P. Zotto ^{a,b}, A. Zucchetta ^{a,b}**INFN Sezione di Pavia^a, Università di Pavia^b, Pavia, Italy**S. Abu Zeid ^{a,49}, C. Aimè ^{a,b}, A. Braghieri ^a, S. Calzaferri ^{a,b}, D. Fiorina ^{a,b}, P. Montagna ^{a,b}, V. Re ^a, C. Riccardi ^{a,b}, P. Salvini ^a, I. Vai ^a, P. Vitulo ^{a,b}**INFN Sezione di Perugia^a, Università di Perugia^b, Perugia, Italy**P. Asenov ^{a,50}, G.M. Bilei ^a, D. Ciangottini ^{a,b}, L. Fanò ^{a,b}, M. Magherini ^{a,b}, G. Mantovani^{a,b}, V. Mariani ^{a,b}, M. Menichelli ^a, F. Moscatelli ^{a,50}, A. Piccinelli ^{a,b}, M. Presilla ^{a,b}, A. Rossi ^{a,b}, A. Santocchia ^{a,b}, D. Spiga ^a, T. Tedeschi ^{a,b}**INFN Sezione di Pisa^a, Università di Pisa^b, Scuola Normale Superiore di Pisa^c, Pisa, Italy; Università di Siena^d, Siena, Italy**P. Azzurri ^a, G. Bagliesi ^a, V. Bertacchi ^{a,c}, R. Bhattacharya ^a, L. Bianchini ^{a,b}, T. Boccali ^a, E. Bossini ^{a,b}, D. Bruschini ^{a,c}, R. Castaldi ^a, M.A. Ciocci ^{a,b}, V. D’Amante ^{a,d}, R. Dell’Orso ^a, M.R. Di Domenico ^{a,d}, S. Donato ^a, A. Giassi ^a, F. Ligabue ^{a,c}, G. Mandorli ^{a,c}, D. Matos Figueiredo ^a, A. Messineo ^{a,b}, M. Musich ^{a,b}, F. Palla ^a, S. Parolia ^{a,b}, G. Ramirez-Sanchez ^{a,c}, A. Rizzi ^{a,b}, G. Rolandi ^{a,c},

S. Roy Chowdhury ^a, T. Sarkar ^a, A. Scribano ^a, N. Shafiei ^{a,b}, P. Spagnolo ^a, R. Tenchini ^a, G. Tonelli ^{a,b}, N. Turini ^{a,d}, A. Venturi ^a, P.G. Verdini ^a

INFN Sezione di Roma^a, Sapienza Università di Roma^b, Roma, Italy

P. Barria ^a, M. Campana ^{a,b}, F. Cavallari ^a, D. Del Re ^{a,b}, E. Di Marco ^a, M. Diemoz ^a, E. Longo ^{a,b}, P. Meridiani ^a, G. Organtini ^{a,b}, F. Pandolfi ^a, R. Paramatti ^{a,b}, C. Quaranta ^{a,b}, S. Rahatlou ^{a,b}, C. Rovelli ^a, F. Santanastasio ^{a,b}, L. Soffi ^a, R. Tramontano ^{a,b}

INFN Sezione di Torino^a, Università di Torino^b, Torino, Italy; Università del Piemonte Orientale^c, Novara, Italy

N. Amapane ^{a,b}, R. Arcidiacono ^{a,c}, S. Argiro ^{a,b}, M. Arneodo ^{a,c}, N. Bartosik ^a, R. Bellan ^{a,b}, A. Bellora ^{a,b}, C. Biino ^a, N. Cartiglia ^a, M. Costa ^{a,b}, R. Covarelli ^{a,b}, N. Demaria ^a, M. Grippo ^{a,b}, B. Kiani ^{a,b}, F. Legger ^a, C. Mariotti ^a, S. Maselli ^a, A. Mecca ^{a,b}, E. Migliore ^{a,b}, E. Monteil ^{a,b}, M. Monteno ^a, M.M. Obertino ^{a,b}, G. Ortona ^a, L. Pacher ^{a,b}, N. Pastrone ^a, M. Pelliccioni ^a, M. Ruspa ^{a,c}, K. Shchelina ^a, F. Siviero ^{a,b}, V. Sola ^a, A. Solano ^{a,b}, D. Soldi ^{a,b}, A. Staiano ^a, M. Tornago ^{a,b}, D. Trocino ^a, G. Umoret ^{a,b}, A. Vagnerini ^{a,b}

INFN Sezione di Trieste^a, Università di Trieste^b, Trieste, Italy

S. Belforte ^a, V. Candelise ^{a,b}, M. Casarsa ^a, F. Cossutti ^a, A. Da Rold ^{a,b}, G. Della Ricca ^{a,b}, G. Sorrentino ^{a,b}

Kyungpook National University, Daegu, Korea

S. Dogra , C. Huh , B. Kim , D.H. Kim , G.N. Kim , J. Kim, J. Lee , S.W. Lee , C.S. Moon , Y.D. Oh , S.I. Pak , M.S. Ryu , S. Sekmen , Y.C. Yang

Chonnam National University, Institute for Universe and Elementary Particles, Kwangju, Korea

H. Kim , D.H. Moon

Hanyang University, Seoul, Korea

E. Asilar , T.J. Kim , J. Park

Korea University, Seoul, Korea

S. Choi , S. Han, B. Hong , K. Lee, K.S. Lee , J. Lim, J. Park, S.K. Park, J. Yoo

Kyung Hee University, Department of Physics, Seoul, Korea

J. Goh

Sejong University, Seoul, Korea

H. S. Kim , Y. Kim, S. Lee

Seoul National University, Seoul, Korea

J. Almond, J.H. Bhyun, J. Choi , S. Jeon , J. Kim , J.S. Kim, S. Ko , H. Kwon , H. Lee , S. Lee, B.H. Oh , M. Oh , S.B. Oh , H. Seo , U.K. Yang, I. Yoon

University of Seoul, Seoul, Korea

W. Jang , D.Y. Kang, Y. Kang , D. Kim , S. Kim , B. Ko, J.S.H. Lee , Y. Lee ,
J.A. Merlin, I.C. Park , Y. Roh, D. Song, I.J. Watson , S. Yang 

Yonsei University, Department of Physics, Seoul, Korea

S. Ha , H.D. Yoo 

Sungkyunkwan University, Suwon, Korea

M. Choi , M.R. Kim , H. Lee, Y. Lee , Y. Lee , I. Yu 

College of Engineering and Technology, American University of the Middle East (AUM), Dasman, Kuwait

T. Beyrouthy, Y. Maghrbi 

Riga Technical University, Riga, Latvia

K. Dreimanis , M. Seidel , T. Torims , V. Veckalns ⁵¹

Vilnius University, Vilnius, Lithuania

M. Ambrozas , A. Carvalho Antunes De Oliveira , A. Juodagalvis , A. Rinkevicius ,
G. Tamulaitis 

National Centre for Particle Physics, Universiti Malaya, Kuala Lumpur, Malaysia

N. Bin Norjoharuddeen , S.Y. Hoh ⁵², I. Yusuff , Z. Zolkapli

Universidad de Sonora (UNISON), Hermosillo, Mexico

J.F. Benitez , A. Castaneda Hernandez , H.A. Encinas Acosta, L.G. Gallegos Maríñez,
M. León Coello , J.A. Murillo Quijada , A. Sehrawat , L. Valencia Palomo 

Centro de Investigacion y de Estudios Avanzados del IPN, Mexico City, Mexico

G. Ayala , H. Castilla-Valdez , I. Heredia-De La Cruz ⁵³, R. Lopez-Fernandez ,
C.A. Mondragon Herrera, D.A. Perez Navarro , A. Sánchez Hernández 

Universidad Iberoamericana, Mexico City, Mexico

C. Oropeza Barrera , F. Vazquez Valencia 

Benemerita Universidad Autonoma de Puebla, Puebla, Mexico

I. Pedraza , H.A. Salazar Ibarguen , C. Uribe Estrada 

University of Montenegro, Podgorica, Montenegro

I. Bubanja, J. Mijuskovic ⁵⁴, N. Raicevic 

National Centre for Physics, Quaid-I-Azam University, Islamabad, Pakistan

A. Ahmad , M.I. Asghar, A. Awais , M.I.M. Awan, M. Gul , H.R. Hoorani , W.A. Khan ,
M. Shoaib , M. Waqas 

**AGH University of Science and Technology Faculty of Computer Science,
Electronics and Telecommunications, Krakow, Poland**

V. Avati, L. Grzanka , M. Malawski 

National Centre for Nuclear Research, Swierk, Poland

H. Bialkowska , M. Bluj , B. Boimska , M. Górski , M. Kazana , M. Szleper ,
P. Zalewski 

**Institute of Experimental Physics, Faculty of Physics, University of Warsaw,
Warsaw, Poland**

K. Bunkowski , K. Doroba , A. Kalinowski , M. Konecki , J. Krolikowski 

**Laboratório de Instrumentação e Física Experimental de Partículas, Lisboa,
Portugal**

M. Araujo , P. Bargassa , D. Bastos , A. Boletti , P. Faccioli , M. Gallinaro , J. Hollar ,
N. Leonardo , T. Niknejad , M. Pisano , J. Seixas , J. Varela 

**VINCA Institute of Nuclear Sciences, University of Belgrade, Belgrade,
Serbia**

P. Adzic , M. Dordevic , P. Milenovic , J. Milosevic 

**Centro de Investigaciones Energéticas Medioambientales y Tecnológicas
(CIEMAT), Madrid, Spain**

M. Aguilar-Benitez, J. Alcaraz Maestre , A. Álvarez Fernández , M. Barrio Luna,
Cristina F. Bedoya , C.A. Carrillo Montoya , M. Cepeda , M. Cerrada , N. Colino ,
B. De La Cruz , A. Delgado Peris , D. Fernández Del Val , J.P. Fernández Ramos ,
J. Flix , M.C. Fouz , O. Gonzalez Lopez , S. Goy Lopez , J.M. Hernandez , M.I. Josa ,
J. León Holgado , D. Moran , C. Perez Dengra , A. Pérez-Calero Yzquierdo ,
J. Puerta Pelayo , I. Redondo , D.D. Redondo Ferrero , L. Romero, S. Sánchez Navas ,
J. Sastre , L. Urda Gómez , J. Vazquez Escobar , C. Willmott

Universidad Autónoma de Madrid, Madrid, Spain

J.F. de Trocóniz 

**Universidad de Oviedo, Instituto Universitario de Ciencias y Tecnologías
Espaciales de Asturias (ICTEA), Oviedo, Spain**

B. Alvarez Gonzalez , J. Cuevas , J. Fernandez Menendez , S. Folgueras ,
I. Gonzalez Caballero , J.R. González Fernández , E. Palencia Cortezon ,
C. Ramón Álvarez , V. Rodríguez Bouza , A. Soto Rodríguez , A. Trapote ,
C. Vico Villalba 

**Instituto de Física de Cantabria (IFCA), CSIC-Universidad de Cantabria,
Santander, Spain**

J.A. Brochero Cifuentes , I.J. Cabrillo , A. Calderon , J. Duarte Campderros ,
M. Fernandez , C. Fernandez Madrazo , A. García Alonso, G. Gomez , C. Lasosa García 

C. Martinez Rivero , P. Martinez Ruiz del Arbol , F. Matorras , P. Matorras Cuevas ,
J. Piedra Gomez , C. Prieels, A. Ruiz-Jimeno , L. Scodellaro , I. Vila , J.M. Vizan Garcia 

University of Colombo, Colombo, Sri Lanka

M.K. Jayananda , B. Kailasapathy ⁵⁶, D.U.J. Sonnadara , D.D.C. Wickramarathna 

University of Ruhuna, Department of Physics, Matara, Sri Lanka

W.G.D. Dharmaratna , K. Liyanage , N. Perera , N. Wickramage 

CERN, European Organization for Nuclear Research, Geneva, Switzerland

D. Abbaneo , J. Alimena , E. Auffray , G. Auzinger , J. Baechler, P. Baillon[†], D. Barney ,
J. Bendavid , M. Bianco , B. Bilin , A. Bocci , E. Brondolin , C. Caillo ,
T. Camporesi , G. Cerminara , N. Chernyavskaya , S.S. Chhibra , S. Choudhury,
M. Cipriani , L. Cristella , D. d'Enterria , A. Dabrowski , A. David , A. De Roeck ,
M.M. Defranchis , M. Deile , M. Dobson , M. Dünser , N. Dupont, A. Elliott-Peisert,
F. Fallavollita⁵⁷, A. Florent , L. Forthomme , G. Franzoni , W. Funk , S. Ghosh ,
S. Giani, D. Gigi, K. Gill , F. Glege , L. Gouskos , E. Govorkova , M. Haranko ,
J. Hegeman , V. Innocente , T. James , P. Janot , J. Kaspar , J. Kieseler ,
N. Kratochwil , S. Laurila , P. Lecoq , E. Leutgeb , A. Lintuluoto , C. Lourenço ,
B. Maier , L. Malgeri , M. Mannelli , A.C. Marini , F. Meijers , S. Mersi , E. Meschi ,
F. Moortgat , M. Mulders , S. Orfanelli, L. Orsini, F. Pantaleo , E. Perez, M. Peruzzi ,
A. Petrilli , G. Petrucciani , A. Pfeiffer , M. Pierini , D. Piparo , M. Pitt , H. Qu ,
T. Quast, D. Rabady , A. Racz, G. Reales Gutierrez, M. Rovere , H. Sakulin ,
J. Salfeld-Nebgen , S. Scarfi , M. Selvaggi , A. Sharma , P. Silva , P. Sphicas ⁵⁸,
A.G. Stahl Leiton , S. Summers , K. Tatar , V.R. Tavolaro , D. Treille , P. Tropea ,
A. Tsirou, J. Wanczyk ⁵⁹, K.A. Wozniak , W.D. Zeuner

Paul Scherrer Institut, Villigen, Switzerland

L. Caminada ⁶⁰, A. Ebrahimi , W. Erdmann , R. Horisberger , Q. Ingram ,
H.C. Kaestli , D. Kotlinski , C. Lange , M. Missiroli ⁶⁰, L. Noehte , T. Rohe 

ETH Zurich — Institute for Particle Physics and Astrophysics (IPA), Zurich, Switzerland

T.K. Arrestad , K. Androsov ⁵⁹, M. Backhaus , P. Berger, A. Calandri , K. Datta ,
A. De Cosa , G. Dissertori , M. Dittmar, M. Donegà , F. Eble , M. Galli , K. Gedia ,
F. Glessgen , T.A. Gómez Espinosa , C. Grab , D. Hits , W. Lustermann , A.-M. Lyon ,
R.A. Manzoni , L. Marchese , C. Martin Perez , A. Mascellani ⁵⁹, M.T. Meinhard ,
F. Nessi-Tedaldi , J. Niedziela , F. Pauss , V. Perovic , S. Pigazzini , M.G. Ratti ,
M. Reichmann , C. Reissel , T. Reitenspiess , B. Ristic , F. Riti , D. Ruini,
D.A. Sanz Becerra , J. Steggemann ⁵⁹, D. Valsecchi ²⁴, R. Wallny 

Universität Zürich, Zurich, Switzerland

C. Amsler ⁶¹, P. Bärtschi , C. Botta , D. Brzhechko, M.F. Canelli , K. Cormier ,
A. De Wit , R. Del Burgo, J.K. Heikkilä , M. Huwiler , W. Jin , A. Jofrehei 

B. Kilminster^{1D}, S. Leontsinis^{1D}, S.P. Liechti^{1D}, A. Macchiolo^{1D}, P. Meiring^{1D}, V.M. Mikuni^{1D}, U. Molinatti^{1D}, I. Neutelings^{1D}, A. Reimers^{1D}, P. Robmann, S. Sanchez Cruz^{1D}, K. Schweiger^{1D}, M. Senger^{1D}, Y. Takahashi^{1D}

National Central University, Chung-Li, Taiwan

C. Adloff⁶², C.M. Kuo, W. Lin, S.S. Yu^{1D}

National Taiwan University (NTU), Taipei, Taiwan

L. Ceard, Y. Chao^{1D}, K.F. Chen^{1D}, P.s. Chen, H. Cheng^{1D}, W.-S. Hou^{1D}, R. Khurana, Y.y. Li^{1D}, R.-S. Lu^{1D}, E. Paganis^{1D}, A. Psallidas, A. Steen^{1D}, H.y. Wu, E. Yazgan^{1D}, P.r. Yu

Chulalongkorn University, Faculty of Science, Department of Physics, Bangkok, Thailand

C. Asawatangtrakuldee^{1D}, N. Srimanobhas^{1D}

Çukurova University, Physics Department, Science and Art Faculty, Adana, Turkey

D. Agyel^{1D}, F. Boran^{1D}, Z.S. Demiroglu^{1D}, F. Dolek^{1D}, I. Dumanoglu^{1D}⁶³, E. Eskut^{1D}, Y. Guler^{1D}⁶⁴, E. Gurpinar Guler^{1D}⁶⁴, C. Isik^{1D}, O. Kara, A. Kayis Topaksu^{1D}, U. Kiminsu^{1D}, G. Onengut^{1D}, K. Ozdemir^{1D}⁶⁵, A. Polatoz^{1D}, A.E. Simsek^{1D}, B. Tali^{1D}⁶⁶, U.G. Tok^{1D}, S. Turkcapar^{1D}, E. Uslan^{1D}, I.S. Zorbakir^{1D}

Middle East Technical University, Physics Department, Ankara, Turkey

G. Karapinar⁶⁷, K. Ocalan^{1D}⁶⁸, M. Yalvac^{1D}⁶⁹

Bogazici University, Istanbul, Turkey

B. Akgun^{1D}, I.O. Atakisi^{1D}, E. Gürmez^{1D}, M. Kaya^{1D}⁷⁰, O. Kaya^{1D}⁷¹, Ö. Özçelik^{1D}, S. Tekten^{1D}⁷²

Istanbul Technical University, Istanbul, Turkey

A. Cakir^{1D}, K. Cankocak^{1D}⁶³, Y. Komurcu^{1D}, S. Sen^{1D}⁶³

Istanbul University, Istanbul, Turkey

O. Aydilek^{1D}, S. Cerci^{1D}⁶⁶, B. Hacisahinoglu^{1D}, I. Hos^{1D}⁷³, B. Isildak^{1D}⁷⁴, B. Kaynak^{1D}, S. Ozkorucuklu^{1D}, C. Simsek^{1D}, D. Sunar Cerci^{1D}⁶⁶

Institute for Scintillation Materials of National Academy of Science of Ukraine, Kharkiv, Ukraine

B. Grynyov^{1D}

National Science Centre, Kharkiv Institute of Physics and Technology, Kharkiv, Ukraine

L. Levchuk^{1D}

University of Bristol, Bristol, United Kingdom

D. Anthony^{1D}, E. Bhal^{1D}, J.J. Brooke^{1D}, A. Bundock^{1D}, E. Clement^{1D}, D. Cussans^{1D}, H. Flacher^{1D}, M. Glowacki, J. Goldstein^{1D}, G.P. Heath, H.F. Heath^{1D}, L. Kreczko^{1D},

B. Krikler¹⁰, S. Paramesvaran¹⁰, S. Seif El Nasr-Storey, V.J. Smith¹⁰, N. Stylianou¹⁰⁷⁵,
K. Walkingshaw Pass, R. White¹⁰

Rutherford Appleton Laboratory, Didcot, United Kingdom

A.H. Ball, K.W. Bell¹⁰, A. Belyaev¹⁰⁷⁶, C. Brew¹⁰, R.M. Brown¹⁰, D.J.A. Cockerill¹⁰,
C. Cooke¹⁰, K.V. Ellis, K. Harder¹⁰, S. Harper¹⁰, M.-L. Holmberg¹⁰⁷⁷, J. Linacre¹⁰,
K. Manolopoulos, D.M. Newbold¹⁰, E. Olaiya, D. Petyt¹⁰, T. Reis¹⁰, G. Salvi¹⁰, T. Schuh,
C.H. Shepherd-Themistocleous¹⁰, I.R. Tomalin¹⁰, T. Williams¹⁰

Imperial College, London, United Kingdom

R. Bainbridge¹⁰, P. Bloch¹⁰, S. Bonomally, J. Borg¹⁰, S. Breeze, C.E. Brown¹⁰, O. Buchmuller,
V. Cacchio, V. Cepaitis¹⁰, G.S. Chahal¹⁰⁷⁸, D. Colling¹⁰, J.S. Dancu, P. Dauncey¹⁰,
G. Davies¹⁰, J. Davies, M. Della Negra¹⁰, S. Fayer, G. Fedi¹⁰, G. Hall¹⁰, M.H. Hassanshahi¹⁰,
A. Howard, G. Iles¹⁰, J. Langford¹⁰, L. Lyons¹⁰, A.-M. Magnan¹⁰, S. Malik, A. Martelli¹⁰,
M. Mieskolainen¹⁰, D.G. Monk¹⁰, J. Nash¹⁰⁷⁹, M. Pesaresi, B.C. Radburn-Smith¹⁰,
D.M. Raymond, A. Richards, A. Rose¹⁰, E. Scott¹⁰, C. Seez¹⁰, A. Shtipliyski, R. Shukla¹⁰,
A. Tapper¹⁰, K. Uchida¹⁰, G.P. Uttley¹⁰, L.H. Vage, T. Virdee¹⁰²⁴, M. Vojinovic¹⁰,
N. Wardle¹⁰, S.N. Webb¹⁰, D. Winterbottom¹⁰

Brunel University, Uxbridge, United Kingdom

K. Coldham, J.E. Cole¹⁰, A. Khan, P. Kyberd¹⁰, I.D. Reid¹⁰

Baylor University, Waco, Texas, U.S.A.

S. Abdullin¹⁰, A. Brinkerhoff¹⁰, B. Caraway¹⁰, J. Dittmann¹⁰, K. Hatakeyama¹⁰,
A.R. Kanuganti¹⁰, B. McMaster¹⁰, M. Saunders¹⁰, S. Sawant¹⁰, C. Sutantawibul¹⁰, J. Wilson¹⁰

Catholic University of America, Washington, DC, U.S.A.

R. Bartek¹⁰, A. Dominguez¹⁰, R. Uniyal¹⁰, A.M. Vargas Hernandez¹⁰

The University of Alabama, Tuscaloosa, Alabama, U.S.A.

A. Buccilli¹⁰, S.I. Cooper¹⁰, D. Di Croce¹⁰, S.V. Gleyzer¹⁰, C. Henderson¹⁰, C.U. Perez¹⁰,
P. Rumerio¹⁰⁸⁰, C. West¹⁰

Boston University, Boston, Massachusetts, U.S.A.

A. Akpinar¹⁰, A. Albert¹⁰, D. Arcaro¹⁰, C. Cosby¹⁰, Z. Demiragli¹⁰, C. Erice¹⁰, E. Fontanesi¹⁰,
D. Gastler¹⁰, S. May¹⁰, J. Rohlf¹⁰, K. Salyer¹⁰, D. Sperka¹⁰, D. Spitzbart¹⁰, I. Suarez¹⁰,
A. Tsatsos¹⁰, S. Yuan¹⁰

Brown University, Providence, Rhode Island, U.S.A.

G. Benelli¹⁰, B. Burkle¹⁰, X. Coubez²⁰, D. Cutts¹⁰, M. Hadley¹⁰, U. Heintz¹⁰, J.M. Hogan¹⁰⁸¹,
T. Kwon¹⁰, G. Landsberg¹⁰, K.T. Lau¹⁰, D. Li¹⁰, J. Luo¹⁰, M. Narain¹⁰, N. Pervan¹⁰,
S. Sagir¹⁰⁸², F. Simpson¹⁰, E. Usai¹⁰, W.Y. Wong, X. Yan¹⁰, D. Yu¹⁰, W. Zhang

University of California, Davis, Davis, California, U.S.A.

J. Bonilla¹⁰, C. Brainerd¹⁰, R. Breedon¹⁰, M. Calderon De La Barca Sanchez¹⁰, M. Chertok¹⁰,
J. Conway¹⁰, P.T. Cox¹⁰, R. Erbacher¹⁰, G. Haza¹⁰, F. Jensen¹⁰, O. Kukral¹⁰, G. Mocellin¹⁰,
M. Mulhearn¹⁰, D. Pellett¹⁰, B. Regnery¹⁰, D. Taylor¹⁰, Y. Yao¹⁰, F. Zhang¹⁰

University of California, Los Angeles, California, U.S.A.

M. Bachtis , R. Cousins , A. Datta , D. Hamilton , J. Hauser , M. Ignatenko ,
 M.A. Iqbal , T. Lam , E. Manca , W.A. Nash , S. Regnard , D. Saltzberg , B. Stone ,
 V. Valuev

University of California, Riverside, Riverside, California, U.S.A.

Y. Chen, R. Clare , J.W. Gary , M. Gordon, G. Hanson , G. Karapostoli , O.R. Long ,
 N. Manganello , W. Si , S. Wimpenny

University of California, San Diego, La Jolla, California, U.S.A.

J.G. Branson , P. Chang , S. Cittolin , S. Cooperstein , D. Diaz , J. Duarte ,
 R. Gerosa , L. Giannini , J. Guiang , R. Kansal , V. Krutelyov , R. Lee , J. Letts ,
 M. Masciovecchio , F. Mokhtar , M. Pieri , B.V. Sathia Narayanan , V. Sharma ,
 M. Tadel , F. Würthwein , Y. Xiang , A. Yagil

University of California, Santa Barbara — Department of Physics, Santa Barbara, California, U.S.A.

N. Amin, C. Campagnari , M. Citron , G. Collura , A. Dorsett , V. Dutta ,
 J. Incandela , M. Kilpatrick , J. Kim , A.J. Li , P. Masterson , H. Mei , M. Oshiro ,
 M. Quinnan , J. Richman , U. Sarica , R. Schmitz , F. Setti , J. Sheplock ,
 P. Siddireddy, D. Stuart , S. Wang

California Institute of Technology, Pasadena, California, U.S.A.

A. Bornheim , O. Cerri, I. Dutta , J.M. Lawhorn , N. Lu , J. Mao , H.B. Newman ,
 T. Q. Nguyen , M. Spiropulu , J.R. Vlimant , C. Wang , S. Xie , R.Y. Zhu

Carnegie Mellon University, Pittsburgh, Pennsylvania, U.S.A.

J. Alison , S. An , M.B. Andrews , P. Bryant , T. Ferguson , A. Harilal , C. Liu ,
 T. Mudholkar , S. Murthy , M. Paulini , A. Roberts , A. Sanchez , W. Terrill

University of Colorado Boulder, Boulder, Colorado, U.S.A.

J.P. Cumalat , W.T. Ford , A. Hassani , G. Karathanasis , E. MacDonald, F. Marini ,
 R. Patel, A. Perloff , C. Savard , N. Schonbeck , K. Stenson , K.A. Ulmer ,
 S.R. Wagner , N. Zipper

Cornell University, Ithaca, New York, U.S.A.

J. Alexander , S. Bright-Thonney , X. Chen , D.J. Cranshaw , J. Fan , X. Fan ,
 D. Gadkari , S. Hogan , J. Monroy , J.R. Patterson , D. Quach , J. Reichert , M. Reid ,
 A. Ryd , J. Thom , P. Wittich , R. Zou

Fermi National Accelerator Laboratory, Batavia, Illinois, U.S.A.

M. Albrow , M. Alyari , G. Apollinari , A. Apresyan , L.A.T. Bauerdi , D. Berry ,
 J. Berryhill , P.C. Bhat , K. Burkett , J.N. Butler , A. Canepa , G.B. Cerati ,
 H.W.K. Cheung , F. Chlebana , K.F. Di Petrillo , J. Dickinson , V.D. Elvira , Y. Feng ,
 J. Freeman , A. Gandrakota , Z. Gecse , L. Gray , D. Green, S. Grünendahl

O. Gutsche^{ID}, R.M. Harris^{ID}, R. Heller^{ID}, T.C. Herwig^{ID}, J. Hirschauer^{ID}, L. Horyn^{ID},
 B. Jayatilaka^{ID}, S. Jindariani^{ID}, M. Johnson^{ID}, U. Joshi^{ID}, T. Klijnsma^{ID}, B. Klima^{ID},
 K.H.M. Kwok^{ID}, S. Lammel^{ID}, D. Lincoln^{ID}, R. Lipton^{ID}, T. Liu^{ID}, C. Madrid^{ID},
 K. Maeshima^{ID}, C. Mantilla^{ID}, D. Mason^{ID}, P. McBride^{ID}, P. Merkel^{ID}, S. Mrenna^{ID}, S. Nahn^{ID},
 J. Ngadiuba^{ID}, D. Noonan^{ID}, V. Papadimitriou^{ID}, N. Pastika^{ID}, K. Pedro^{ID}, C. Pena^{ID}⁸³,
 F. Ravera^{ID}, A. Reinsvold Hall^{ID}⁸⁴, L. Ristori^{ID}, E. Sexton-Kennedy^{ID}, N. Smith^{ID}, A. Soha^{ID},
 L. Spiegel^{ID}, J. Strait^{ID}, L. Taylor^{ID}, S. Tkaczyk^{ID}, N.V. Tran^{ID}, L. Uplegger^{ID},
 E.W. Vaandering^{ID}, H.A. Weber^{ID}, I. Zoi^{ID}

University of Florida, Gainesville, Florida, U.S.A.

P. Avery^{ID}, D. Bourilkov^{ID}, L. Cadamuro^{ID}, V. Cherepanov^{ID}, R.D. Field, D. Guerrero^{ID},
 M. Kim, E. Koenig^{ID}, J. Konigsberg^{ID}, A. Korytov^{ID}, K.H. Lo, K. Matchev^{ID}, N. Menendez^{ID},
 G. Mitselmakher^{ID}, A. Muthirakalayil Madhu^{ID}, N. Rawal^{ID}, D. Rosenzweig^{ID}, S. Rosenzweig^{ID},
 K. Shi^{ID}, J. Wang^{ID}, Z. Wu^{ID}

Florida State University, Tallahassee, Florida, U.S.A.

T. Adams^{ID}, A. Askew^{ID}, R. Habibullah^{ID}, V. Hagopian^{ID}, T. Kolberg^{ID}, G. Martinez,
 H. Prosper^{ID}, C. Schiber, O. Viazlo^{ID}, R. Yohay^{ID}, J. Zhang

Florida Institute of Technology, Melbourne, Florida, U.S.A.

M.M. Baarmand^{ID}, S. Butalla^{ID}, T. Elkafrawy^{ID}⁴⁹, M. Hohlmann^{ID}, R. Kumar Verma^{ID},
 M. Rahmani, F. Yumiceva^{ID}

University of Illinois at Chicago (UIC), Chicago, Illinois, U.S.A.

M.R. Adams^{ID}, H. Becerril Gonzalez^{ID}, R. Cavanaugh^{ID}, S. Dittmer^{ID}, O. Evdokimov^{ID},
 C.E. Gerber^{ID}, D.J. Hofman^{ID}, D. S. Lemos^{ID}, A.H. Merrit^{ID}, C. Mills^{ID}, G. Oh^{ID}, T. Roy^{ID},
 S. Rudrabhatla^{ID}, M.B. Tonjes^{ID}, N. Varelas^{ID}, X. Wang^{ID}, Z. Ye^{ID}, J. Yoo^{ID}

The University of Iowa, Iowa City, Iowa, U.S.A.

M. Alhusseini^{ID}, K. Dilsiz^{ID}⁸⁵, L. Emediato^{ID}, R.P. Gandrajula^{ID}, G. Karaman^{ID},
 O.K. Köseyan^{ID}, J.-P. Merlo, A. Mestvirishvili^{ID}⁸⁶, J. Nachtman^{ID}, O. Neogi, H. Ogul^{ID}⁸⁷,
 Y. Onel^{ID}, A. Penzo^{ID}, C. Snyder, E. Tiras^{ID}⁸⁸

Johns Hopkins University, Baltimore, Maryland, U.S.A.

O. Amram^{ID}, B. Blumenfeld^{ID}, L. Corcodilos^{ID}, J. Davis^{ID}, A.V. Gritsan^{ID}, S. Kyriacou^{ID},
 P. Maksimovic^{ID}, J. Roskes^{ID}, S. Sekhar^{ID}, M. Swartz^{ID}, T.Á. Vámi^{ID}

The University of Kansas, Lawrence, Kansas, U.S.A.

A. Abreu^{ID}, L.F. Alcerro Alcerro^{ID}, J. Anguiano^{ID}, P. Baringer^{ID}, A. Bean^{ID}, Z. Flowers^{ID},
 T. Isidori^{ID}, S. Khalil^{ID}, J. King^{ID}, G. Krintiras^{ID}, M. Lazarovits^{ID}, C. Le Mahieu^{ID}, C. Lindsey,
 J. Marquez^{ID}, N. Minafra^{ID}, M. Murray^{ID}, M. Nickel^{ID}, C. Rogan^{ID}, C. Royon^{ID}, R. Salvatico^{ID},
 S. Sanders^{ID}, C. Smith^{ID}, Q. Wang^{ID}, J. Williams^{ID}, G. Wilson^{ID}

Kansas State University, Manhattan, Kansas, U.S.A.

B. Allmond^{ID}, S. Duric, A. Ivanov^{ID}, K. Kaadze^{ID}, D. Kim, Y. Maravin^{ID}, T. Mitchell,
 A. Modak, K. Nam, D. Roy^{ID}

Lawrence Livermore National Laboratory, Livermore, California, U.S.A.F. Rebassoo^{ID}, D. Wright^{ID}**University of Maryland, College Park, Maryland, U.S.A.**E. Adams^{ID}, A. Baden^{ID}, O. Baron, A. Belloni^{ID}, A. Bethani^{ID}, S.C. Eno^{ID}, N.J. Hadley^{ID}, S. Jabeen^{ID}, R.G. Kellogg^{ID}, T. Koeth^{ID}, Y. Lai^{ID}, S. Lascio^{ID}, A.C. Mignerey^{ID}, S. Nabili^{ID}, C. Palmer^{ID}, C. Papageorgakis^{ID}, L. Wang^{ID}, K. Wong^{ID}**Massachusetts Institute of Technology, Cambridge, Massachusetts, U.S.A.**D. Abercrombie, W. Busza^{ID}, I.A. Cali^{ID}, Y. Chen^{ID}, M. D’Alfonso^{ID}, J. Eysermans^{ID}, C. Freer^{ID}, G. Gomez-Ceballos^{ID}, M. Goncharov, P. Harris, M. Hu^{ID}, D. Kovalskyi^{ID}, J. Krupa^{ID}, Y.-J. Lee^{ID}, K. Long^{ID}, C. Mironov^{ID}, C. Paus^{ID}, D. Rankin^{ID}, C. Roland^{ID}, G. Roland^{ID}, Z. Shi^{ID}, G.S.F. Stephans^{ID}, J. Wang, Z. Wang^{ID}, B. Wyslouch^{ID}**University of Minnesota, Minneapolis, Minnesota, U.S.A.**R.M. Chatterjee, A. Evans^{ID}, J. Hiltbrand^{ID}, Sh. Jain^{ID}, B.M. Joshi^{ID}, C. Kapsiak^{ID}, M. Krohn^{ID}, Y. Kubota^{ID}, J. Mans^{ID}, M. Revering^{ID}, R. Rusack^{ID}, R. Saradhy^{ID}, N. Schroeder^{ID}, N. Strobbe^{ID}, M.A. Wadud^{ID}**University of Mississippi, Oxford, Mississippi, U.S.A.**L.M. Cremaldi^{ID}**University of Nebraska-Lincoln, Lincoln, Nebraska, U.S.A.**K. Bloom^{ID}, M. Bryson, D.R. Claes^{ID}, C. Fangmeier^{ID}, L. Finco^{ID}, F. Golf^{ID}, C. Joo^{ID}, R. Kamaliuddin, I. Kravchenko^{ID}, I. Reed^{ID}, J.E. Siado^{ID}, G.R. Snow[†], W. Tabb^{ID}, A. Wightman^{ID}, F. Yan^{ID}, A.G. Zecchinelli^{ID}**State University of New York at Buffalo, Buffalo, New York, U.S.A.**G. Agarwal^{ID}, H. Bandyopadhyay^{ID}, L. Hay^{ID}, I. Iashvili^{ID}, A. Kharchilava^{ID}, C. McLean^{ID}, M. Morris^{ID}, D. Nguyen^{ID}, J. Pekkanen^{ID}, S. Rappoccio^{ID}, A. Williams^{ID}**Northeastern University, Boston, Massachusetts, U.S.A.**G. Alverson^{ID}, E. Barberis^{ID}, Y. Haddad^{ID}, Y. Han^{ID}, A. Krishna^{ID}, J. Li^{ID}, J. Lidrych^{ID}, G. Madigan^{ID}, B. Marzocchi^{ID}, D.M. Morse^{ID}, V. Nguyen^{ID}, T. Orimoto^{ID}, A. Parker^{ID}, L. Skinnari^{ID}, A. Tishelman-Charny^{ID}, T. Wamorkar^{ID}, B. Wang^{ID}, A. Wisecarver^{ID}, D. Wood^{ID}**Northwestern University, Evanston, Illinois, U.S.A.**S. Bhattacharya^{ID}, J. Bueghly, Z. Chen^{ID}, A. Gilbert^{ID}, K.A. Hahn^{ID}, Y. Liu^{ID}, N. Odell^{ID}, M.H. Schmitt^{ID}, M. Velasco**University of Notre Dame, Notre Dame, Indiana, U.S.A.**R. Band^{ID}, R. Bucci, M. Cremonesi, A. Das^{ID}, R. Goldouzian^{ID}, M. Hildreth^{ID}, K. Hurtado Anampa^{ID}, C. Jessop^{ID}, K. Lannon^{ID}, J. Lawrence^{ID}, N. Loukas^{ID}, L. Lutton^{ID}, J. Mariano, N. Marinelli, I. Mcalister, T. McCauley^{ID}, C. Mcgrady^{ID}, K. Mohrman^{ID}, C. Moore^{ID}, Y. Musienko^{ID}¹², R. Ruchti^{ID}, A. Townsend^{ID}, M. Wayne^{ID}, H. Yockey, M. Zarucki^{ID}, L. Zygal^{ID}

The Ohio State University, Columbus, Ohio, U.S.A.

B. Bylsma, M. Carrigan¹⁰, L.S. Durkin¹⁰, B. Francis¹⁰, C. Hill¹⁰, A. Lesauvage¹⁰,
 M. Nunez Ornelas¹⁰, K. Wei, B.L. Winer¹⁰, B. R. Yates¹⁰

Princeton University, Princeton, New Jersey, U.S.A.

F.M. Addesa¹⁰, P. Das¹⁰, G. Dezoort¹⁰, P. Elmer¹⁰, A. Frankenthal¹⁰, B. Greenberg¹⁰,
 N. Haubrich¹⁰, S. Higginbotham¹⁰, A. Kalogeropoulos¹⁰, G. Kopp¹⁰, S. Kwan¹⁰, D. Lange¹⁰,
 D. Marlow¹⁰, K. Mei¹⁰, I. Ojalvo¹⁰, J. Olsen¹⁰, D. Stickland¹⁰, C. Tully¹⁰

University of Puerto Rico, Mayaguez, Puerto Rico, U.S.A.

S. Malik¹⁰, S. Norberg

Purdue University, West Lafayette, Indiana, U.S.A.

A.S. Bakshi¹⁰, V.E. Barnes¹⁰, R. Chawla¹⁰, S. Das¹⁰, L. Gutay, M. Jones¹⁰, A.W. Jung¹⁰,
 D. Kondratyev¹⁰, A.M. Koshy, M. Liu¹⁰, G. Negro¹⁰, N. Neumeister¹⁰, G. Paspalaki¹⁰,
 S. Piperov¹⁰, A. Purohit¹⁰, J.F. Schulte¹⁰, M. Stojanovic¹⁰¹⁵, J. Thieman¹⁰, F. Wang¹⁰,
 R. Xiao¹⁰, W. Xie¹⁰

Purdue University Northwest, Hammond, Indiana, U.S.A.

J. Dolen¹⁰, N. Parashar¹⁰

Rice University, Houston, Texas, U.S.A.

D. Acosta¹⁰, A. Baty¹⁰, T. Carnahan¹⁰, M. Decaro, S. Dildick¹⁰, K.M. Ecklund¹⁰,
 P.J. Fernández Manteca¹⁰, S. Freed, P. Gardner, F.J.M. Geurts¹⁰, A. Kumar¹⁰, W. Li¹⁰,
 B.P. Padley¹⁰, R. Redjimi, J. Rotter¹⁰, W. Shi¹⁰, S. Yang¹⁰, E. Yigitbasi¹⁰, L. Zhang⁸⁹,
 Y. Zhang¹⁰

University of Rochester, Rochester, New York, U.S.A.

A. Bodek¹⁰, P. de Barbaro¹⁰, R. Demina¹⁰, J.L. Dulemba¹⁰, C. Fallon, T. Ferbel¹⁰, M. Galanti,
 A. Garcia-Bellido¹⁰, O. Hindrichs¹⁰, A. Khukhunaishvili¹⁰, E. Ranken¹⁰, R. Taus¹⁰,
 G.P. Van Onsem¹⁰

The Rockefeller University, New York, New York, U.S.A.

K. Goulianatos¹⁰

Rutgers, The State University of New Jersey, Piscataway, New Jersey, U.S.A.

B. Chiarito, J.P. Chou¹⁰, Y. Gershtein¹⁰, E. Halkiadakis¹⁰, A. Hart¹⁰, M. Heindl¹⁰,
 D. Jaroslawski¹⁰, O. Karacheban¹⁰²², I. Laflotte¹⁰, A. Lath¹⁰, R. Montalvo, K. Nash,
 M. Osherson¹⁰, S. Salur¹⁰, S. Schnetzer, S. Somalwar¹⁰, R. Stone¹⁰, S.A. Thayil¹⁰, S. Thomas,
 H. Wang¹⁰

University of Tennessee, Knoxville, Tennessee, U.S.A.

H. Acharya, A.G. Delannoy¹⁰, S. Fiorendi¹⁰, T. Holmes¹⁰, E. Nibigira¹⁰, S. Spanier¹⁰

Texas A&M University, College Station, Texas, U.S.A.

O. Bouhalil⁹⁰, M. Dalchenko^{ID}, A. Delgado^{ID}, R. Eusebi^{ID}, J. Gilmore^{ID}, T. Huang^{ID}, T. Kamon^{ID}⁹¹, H. Kim^{ID}, S. Luo^{ID}, S. Malhotra, R. Mueller^{ID}, D. Overton^{ID}, D. Rathjens^{ID}, A. Safonov^{ID}

Texas Tech University, Lubbock, Texas, U.S.A.

N. Akchurin^{ID}, J. Damgov^{ID}, V. Hegde^{ID}, K. Lamichhane^{ID}, S.W. Lee^{ID}, T. Mengke, S. Muthumuni^{ID}, T. Peltola^{ID}, I. Volobouev^{ID}, Z. Wang, A. Whitbeck^{ID}

Vanderbilt University, Nashville, Tennessee, U.S.A.

E. Appelt^{ID}, S. Greene, A. Gurrola^{ID}, W. Johns^{ID}, A. Melo^{ID}, F. Romeo^{ID}, P. Sheldon^{ID}, S. Tuo^{ID}, J. Velkovska^{ID}, J. Viinikainen^{ID}

University of Virginia, Charlottesville, Virginia, U.S.A.

B. Cardwell^{ID}, B. Cox^{ID}, G. Cummings^{ID}, J. Hakala^{ID}, R. Hirosky^{ID}, M. Joyce^{ID}, A. Ledovskoy^{ID}, A. Li^{ID}, C. Neu^{ID}, C.E. Perez Lara^{ID}, B. Tannenwald^{ID}

Wayne State University, Detroit, Michigan, U.S.A.

P.E. Karchin^{ID}, N. Poudyal^{ID}

University of Wisconsin — Madison, Madison, Wisconsin, U.S.A.

S. Banerjee^{ID}, K. Black^{ID}, T. Bose^{ID}, S. Dasu^{ID}, I. De Bruyn^{ID}, P. Everaerts^{ID}, C. Galloni, H. He^{ID}, M. Herndon^{ID}, A. Herve^{ID}, C.K. Koraka^{ID}, A. Lanaro, A. Loeliger^{ID}, R. Loveless^{ID}, J. Madhusudanan Sreekala^{ID}, A. Mallampalli^{ID}, A. Mohammadi^{ID}, S. Mondal, G. Parida^{ID}, D. Pinna, A. Savin, V. Shang^{ID}, V. Sharma^{ID}, W.H. Smith^{ID}, D. Teague, H.F. Tsoi^{ID}, W. Vetens^{ID}

Authors affiliated with an institute or an international laboratory covered by a cooperation agreement with CERN

S. Afanasiev^{ID}, V. Andreev^{ID}, Yu. Andreev^{ID}, T. Aushev^{ID}, M. Azarkin^{ID}, A. Babaev^{ID}, A. Belyaev^{ID}, V. Blinov⁹², E. Boos^{ID}, V. Borshch^{ID}, D. Budkouski^{ID}, V. Bunichev^{ID}, O. Bychkova, M. Chadeeva^{ID}⁹², V. Chekhovsky, A. Dermenev^{ID}, T. Dimova^{ID}⁹², I. Dremin^{ID}, M. Dubinin^{ID}⁸³, L. Dudko^{ID}, V. Epshteyn^{ID}, A. Ershov^{ID}, G. Gavrilov^{ID}, V. Gavrilov^{ID}, S. Gninenko^{ID}, V. Golovtcov^{ID}, N. Golubev^{ID}, I. Golutvin^{ID}, I. Gorbunov^{ID}, A. Gribushin^{ID}, V. Ivanchenko^{ID}, Y. Ivanov^{ID}, V. Kachanov^{ID}, L. Kardapoltsev^{ID}⁹², V. Karjavine^{ID}, A. Karneyeu^{ID}, V. Kim^{ID}⁹², M. Kirakosyan, D. Kirpichnikov^{ID}, M. Kirsanov^{ID}, V. Klyukhin^{ID}, O. Kodolova^{ID}⁹³, D. Konstantinov^{ID}, V. Korenkov^{ID}, A. Kozyrev^{ID}⁹², N. Krasnikov^{ID}, E. Kuznetsova^{ID}⁹⁴, A. Lanev^{ID}, P. Levchenko^{ID}, A. Litomin, N. Lychkovskaya^{ID}, V. Makarenko^{ID}, A. Malakhov^{ID}, V. Matveev^{ID}⁹², V. Murzin^{ID}, A. Nikitenko^{ID}⁹⁵, S. Obraztsov^{ID}, V. Okhotnikov^{ID}, I. Ovtin^{ID}⁹², V. Palichik^{ID}, P. Parygin^{ID}, V. Perelygin^{ID}, M. Perfilov, G. Pivovarov^{ID}, V. Popov, E. Popova^{ID}, O. Radchenko^{ID}⁹², V. Rusinov, M. Savina^{ID}, V. Savrin^{ID}, D. Selivanova^{ID}, V. Shalaev^{ID}, S. Shmatov^{ID}, S. Shulha^{ID}, Y. Skovpen^{ID}⁹², S. Slabospitskii^{ID}, V. Smirnov^{ID}, D. Sosnov^{ID}, A. Stepennov^{ID}, V. Sulimov^{ID}, E. Tcherniaev^{ID}, A. Terkulov^{ID}, O. Teryaev^{ID}, I. Tlisova^{ID}, M. Toms^{ID}, A. Toropin^{ID},

L. Uvarov¹⁰, A. Uzunian¹⁰, E. Vlasov¹⁰, A. Vorobyev, N. Voytishin¹⁰, B.S. Yuldashev⁹⁶,
A. Zarubin¹⁰, I. Zhizhin¹⁰, A. Zhokin¹⁰

[†] Deceased

¹ Also at TU Wien, Vienna, Austria

² Also at Institute of Basic and Applied Sciences, Faculty of Engineering, Arab Academy for Science, Technology and Maritime Transport, Alexandria, Egypt

³ Also at Université Libre de Bruxelles, Bruxelles, Belgium

⁴ Also at Universidade Estadual de Campinas, Campinas, Brazil

⁵ Also at Federal University of Rio Grande do Sul, Porto Alegre, Brazil

⁶ Also at UFMS, Nova Andradina, Brazil

⁷ Also at The University of the State of Amazonas, Manaus, Brazil

⁸ Also at University of Chinese Academy of Sciences, Beijing, China

⁹ Also at Nanjing Normal University, Nanjing, China

¹⁰ Now at The University of Iowa, Iowa City, Iowa, U.S.A.

¹¹ Also at University of Chinese Academy of Sciences, Beijing, China

¹² Also at an institute or an international laboratory covered by a cooperation agreement with CERN

¹³ Now at British University in Egypt, Cairo, Egypt

¹⁴ Now at Cairo University, Cairo, Egypt

¹⁵ Also at Purdue University, West Lafayette, Indiana, U.S.A.

¹⁶ Also at Université de Haute Alsace, Mulhouse, France

¹⁷ Also at Department of Physics, Tsinghua University, Beijing, China

¹⁸ Also at Erzincan Binali Yildirim University, Erzincan, Turkey

¹⁹ Also at University of Hamburg, Hamburg, Germany

²⁰ Also at RWTH Aachen University, III. Physikalisches Institut A, Aachen, Germany

²¹ Also at Isfahan University of Technology, Isfahan, Iran

²² Also at Brandenburg University of Technology, Cottbus, Germany

²³ Also at Forschungszentrum Jülich, Juelich, Germany

²⁴ Also at CERN, European Organization for Nuclear Research, Geneva, Switzerland

²⁵ Also at Physics Department, Faculty of Science, Assiut University, Assiut, Egypt

²⁶ Also at Karoly Robert Campus, MATE Institute of Technology, Gyongyos, Hungary

²⁷ Also at Wigner Research Centre for Physics, Budapest, Hungary

²⁸ Also at Institute of Physics, University of Debrecen, Debrecen, Hungary

²⁹ Also at Institute of Nuclear Research ATOMKI, Debrecen, Hungary

³⁰ Now at Universitatea Babes-Bolyai — Facultatea de Fizica, Cluj-Napoca, Romania

³¹ Also at Faculty of Informatics, University of Debrecen, Debrecen, Hungary

³² Also at Punjab Agricultural University, Ludhiana, India

³³ Also at UPES — University of Petroleum and Energy Studies, Dehradun, India

³⁴ Also at University of Visva-Bharati, Santiniketan, India

³⁵ Also at University of Hyderabad, Hyderabad, India

³⁶ Also at Indian Institute of Science (IISc), Bangalore, India

³⁷ Also at Indian Institute of Technology (IIT), Mumbai, India

³⁸ Also at IIT Bhubaneswar, Bhubaneswar, India

³⁹ Also at Institute of Physics, Bhubaneswar, India

⁴⁰ Also at Deutsches Elektronen-Synchrotron, Hamburg, Germany

⁴¹ Now at Department of Physics, Isfahan University of Technology, Isfahan, Iran

⁴² Also at Sharif University of Technology, Tehran, Iran

⁴³ Also at Department of Physics, University of Science and Technology of Mazandaran, Behshahr, Iran

⁴⁴ Also at Italian National Agency for New Technologies, Energy and Sustainable Economic

Development, Bologna, Italy

⁴⁵ Also at Centro Siciliano di Fisica Nucleare e di Struttura Della Materia, Catania, Italy

⁴⁶ Also at Scuola Superiore Meridionale, Università di Napoli ‘Federico II’, Napoli, Italy

⁴⁷ Also at Fermi National Accelerator Laboratory, Batavia, Illinois, U.S.A.

- ⁴⁸ Also at Università di Napoli ‘Federico II’, Napoli, Italy
- ⁴⁹ Also at Ain Shams University, Cairo, Egypt
- ⁵⁰ Also at Consiglio Nazionale delle Ricerche — Istituto Officina dei Materiali, Perugia, Italy
- ⁵¹ Also at Riga Technical University, Riga, Latvia
- ⁵² Also at Department of Applied Physics, Faculty of Science and Technology, Universiti Kebangsaan Malaysia, Bangi, Malaysia
- ⁵³ Also at Consejo Nacional de Ciencia y Tecnología, Mexico City, Mexico
- ⁵⁴ Also at IRFU, CEA, Université Paris-Saclay, Gif-sur-Yvette, France
- ⁵⁵ Also at Faculty of Physics, University of Belgrade, Belgrade, Serbia
- ⁵⁶ Also at Trincomalee Campus, Eastern University, Sri Lanka, Nilaveli, Sri Lanka
- ⁵⁷ Also at INFN Sezione di Pavia, Università di Pavia, Pavia, Italy
- ⁵⁸ Also at National and Kapodistrian University of Athens, Athens, Greece
- ⁵⁹ Also at Ecole Polytechnique Fédérale Lausanne, Lausanne, Switzerland
- ⁶⁰ Also at Universität Zürich, Zurich, Switzerland
- ⁶¹ Also at Stefan Meyer Institute for Subatomic Physics, Vienna, Austria
- ⁶² Also at Laboratoire d’Annecy-le-Vieux de Physique des Particules, IN2P3-CNRS, Annecy-le-Vieux, France
- ⁶³ Also at Near East University, Research Center of Experimental Health Science, Mersin, Turkey
- ⁶⁴ Also at Konya Technical University, Konya, Turkey
- ⁶⁵ Also at Izmir Bakircay University, Izmir, Turkey
- ⁶⁶ Also at Adiyaman University, Adiyaman, Turkey
- ⁶⁷ Also at Istanbul Gedik University, Istanbul, Turkey
- ⁶⁸ Also at Necmettin Erbakan University, Konya, Turkey
- ⁶⁹ Also at Bozok Üniversitesi Rektörlüğü, Yozgat, Turkey
- ⁷⁰ Also at Marmara University, Istanbul, Turkey
- ⁷¹ Also at Milli Savunma University, Istanbul, Turkey
- ⁷² Also at Kafkas University, Kars, Turkey
- ⁷³ Also at Istanbul University — Cerrahpasa, Faculty of Engineering, Istanbul, Turkey
- ⁷⁴ Also at Yildiz Technical University, Istanbul, Turkey
- ⁷⁵ Also at Vrije Universiteit Brussel, Brussel, Belgium
- ⁷⁶ Also at School of Physics and Astronomy, University of Southampton, Southampton, United Kingdom
- ⁷⁷ Also at University of Bristol, Bristol, United Kingdom
- ⁷⁸ Also at IPPP Durham University, Durham, United Kingdom
- ⁷⁹ Also at Monash University, Faculty of Science, Clayton, Australia
- ⁸⁰ Also at Università di Torino, Torino, Italy
- ⁸¹ Also at Bethel University, St. Paul, Minnesota, U.S.A.
- ⁸² Also at Karamanoğlu Mehmetbey University, Karaman, Turkey
- ⁸³ Also at California Institute of Technology, Pasadena, California, U.S.A.
- ⁸⁴ Also at United States Naval Academy, Annapolis, Maryland, U.S.A.
- ⁸⁵ Also at Bingöl University, Bingöl, Turkey
- ⁸⁶ Also at Georgian Technical University, Tbilisi, Georgia
- ⁸⁷ Also at Sinop University, Sinop, Turkey
- ⁸⁸ Also at Erciyes University, Kayseri, Turkey
- ⁸⁹ Also at Institute of Modern Physics and Key Laboratory of Nuclear Physics and Ion-beam Application (MOE) — Fudan University, Shanghai, China
- ⁹⁰ Also at Texas A&M University at Qatar, Doha, Qatar
- ⁹¹ Also at Kyungpook National University, Daegu, Korea
- ⁹² Also at another institute or international laboratory covered by a cooperation agreement with CERN
- ⁹³ Also at Yerevan Physics Institute, Yerevan, Armenia
- ⁹⁴ Also at University of Florida, Gainesville, Florida, U.S.A.
- ⁹⁵ Also at Imperial College, London, United Kingdom
- ⁹⁶ Also at Institute of Nuclear Physics of the Uzbekistan Academy of Sciences, Tashkent, Uzbekistan