

RECEIVED: September 27, 2023

ACCEPTED: November 30, 2023

PUBLISHED: December 12, 2023

Search for direct production of GeV-scale resonances decaying to a pair of muons in proton-proton collisions at $\sqrt{s} = 13$ TeV



The CMS collaboration

E-mail: cms-publication-committee-chair@cern.ch

ABSTRACT: A search for direct production of low-mass dimuon resonances is performed using $\sqrt{s} = 13$ TeV proton-proton collision data collected by the CMS experiment during the 2017–2018 operation of the CERN LHC with an integrated luminosity of 96.6 fb^{-1} . The search exploits a dedicated high-rate trigger stream that records events with two muons with transverse momenta as low as 3 GeV but does not include the full event information. The search is performed by looking for narrow peaks in the dimuon mass spectrum in the ranges of 1.1–2.6 GeV and 4.2–7.9 GeV. No significant excess of events above the expectation from the standard model background is observed. Model-independent limits on production rates of dimuon resonances within the experimental fiducial acceptance are set. Competitive or world’s best limits are set at 90% confidence level for a minimal dark photon model and for a scenario with two Higgs doublets and an extra complex scalar singlet (2HDM+S). Values of the squared kinetic mixing coefficient ε^2 in the dark photon model above 10^{-6} are excluded over most of the mass range of the search. In the 2HDM+S, values of the mixing angle $\sin(\theta_H)$ above 0.08 are excluded over most of the mass range of the search with a fixed ratio of the Higgs doublets vacuum expectation $\tan\beta = 0.5$.

KEYWORDS: Beyond Standard Model, Hadron-Hadron Scattering

ARXIV EPRINT: [2309.16003](https://arxiv.org/abs/2309.16003)

Contents

1	Introduction	1
2	The CMS detector	3
3	Triggers and data sets	3
4	Event simulation	4
5	Event selection	5
6	Signal and background model	8
7	Systematic uncertainties	10
8	Results	11
9	Summary	14
	The CMS collaboration	19

1 Introduction

Searches for resonant particle pair production in accelerator experiments have played a significant role in the development of the standard model (SM). The observation of resonant production of electron-positron pairs and muon-antimuon pairs signaled the discovery of the charm [1, 2] and bottom [3] quarks, respectively. The Z boson was discovered through the observation of resonant production of both electron-positron and muon-antimuon pairs [4]. More recently, findings of resonant production of photon pairs and Z boson pairs were central to the discovery of the Higgs boson at the CERN LHC by the ATLAS [5] and CMS [6, 7] Collaborations. Searches for new particles at the TeV scale are an important component of the LHC physics program. Resonant searches in dilepton channels are a large part of this effort, however, they have so far yielded results consistent with SM predictions [8, 9]. Interest has therefore grown in extending resonance searches to explore the phase space of lower masses and very small effective couplings to SM particles, which is becoming possible thanks to the large amount of data delivered by the LHC. The GeV scale is particularly interesting because a new vector boson in that mass range could provide an explanation for several astrophysical observations [10].

This paper describes a search for production of a narrow resonance that decays promptly to a pair of oppositely charged muons using proton-proton (pp) collision data

recorded by the CMS experiment. The explored mass ranges are 1.1–2.6 GeV and 4.2–7.9 GeV. These ranges exclude the regions around the J/ψ , $\psi(2S)$, and $\Upsilon(1S)$ resonances. Previous limits on dimuon resonance production in these mass ranges have been set by the BaBar [11] and LHCb [12–14] Collaborations. The latter also probes the mass range below 1 GeV and searches for long-lived particles, which are not covered in this paper. This search is an extension of a previous CMS analysis for higher masses [15], with trigger selection and muon identification optimized for the very low-mass region. It assumes that the natural width of the resonance is much smaller than the detector dimuon mass resolution ($\approx 1.3\%$), and that its decay is consistent with originating from the primary pp collision vertex within the instrumental resolution. The sensitivity of this analysis is enhanced by the use of high-rate (“scouting”) triggers [16] that select a larger fraction of signal-like events as compared with the standard CMS triggers, by recording only a small part of the event information, consisting of trigger objects as opposed to the full detector readout. The previous CMS analysis at higher masses utilized both the standard CMS triggers and the scouting triggers, while this search uses the scouting triggers exclusively.

We also provide an interpretation of the result in the context of two specific beyond the SM scenarios. The first scenario involves the introduction of a new gauge field $U(1)_D$ that undergoes kinetic mixing with the SM hypercharge gauge field $U(1)_Y$ [17, 18]. Such mixing, controlled by the parameter ϵ , results in a dark photon (Z_D), a possibly massive, neutral vector boson. In the mass ranges probed by this search, it is coupled dominantly to the electric charges of SM particles with a coupling strength suppressed by a factor ϵ . It is assumed to be produced via the Drell-Yan (DY) mechanism, which is the dominant production mechanism in the mass range of interest [19]. The second scenario is a two Higgs doublet model with an extra complex scalar singlet (2HDM+S) [20]. For certain model parameters [21], this scenario can accommodate a light pseudoscalar boson (a) that can decay into a dimuon pair. In this model, the couplings of the light pseudoscalar a to the SM particles are determined by its mixing with the Higgs doublets and parameterized by the mixing angle (θ_H) and the ratio of the Higgs doublet vacuum expectation values ($\tan\beta$). It is assumed that the light pseudoscalar a is predominantly produced via gluon-gluon fusion (ggF). The interpretation is provided in the specific 2HDM+S scenario of type IV with $\tan\beta = 0.5$ given the enhanced coupling to the up-type quarks, predicted in this model. The same model is targeted by the LHCb experiment [14] because it represents the most sensitive one in experiments at pp colliders and allows meaningful bounds to be set on the sine of the mixing angle θ_H . Example Feynman diagrams for the production of both particles are presented in figure 1.

This paper is organized as follows. We begin with the CMS detector description in section 2 and elaborate on the relevant triggers (section 3) and the simulated data samples used in the analysis (section 4). Sections 5 and 6 detail the event selection and the signal and background modeling, respectively. Section 7 explains the systematic uncertainties and the results are presented in section 8, followed by a summary in section 9. Tabulated results are provided in the HEPData record for this analysis [22].




Figure 1. Expected dominant production modes for beyond-SM particles: a dark photon Z_D (left) and light pseudoscalar boson a (right).

2 The CMS detector

The central feature of the CMS apparatus is a superconducting solenoid of 6 m internal diameter, providing a magnetic field of 3.8 T. Within the solenoid volume are a silicon pixel and strip tracker, and both a lead tungstate crystal electromagnetic and a brass/scintillator calorimeter, which consist of a barrel and two endcap sections. Forward calorimeters extend the pseudorapidity η coverage provided by the barrel and endcap detectors. Muons are detected in gas-ionization chambers embedded in the steel flux-return yoke outside the solenoid. A more detailed description of the CMS detector, together with a definition of the coordinate system used and the relevant kinematic variables, is given in ref. [23].

Muons are measured in the pseudorapidity range $|\eta| < 2.4$, with detection planes made using three technologies: drift tubes, cathode strip chambers, and resistive plate chambers. Matching muons to tracks measured in the silicon tracker results in a relative transverse momentum (p_T) resolution of 1% in the barrel and 3% in the endcaps, for muons with p_T up to 100 GeV.

3 Triggers and data sets

Events of interest are selected using a two-tiered trigger system [24], the level-1 (L1) and high-level (HLT) triggers. The L1, composed of custom hardware processors, uses information from the calorimeters and muon detectors to select events at a rate of around 100 kHz. The HLT consists of a farm of CPU processors running a version of the CMS full event reconstruction software optimized for fast event data processing. Events passing the full set of HLT selection criteria are saved for permanent storage at an overall rate of about 1 kHz.

A set of four different L1 trigger algorithms are utilized in this analysis, defined by requiring the presence of at least two muons with various additional requirements. The simplest of these (1) only requires two muons with $p_T > 15$ GeV and > 7 GeV, respectively. The remaining three triggers all share the requirement that the two muons be oppositely charged, together with the additional conditions: (2) each muon has $|\eta| < 2.0$ and $p_T > 4.5$ GeV, and that the dimuon mass ($m_{\mu\mu}$) is between 7–18 GeV; or (3) each muon has $p_T > 4.0$ (4.5) GeV in 2017 (2018) and the two muons must have an angular separation $\Delta R = \sqrt{(\Delta\eta)^2 + (\Delta\phi)^2} < 1.2$ and ϕ is the azimuthal angle; or (4) both muons have

L1 path	p_T [GeV]	$ \eta $	ΔR	$m_{\mu\mu}$ [GeV]	Charge
(1)	$>15/7$	—	—	—	—
(2)	>4.5	<2.0	—	7–18	Opposite sign
(3)	>4.0 (2017)/ 4.5 (2018)	—	<1.2	—	Opposite sign
(4)	—	<1.5	<1.4	—	Opposite sign

Table 1. The set of dimuon L1 requirements applied in the high-rate triggers.

$|\eta| < 1.5$ and are separated by $\Delta R < 1.4$, with no restriction on p_T . These criteria are summarized in table 1.

At the HLT, our analysis makes use of the CMS high-rate data scouting trigger system [16] in which the information associated with the selected events is much reduced, permitting the recording of larger data samples. This technique has been used in previous resonant searches performed by the CMS experiment including dimuon [15] and hadronic [25] final states. Events are directly reconstructed in the HLT and are required to have at least two muons with $p_T > 3$ GeV. The dimuon events selected by the data scouting trigger are written to disk with an event size of about 4 (8) kB for data taken in 2017 (2018). The stored information includes tracking hits, calorimeter clusters, muon hits, and trigger primitives, along with their reconstructed quantities. By contrast the size of a standard raw event is closer to 1 MB. These dimuon scouting triggers were fully commissioned during 2017 [26] and recorded events at a rate of approximately 2 kHz at the peak instantaneous luminosity of $2 \times 10^{34} \text{ cm}^{-2} \text{s}^{-1}$, whereas the standard dimuon triggers recorded approximately 0.45 kHz at the same instantaneous luminosity. Data corresponding to an integrated luminosity of 96.6 fb^{-1} were collected with these triggers in 2017–2018. Figure 2 shows the efficiency of the dimuon scouting triggers in conjunction with the four L1 triggers for both data periods. This is measured using an independent CMS data set collected with a trigger requiring large energy deposits in the electromagnetic calorimeter. This data set is agnostic to the presence or absence of muons in the event and therefore is not correlated with the activation of the dimuon scouting trigger. By sampling this data set for events that contain pairs of muons, we assemble an unbiased set of dimuon events to gauge the performance of the dimuon scouting trigger. For this sample of unbiased muon pairs, we count the number that also triggered the dimuon scouting path and at least one of the four L1 triggers, and compare the result to the total number of reconstructed dimuon events in that data set that pass our fiducial selections. The ratio of these two counts is the efficiency of our trigger. The trigger efficiency in 2018 is higher than that of 2017, due to updates in the HLT muon reconstruction algorithm for the 2018 data and efficiency loss in the later part of 2017 caused by a powering problem affecting part of the pixel detector [27].

4 Event simulation

The analysis employs an estimation of the SM background based on recorded data. However, simulations of certain SM processes are used for the validation of selection criteria,




Figure 2. The measured 2017 (left) and 2018 (right) efficiencies of the dimuon scouting trigger and logical OR of all L1 triggers using 2017–2018 data. Each cell value represents the probability that a pair of muons meeting the trigger requirements will activate the dimuon scouting trigger. The x -axis shows the dimuon mass and includes the entire relevant range for this analysis. The y -axis shows the angular separation, ΔR , between the two muons. Statistical uncertainty on the value of each cell is less than 5%.

corrections to data driven efficiency measurements, as well as the corresponding systematic uncertainties in the signal predictions. These processes include DY production of dimuon pairs in the low $m_{\mu\mu}$ region ($1 < m_{\mu\mu} < 10$ GeV), as well as resonant dimuon production via J/ψ and $\Upsilon(1S)$ mesons. These processes are simulated at leading order (LO) in perturbative quantum chromodynamics with PYTHIA 8.230 [28]. In the 2HDM+S scenario, the production of a BSM pseudoscalar Higgs via ggF, with modified couplings to the up-type and down-type quarks, is simulated at LO with PYTHIA 8.230 and is used as a proxy for the production of a light pseudoscalar of the type IV model. The Z_D signal is simulated at next-to-LO in perturbative quantum chromodynamics using the hidden Abelian Higgs model [19, 29, 30] implemented in the MADGRAPH5_aMC@NLO v3.4.1 generator [31], with up to two additional partons in the matrix element calculations. The parton distribution function (PDF) set is NNPDF3.1 in the four-flavor scheme [32].

For all simulated samples, the parton showering and hadronization processes are performed with PYTHIA 8.230 using the CP5 underlying event tune [33]. Events are passed through a full detector simulation performed with GEANT4 [34]. The simulated samples include additional pp interactions per bunch crossing, referred to as “pileup”. The effect of pileup is taken into account by generating minimum-bias collision events with PYTHIA. The simulated events are weighted such that the distribution of the number of pileup interactions matches that in data.

5 Event selection

A complete event reconstruction is performed by the HLT. The muons in the scouting data stream are reconstructed by combining the information from the silicon tracker and muon systems, and their momentum is determined from the curvature of the muon track. The

muon momentum obtained with the HLT event reconstruction agrees with that obtained with the full CMS offline event reconstruction within 1–2%.

The event candidate is required to have at least one primary vertex (PV) reconstructed by the HLT system and to contain a pair of oppositely charged muons originating from this vertex. The absolute transverse distance of the PV to the beam axis, referred to as the vertex transverse displacement L_{xy} , is required to be less than 0.2 cm. Both muons are required to have $p_T > 4 \text{ GeV}$ and $|\eta| < 1.9$. The restriction on η is imposed to ensure optimal dimuon mass resolution without incurring a significant loss in acceptance. The muons are required to satisfy a muon identification based on multivariate analysis (MVA) techniques. Two MVA discriminants are used depending on the reconstructed $m_{\mu\mu}$, optimized for the signal kinematic properties in each mass range. The MVA identification utilizes information on the quality of the muon tracks, the relative isolation of the muon, and the vertex that the muons are associated with. The muon track quality parameters include the number of hits in the pixel detector, number of contributing layers in the strip tracker, and χ^2 of the muon track fit. The relative isolation of the muon is computed as the scalar p_T sum of reconstructed tracks in a cone of $\Delta R = 0.3$ around the muon candidate, divided by the muon p_T . The dimuon vertex information includes the χ^2 of the common vertex fit and L_{xy} . These parameters are combined into a single discriminator using a boosted decision tree (BDT) [35]. The choice of input parameters to the BDT and the requirement on the discriminator are optimized to achieve the highest expected sensitivity to signal-like events.

The mass range 2.6–4.2 GeV is excluded because of the presence of the J/ψ and $\psi(2S)$ resonances. The MVA identification for $m_{\mu\mu} < 2.6 \text{ GeV}$ is trained using a partial J/ψ sample, whereas the identification for $m_{\mu\mu} > 4.2 \text{ GeV}$ is trained using an $\Upsilon(1S)$ sample. The efficiency of the muon MVA identification is determined from J/ψ and Υ events using the “tag-and-probe technique” [36]. Background samples enriched with nonprompt and misidentified muons for the BDT training are extracted from the data events with only one pair of same-sign muons and a mass of $3 < m_{\mu\mu} < 10 \text{ GeV}$.

At smaller values of $m_{\mu\mu}$, the dimuon system is more Lorentz boosted and the muons become more collinear, resulting in larger uncertainties in the reconstructed PV position. The L_{xy} distribution becomes broader, whereas the distribution of the vertex displacement significance remains stable. Therefore, we employ different displacement criteria depending on the mass value. For events with $m_{\mu\mu} < 2.6 \text{ GeV}$ we further require that L_{xy} divided by its uncertainty ($\sigma_{L_{xy}}$) be less than 3.5, and the L_{xy} value is not used in the BDT. Events with $m_{\mu\mu} > 4.2 \text{ GeV}$ are required to have $L_{xy} < 0.015 \text{ cm}$ to further suppress nonprompt backgrounds. Figure 3 shows the $m_{\mu\mu}$ distribution obtained from the dimuon scouting triggers. The average efficiency of the $\Upsilon(1S)$ (J/ψ)-trained muon MVA identification is around 85 (90)% estimated from the $\Upsilon(1S)$ (J/ψ) resonance. The $\Upsilon(1S)$ -trained muon MVA identification exhibits better performance for $m_{\mu\mu} > 4.2 \text{ GeV}$, as it suppresses more background events, while maintaining high signal efficiency. Compared with the simple cut-based selection used in the previous CMS search [15], the MVA identification provides a higher efficiency while still keeping the misidentification rate low, resulting in improvements to the final limits on the cross section by about 30%. We refer to this as the inclusive selection.




Figure 3. The $m_{\mu\mu}$ distribution obtained with the muon scouting data collected during 2017–2018 with two sets of selections: the $\Upsilon(1S)$ -trained muon MVA identification with $L_{xy} < 0.015$ cm (blue, solid), and the J/ψ -trained muon MVA identification with $L_{xy}/\sigma_{L_{xy}} < 3.5$ (red, dashed). The J/ψ -trained and $\Upsilon(1S)$ -trained selections are used in the mass range 1.1–2.6 and 4.2–7.9 GeV, respectively.

Preselection	$L_{xy} < 0.2$ cm, $ \eta^\mu < 1.9$, Opposite sign			
Signal selection	Inclusive			high- p_T
Mass	$m_{\mu\mu} < 4$ GeV	$m_{\mu\mu} > 4$ GeV	$m_{\mu\mu} < 4$ GeV	$m_{\mu\mu} > 4$ GeV
p_T^μ		>4 GeV		>5 GeV
Muon	J/ψ ID	Υ ID	J/ψ ID	J/ψ ID
Vertex	$L_{xy}/\sigma_{L_{xy}} < 3.5$	$L_{xy} < 0.015$ cm	$L_{xy}/\sigma_{L_{xy}} < 3.5$	$L_{xy}/\sigma_{L_{xy}} < 3.5$
$p_T^{\mu\mu}$	—	—	>35 GeV	>20 GeV

Table 2. Summary of all selection criteria for an event to enter the analysis in inclusive and high- p_T search categories.

In the search for a light pseudoscalar particle (the 2HDM+S scenario), the signal is produced via the ggF process, which results in a harder p_T distribution compared to the DY process, due to the enhanced probability for a jet to scatter off of an initial state gluon. Higher values of muon p_T are thus required to further improve the sensitivity for the pseudoscalar signal. The muons are required to have $p_T > 5$ GeV and the muon pair is required to have $p_T > 20$ (35) GeV for $m_{\mu\mu} > 4.2$ GeV (< 2.6 GeV). The J/ψ -trained muon MVA identification and $L_{xy}/\sigma_{L_{xy}} < 3.5$ are required for both $m_{\mu\mu} < 2.6$ GeV and $m_{\mu\mu} > 4.2$ GeV. We refer to this as the high- p_T selection. All selection criteria for both categories are summarized in table 2.




Figure 4. The signal acceptance and reconstruction efficiency are extracted from the dark photon (purple, dark) and pseudoscalar (cyan, light) simulations. The region at 2.6–4.2 GeV is excluded because of the presence of the J/ψ and $\psi(2\text{S})$ resonances.

The muon reconstruction and identification efficiency measured from data and the acceptance extracted from signal simulations are shown in figure 4. The efficiency of the vertex displacement requirement of the signal is corrected using the J/ψ and $\Upsilon(1\text{S})$ samples from data and simulation. The total identification efficiency of the event is the product of the identification efficiencies of the two muons.

6 Signal and background model

The signal is extracted by performing simultaneous signal plus background maximum likelihood fits to the $m_{\mu\mu}$ distribution. For each mass hypothesis, the fit is performed over a mass window spanning ± 5 times the mass resolution around the nominal resonance mass. The mass resolution is determined from data to be roughly 1.3% across the entire mass range. This window size ensures sufficient sidebands for the background model fit, such that it will not be influenced by the presence of a signal at the central value. When fitting the mass distributions in the high- p_T selection category, the size of the window is increased to ± 8 times the mass resolution, to offset the fact that the number of events in the sideband is significantly reduced by the high p_T requirements. The probability density function (pdf) chosen to model the signal distribution is the sum of a double Crystal Ball [37, 38] and a Gaussian distribution. We limit the search to the case of narrow resonances with an intrinsic width much smaller than the experimental resolution. Since this same approxi-

mation holds for charmonium and Y resonances, the shape parameters for the signal pdf are determined from least square mass fits for the J/ψ , $\psi(2S)$, and $Y((1S), (2S), \text{ and } (3S))$ resonances. The signal pdf parameters, including the mass resolution, are found to have little dependence on the resonance mass or the selection categories. The only free parameter of the signal pdf in the final fit is the overall normalization, which is proportional to the production cross section of the new particle.

Several possible functions to model the background are investigated. We use empirical functions because the shape of the background in the search region is complicated by the presence of multiple known resonances and by the acceptance of the various triggers. The families of functions considered are Bernstein polynomials [39], a polynomial times an exponential, a sum of exponentials, and a sum of a Bernstein polynomial and a power law function. The optimal order for each function family is chosen by performing a Fisher F -test [40] with background-only fits to the dimuon mass distribution for each mass hypothesis. The order of the function family is chosen as the lowest one that keeps the number of fits with an F value above the critical value below the false-rejection probability ($\alpha < 0.05$). Across all the considered families, the optimal order for the polynomial component is always 4th order or lower. For the sum-of-exponentials function, the optimal number of terms is found to be 7. The potential bias of each function family is evaluated using simulations assuming either no signal or the presence of a small hypothetical signal. The bias for each function type, defined as the difference between the assumed and measured signal strength, is less than $\approx 20\%$ of the statistical uncertainty in the signal strength. Therefore, the added uncertainty due to the bias of the background function is negligible and is not included in the final fit. The discrete profiling method [41] is used to further suppress the bias, which arises from the choice of the background function, by allowing the signal shape to vary between the options listed above during the maximum likelihood fit. The background model fit of the mass distribution becomes unreliable when the tails of J/ψ and $\psi(2S)$ resonances enter the mass window, so the mass range 2.6–4.2 GeV is excluded from the search.

Among all the known resonances, the D^0 meson contributes an apparent peak in the $m_{\mu\mu}$ continuum in the search range. When the pions and kaons from the $D^0 \rightarrow K^+K^-$ or $D^0 \rightarrow K^-\pi^+$ decays are misreconstructed as muons, the $m_{\mu\mu}$ distribution exhibits an enhancement at 1.72 or 1.58 GeV, respectively. The contamination of these processes in the signal region is estimated from a control region by a simultaneous fit to both regions. The control region is defined by requiring an $L_{xy}/\sigma_{L_{xy}}$ to be within 3.5–11.0. The pdf of the $D^0 \rightarrow K^+K^- (K^-\pi^+)$ dimuon peak is derived from simulation. The normalization is a free parameter that is constrained by a transfer factor between the signal and control regions from the simulation. The uncertainty in the transfer factor includes the statistical uncertainty of the simulated samples and the mismodeling of $L_{xy}/\sigma_{L_{xy}}$, which is quantified using the J/ψ peak in both data and simulation. The J/ψ events in the data are determined by subtracting the contribution from background and nonprompt J/ψ production, estimated through extrapolation from the distribution, where $8 < L_{xy}/\sigma_{L_{xy}} < 20$. The contribution from the $D^0 \rightarrow \pi^+\pi^-$ process is not modeled because pions are much less likely to be misidentified as muons compared to kaons and there are insufficient $D^0 \rightarrow \pi^+\pi^-$ yields in the control region to properly constrain the process.

7 Systematic uncertainties

The signal extraction in this analysis is affected by several sources of systematic uncertainties. The measured integrated luminosity has an uncertainty of 2.3 (2.5)% for data collected in 2017 (2018) [42–44]. Roughly half of these values are attributed to correlated systematic effects between the years, and the remaining portion is uncorrelated, giving a combined uncertainty of about 2.0% between the two years. The uncertainty in the efficiency of the PV displacement requirement is 3% for $m_{\mu\mu} > 4.2 \text{ GeV}$ and is negligible for $m_{\mu\mu} < 2.6 \text{ GeV}$.

The efficiency of the dimuon scouting triggers is measured with respect to fully reconstructed muons using events selected with independent standard triggers. The uncertainties in the trigger efficiency depend on the dimuon mass and vary from 1–16%, as determined from the difference between the trigger efficiency in data and simulation in each mass bin.

The efficiency of muon identification is measured with respect to trigger requirements. The uncertainties in the identification efficiency of the two muons vary from 4–20%, and include the statistical uncertainty of the measurement sample and the uncertainty associated with the modeling in the input parameters of the muon MVA identification between the data and simulation. The uncertainty is largest for muons with low p_T and low ΔR because of the reduced number of muons in this region in the simulated samples.

The per muon identification efficiency is applied to calculate the total reconstruction efficiency. The identification efficiency is measured from the resonances produced mainly through the gluon splitting process, though the signal production may proceed via other mechanisms. The identification efficiencies between the two muons are correlated because event observables are used in the MVA identification, whereas the efficiencies are treated as uncorrelated when calculating the total efficiency. To account for the efficiency variation caused by these effects, an additional 4 (8)% systematic uncertainty is assigned for $m_{\mu\mu} > 4.2 \text{ GeV}$ ($< 2.6 \text{ GeV}$). The uncertainty in the transfer factor that connects the normalizations between the signal and control regions of the $D^0 \rightarrow K^+K^- (K^-\pi^+)$ backgrounds is 20 (25)%.

The mass resolution of the signal is measured using different resonances in data. The variation in the observed mass resolutions is around 20% and is assigned as the systematic uncertainty. The signal shape uncertainty, which varies from 1.3–2.7%, is derived by using a double Gaussian as the signal template in the fit. As described in section 6, the uncertainty due to the bias of all of the candidate background functions is negligible. The implicit bias associated with having chosen empirical functions to model the background is accounted for via the implementation of the discrete profiling method. This methodology introduces an additional discrete nuisance parameter, associated with the choice of background function, into the maximum likelihood fit. Table 3 summarizes the systematic uncertainties in the search.

The theoretical uncertainties in the signal models are not considered in the model-independent search since the propagated kinematic variations exert a negligible influence on the reconstruction efficiencies. Likewise, these uncertainties are not included in the model-dependent limits, to factorize the theoretical and experimental uncertainties.

Effect	$m_{\mu\mu} < 2.6 \text{ GeV}$	$m_{\mu\mu} > 4.2 \text{ GeV}$
Integrated luminosity		2.3–2.5%
Mass resolution		20%
Signal shape	1.3–2.7%	1.4–1.7%
Trigger efficiency		1–16%
Muon ID efficiency	4–9%	12–20%
Vertex selection	—	3%
Efficiency application	8%	4%
D^0 meson normalization	20–25%	—

Table 3. Summary of the experimental systematic uncertainties for a signal model in the model independent search for a dimuon resonance.

8 Results

The data are consistent with the continuum dimuon production. Limits at 95% confidence level (CL) are set on the product of the signal cross section, the branching fraction to a muon pair, and fiducial acceptance of a hypothetical resonance with a mass from 1.1–2.6 GeV or from 4.2–7.9 GeV. The computation of exclusion upper limits is performed using the CL_s criterion in the asymptotic approximation [45–47]. A profile likelihood ratio test statistic is built including the systematic uncertainties presented above as nuisance parameters. A log-normal pdf is used to describe each constrained nuisance parameter.

Figure 5 shows the expected and observed model independent limits for the inclusive and high- p_T selections. The 95% confidence level (CL) upper limit for the inclusive selection is of order 0.10 (0.15) pb in the mass range 1.1–2.6 (4.2–7.9) GeV. The 95% CL upper limit for the high- p_T selection is about 0.015 (0.024) pb in the mass range 1.1–2.6 (4.2–7.9) GeV. The discontinuities in the expected limits arise from the discrete profiling method applied when different background functions are favored during the fitting process. The upward deviations in the expected limits below 2 GeV are due to the peaking background associated with D^0 meson decays. The excess at 1.867 GeV in the inclusive selection may correspond to the $D^0 \rightarrow \pi^+\pi^-$ or $D^0 \rightarrow \mu^+\mu^-$ processes. Distinguishing between these two possibilities is beyond the scope of this paper. The observed limits exhibit oscillatory structures within some mass ranges, which result from statistical fluctuations modulated by the sliding mass window. No significant excess is observed beyond 2.6 (0.6) standard deviations local (global) significance from the SM background expectation in the high- p_T category. The model independent limits are exploited to constrain the dark photon [17, 18, 48] and 2HDM+S [20] scenarios.

In the first case, limits on the squared kinetic mixing coefficient ε^2 are extracted as a function of the dark photon mass (m_{Z_D}) according to the following relation:

$$\sigma_{pp \rightarrow Z_D} \varepsilon^2 \mathcal{B}A = \sigma_{\text{limit}}, \quad (8.1)$$




Figure 5. Left: expected and observed model-independent upper limits at 95% CL on the product of the signal cross section (σ), the branching fraction to a pair of muons for the inclusive dimuon selection (\mathcal{B}), and fiducial acceptance (A). Right: the model-independent limits for the high- p_T selection. The mass region dominated by the J/ψ and $\psi(2S)$ resonances is excluded from the search.

where $\sigma_{pp \rightarrow Z_D}$ is the dark photon cross section with $\varepsilon = 1$, \mathcal{B} is the branching fraction of $Z_D \rightarrow \mu^+ \mu^-$, σ_{limit} is the model independent limit, and A is the acceptance. We use the numerical software DYTURBO [49] to generate the DY process at next-to-next-to-LO (NNLO) with the renormalization and factorization scales set to $\mu = 0.5\sqrt{m_{\mu\mu}^2 + p_{T\mu\mu}^2}$ and to correct the dark photon cross section and extract the acceptance, because the dark photon is produced through the DY-like process. The interpretation of the cross section limit as a limit on the mixing coefficient ε^2 is subject to theoretical uncertainties. Uncertainties in the theoretical cross section are found by varying the renormalization and factorization scales up and down by a factor of 2, and the uncertainty in the acceptance is found by comparing the results between the DYTURBO and MADGRAPH5_aMC@NLO generators. The scale variations are between 3.5 and 9.7%, whereas the difference in acceptance is up to 31%. Upper limits at 90% CL on the mixing coefficient ε^2 are presented in figure 6, together with the LHCb and BaBar limits reinterpreted by DARKCAST [50]. The sensitivity becomes better at low masses because of the larger DY production cross section at lower energy scale. The kinetic mixing coefficient ε reflects the degree of mixing and the strength of the coupling of dark photon to other SM particles. The mixing coefficient ε^2 above $2(9) \times 10^{-7}$ is excluded at $m_{Z_D} = 2(7)$ GeV. These results complement the previous CMS search at larger masses [15].

In the 2HDM+S scenarios, the limits from the high- p_T selection are converted to an upper limit on the mixing angle θ_H as a function of the light pseudoscalar-boson mass. The recasting of the limit is performed under the assumption of $\tan \beta = 0.5$ according to the following relation:

$$\sigma_{pp \rightarrow a} \sin^2(\theta_H) \mathcal{B} A = \sigma_{\text{limit}}, \quad (8.2)$$

where $\sigma_{pp \rightarrow a}$ is the production cross section of the light pseudoscalar a , σ_{limit} is the model independent limit, \mathcal{B} is the branching fraction of $a \rightarrow \mu^+ \mu^-$, and A is the acceptance. The $\sigma_{pp \rightarrow a}$ is computed from the HIGLU generator at NNLO using the NNPDF3.0 PDF set [32] and assuming renormalization and factorization scales $\mu = 0.5m_a$, as in the LHCb




Figure 6. Observed upper limits at 90% CL on the square of the kinetic mixing coefficient ε in the minimal model of a dark photon from the CMS search in the mass ranges of 1.1–2.6 GeV and 4.2–7.9 GeV (pink). The CMS limits are compared with the existing limits at 90% CL provided by LHCb [13] (blue) and BaBar [11] (gray). The LHCb exclusion limit also extends below 1 GeV and to smaller couplings through its search for long-lived signals.




Figure 7. Observed upper limits at 90% CL on the mixing angle θ_H for the 2HDM+S scenario from the CMS search in the mass ranges of 1.1–2.6 GeV and 4.2–7.9 GeV (pink). The CMS limits are compared with the existing limits at 90% CL provided by LHCb [14] (blue) and BaBar [11] (gray).

search [14]. Uncertainties in the theoretical cross section for pseudoscalar production via ggF are found by varying the renormalization and factorization scales by a factor of 2. These uncertainties are around 90% at $m_a = 1.18$ GeV and gradually reduce to 10% at $m_a > 4.2$ GeV. The uncertainty in the ggF acceptance is estimated by comparing the values obtained with MADGRAPH5_aMC@NLO and PYTHIA, and is about 30%.

Observed upper limits at 90% CL are presented in figure 7. Values of $\sin(\theta_H)$ above ≈ 0.01 (0.02) are excluded at $m_a = 2$ (7) GeV with fixed $\tan \beta = 0.5$. The limits derived from this search in the low-mass region are competitive with recently reported results from the LHCb experiment [14] below the charmonium peaks and better above them.

9 Summary

A search for direct production of a narrow resonance decaying to a pair of muons has been presented using proton-proton collision data recorded by the CMS experiment at $\sqrt{s} = 13\text{ TeV}$ in 2017–2018. The search is performed in the dimuon mass regions of 1.1–2.6 GeV and 4.2–7.9 GeV using data collected in a dedicated high-rate trigger stream, corresponding to an integrated luminosity of 96.6 fb^{-1} . A multivariate analysis method is used to identify muons to achieve a higher sensitivity. No significant excess of events above the expectation from the standard model background is observed. Model-independent limits on production rates of dimuon resonances within the experimental fiducial acceptance are set. Competitive or world’s best limits are set at 90% confidence level for a minimal dark photon model and for a scenario with two Higgs doublets and an extra complex scalar singlet (2HDM+S). Values of the squared kinetic mixing coefficient ε^2 in the dark photon model above 10^{-6} are excluded over most of the mass range of the search. In the 2HDM+S, values of the mixing angle $\sin(\theta_H)$ above 0.08 are excluded over most of the mass range of the search with a fixed ratio of the Higgs doublets vacuum expectation $\tan\beta = 0.5$.

Acknowledgments

We thank Stefano Camarda for suggesting the DYTURBO settings for calculating the DY cross section at low mass, for the dark photon interpretation. We also thank Philip Ilten for providing the details of the $\text{pp} \rightarrow a$ cross section calculation using the HIGLU generator, for the comparison of CMS and LHCb results.

We congratulate our colleagues in the CERN accelerator departments for the excellent performance of the LHC and thank the technical and administrative staffs at CERN and at other CMS institutes for their contributions to the success of the CMS effort. In addition, we gratefully acknowledge the computing centers and personnel of the Worldwide LHC Computing Grid and other centers for delivering so effectively the computing infrastructure essential to our analyses. Finally, we acknowledge the enduring support for the construction and operation of the LHC, the CMS detector, and the supporting computing infrastructure provided by the following funding agencies: SC (Armenia), BMBWF and FWF (Austria); FNRS and FWO (Belgium); CNPq, CAPES, FAPERJ, FAPERGS, and FAPESP (Brazil); MES and BNSF (Bulgaria); CERN; CAS, MoST, and NSFC (China); MINCIENCIAS (Colombia); MSES and CSF (Croatia); RIF (Cyprus); SENESCYT (Ecuador); MoER, ERC PUT and ERDF (Estonia); Academy of Finland, MEC, and HIP (Finland); CEA and CNRS/IN2P3 (France); SRNSF (Georgia); BMBF, DFG, and HGF (Germany); GSRI (Greece); NKFIH (Hungary); DAE and DST (India); IPM (Iran); SFI (Ireland); INFN (Italy); MSIP and NRF (Republic of Korea); MES (Latvia); LAS (Lithuania); MOE and UM (Malaysia); BUAP, CINVESTAV, CONACYT, LNS, SEP, and UASLP-FAI (Mexico); MOS (Montenegro); MBIE (New Zealand); PAEC (Pakistan); MES and NSC (Poland); FCT (Portugal); MESTD (Serbia); MCIN/AEI and PCTI (Spain); MOSTR (Sri Lanka); Swiss Funding Agencies (Switzerland); MST (Taipei); MHESI and NSTDA (Thailand); TUBITAK and TENMAK (Turkey); NASU (Ukraine); STFC (United Kingdom); DOE and NSF (U.S.A.).

Individuals have received support from the Marie-Curie program and the European Research Council and Horizon 2020 Grant, contract Nos. 675440, 724704, 752730, 758316, 765710, 824093, and COST Action CA16108 (European Union); the Leventis Foundation; the Alfred P. Sloan Foundation; the Alexander von Humboldt Foundation; the Science Committee, project no. 22rl-037 (Armenia); the Belgian Federal Science Policy Office; the Fonds pour la Formation à la Recherche dans l’Industrie et dans l’Agriculture (FRIA-Belgium); the Agentschap voor Innovatie door Wetenschap en Technologie (IWT-Belgium); the F.R.S.-FNRS and FWO (Belgium) under the “Excellence of Science — EOS” — be.h project n. 30820817; the Beijing Municipal Science & Technology Commission, No. Z191100007219010 and Fundamental Research Funds for the Central Universities (China); the Ministry of Education, Youth and Sports (MEYS) of the Czech Republic; the Shota Rustaveli National Science Foundation, grant FR-22-985 (Georgia); the Deutsche Forschungsgemeinschaft (DFG), under Germany’s Excellence Strategy — EXC 2121 “Quantum Universe” — 390833306, and under project number 400140256 — GRK2497; the Hellenic Foundation for Research and Innovation (HFRI), Project Number 2288 (Greece); the Hungarian Academy of Sciences, the New National Excellence Program — ÚNKP, the NKFIH research grants K 124845, K 124850, K 128713, K 128786, K 129058, K 131991, K 133046, K 138136, K 143460, K 143477, 2020-2.2.1-ED-2021-00181, and TKP2021-NKTA-64 (Hungary); the Council of Science and Industrial Research, India; the Latvian Council of Science; the Ministry of Education and Science, project no. 2022/WK/14, and the National Science Center, contracts Opus 2021/41/B/ST2/01369 and 2021/43/B/ST2/01552 (Poland); the Fundação para a Ciência e a Tecnologia, grant CEECIND/01334/2018 (Portugal); the National Priorities Research Program by Qatar National Research Fund; MCIN/AEI/10.13039/501100011033, ERDF “a way of making Europe”, and the Programa Estatal de Fomento de la Investigación Científica y Técnica de Excelencia María de Maeztu, grant MDM-2017-0765 and Programa Severo Ochoa del Principado de Asturias (Spain); the Chulalongkorn Academic into Its 2nd Century Project Advancement Project, and the National Science, Research and Innovation Fund via the Program Management Unit for Human Resources & Institutional Development, Research and Innovation, grant B05F650021 (Thailand); the Kavli Foundation; the Nvidia Corporation; the SuperMicro Corporation; the Welch Foundation, contract C-1845; and the Weston Havens Foundation (U.S.A.).

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](#)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] E598 collaboration, *Experimental observation of a heavy particle J*, *Phys. Rev. Lett.* **33** (1974) 1404 [[INSPIRE](#)].
- [2] SLAC-SP-017 collaboration, *Discovery of a narrow resonance in e^+e^- annihilation*, *Phys. Rev. Lett.* **33** (1974) 1406 [[INSPIRE](#)].
- [3] E288 collaboration, *Observation of a dimuon resonance at 9.5 GeV in 400 GeV proton-nucleus collisions*, *Phys. Rev. Lett.* **39** (1977) 252 [[INSPIRE](#)].

- [4] UA1 collaboration, *Experimental observation of lepton pairs of invariant mass around 95 GeV/c² at the CERN SPS collider*, *Phys. Lett. B* **126** (1983) 398 [[INSPIRE](#)].
- [5] ATLAS collaboration, *Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC*, *Phys. Lett. B* **716** (2012) 1 [[arXiv:1207.7214](#)] [[INSPIRE](#)].
- [6] CMS collaboration, *Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC*, *Phys. Lett. B* **716** (2012) 30 [[arXiv:1207.7235](#)] [[INSPIRE](#)].
- [7] CMS collaboration, *Observation of a new boson with mass near 125 GeV in pp collisions at $\sqrt{s} = 7$ and 8 TeV*, *JHEP* **06** (2013) 081 [[arXiv:1303.4571](#)] [[INSPIRE](#)].
- [8] ATLAS collaboration, *Search for high-mass dilepton resonances using 139 fb⁻¹ of pp collision data collected at $\sqrt{s} = 13$ TeV with the ATLAS detector*, *Phys. Lett. B* **796** (2019) 68 [[arXiv:1903.06248](#)] [[INSPIRE](#)].
- [9] CMS collaboration, *Search for resonant and nonresonant new phenomena in high-mass dilepton final states at $\sqrt{s} = 13$ TeV*, *JHEP* **07** (2021) 208 [[arXiv:2103.02708](#)] [[INSPIRE](#)].
- [10] N. Arkani-Hamed, D.P. Finkbeiner, T.R. Slatyer and N. Weiner, *A theory of dark matter*, *Phys. Rev. D* **79** (2009) 015014 [[arXiv:0810.0713](#)] [[INSPIRE](#)].
- [11] BABAR collaboration, *Search for a dark photon in e⁺e⁻ collisions at BaBar*, *Phys. Rev. Lett.* **113** (2014) 201801 [[arXiv:1406.2980](#)] [[INSPIRE](#)].
- [12] LHCb collaboration, *Search for dark photons produced in 13 TeV pp collisions*, *Phys. Rev. Lett.* **120** (2018) 061801 [[arXiv:1710.02867](#)] [[INSPIRE](#)].
- [13] LHCb collaboration, *Search for A' $\rightarrow \mu^+ \mu^-$ decays*, *Phys. Rev. Lett.* **124** (2020) 041801 [[arXiv:1910.06926](#)] [[INSPIRE](#)].
- [14] LHCb collaboration, *Searches for low-mass dimuon resonances*, *JHEP* **10** (2020) 156 [[arXiv:2007.03923](#)] [[INSPIRE](#)].
- [15] CMS collaboration, *Search for a narrow resonance lighter than 200 GeV decaying to a pair of muons in proton-proton collisions at $\sqrt{s} = 13$ TeV*, *Phys. Rev. Lett.* **124** (2020) 131802 [[arXiv:1912.04776](#)] [[INSPIRE](#)].
- [16] CMS collaboration, *Search for narrow resonances in dijet final states at $\sqrt{s} = 8$ TeV with the novel CMS technique of data scouting*, *Phys. Rev. Lett.* **117** (2016) 031802 [[arXiv:1604.08907](#)] [[INSPIRE](#)].
- [17] P. Galison and A. Manohar, *Two Z's or not two Z's?*, *Phys. Lett. B* **136** (1984) 279 [[INSPIRE](#)].
- [18] B. Holdom, *Two U(1)'s and ϵ charge shifts*, *Phys. Lett. B* **166** (1986) 196 [[INSPIRE](#)].
- [19] D. Curtin, R. Essig, S. Gori and J. Shelton, *Illuminating dark photons with high-energy colliders*, *JHEP* **02** (2015) 157 [[arXiv:1412.0018](#)] [[INSPIRE](#)].
- [20] U. Haisch, J.F. Kamenik, A. Malinauskas and M. Spira, *Collider constraints on light pseudoscalars*, *JHEP* **03** (2018) 178 [[arXiv:1802.02156](#)] [[INSPIRE](#)].
- [21] S. Argyropoulos and U. Haisch, *Benchmarking LHC searches for light 2HDM+a pseudoscalars*, *SciPost Phys.* **13** (2022) 007 [[arXiv:2202.12631](#)] [[INSPIRE](#)].
- [22] *HEPData record for this analysis*, CMS-EXO-21-005, (2023) [[DOI:10.17182/HEPDATA.140424](#)].

- [23] CMS collaboration, *The CMS experiment at the CERN LHC*, 2008 *JINST* **3** S08004 [[INSPIRE](#)].
- [24] CMS collaboration, *The CMS trigger system*, 2017 *JINST* **12** P01020 [[arXiv:1609.02366](#)] [[INSPIRE](#)].
- [25] CMS collaboration, *Search for dijet resonances in proton-proton collisions at $\sqrt{s} = 13$ TeV and constraints on dark matter and other models*, *Phys. Lett. B* **769** (2017) 520 [[arXiv:1611.03568](#)] [[INSPIRE](#)].
- [26] CMS collaboration, *Data scouting: a new trigger paradigm*, in the proceedings of the 5th Large Hadron Collider physics conference, (2017) [[arXiv:1708.06925](#)] [[INSPIRE](#)].
- [27] CMS TRACKER GROUP collaboration, *The CMS phase-1 pixel detector upgrade*, 2021 *JINST* **16** P02027 [[arXiv:2012.14304](#)] [[INSPIRE](#)].
- [28] T. Sjöstrand et al., *An introduction to PYTHIA 8.2*, *Comput. Phys. Commun.* **191** (2015) 159 [[arXiv:1410.3012](#)] [[INSPIRE](#)].
- [29] S. Gopalakrishna, S. Jung and J.D. Wells, *Higgs boson decays to four fermions through an Abelian hidden sector*, *Phys. Rev. D* **78** (2008) 055002 [[arXiv:0801.3456](#)] [[INSPIRE](#)].
- [30] D. Curtin et al., *Exotic decays of the 125 GeV Higgs boson*, *Phys. Rev. D* **90** (2014) 075004 [[arXiv:1312.4992](#)] [[INSPIRE](#)].
- [31] J. Alwall et al., *The automated computation of tree-level and next-to-leading order differential cross sections, and their matching to parton shower simulations*, *JHEP* **07** (2014) 079 [[arXiv:1405.0301](#)] [[INSPIRE](#)].
- [32] NNPDF collaboration, *Parton distributions for the LHC run II*, *JHEP* **04** (2015) 040 [[arXiv:1410.8849](#)] [[INSPIRE](#)].
- [33] CMS collaboration, *Extraction and validation of a new set of CMS PYTHIA8 tunes from underlying-event measurements*, *Eur. Phys. J. C* **80** (2020) 4 [[arXiv:1903.12179](#)] [[INSPIRE](#)].
- [34] GEANT4 collaboration, *GEANT4 — a simulation toolkit*, *Nucl. Instrum. Meth. A* **506** (2003) 250 [[INSPIRE](#)].
- [35] B.P. Roe et al., *Boosted decision trees, an alternative to artificial neural networks*, *Nucl. Instrum. Meth. A* **543** (2005) 577 [[physics/0408124](#)] [[INSPIRE](#)].
- [36] CMS collaboration, *Performance of CMS muon reconstruction in pp collision events at $\sqrt{s} = 7$ TeV*, 2012 *JINST* **7** P10002 [[arXiv:1206.4071](#)] [[INSPIRE](#)].
- [37] J.E. Gaiser, *Charmonium spectroscopy from radiative decays of the J/ψ and ψ'* , Ph.D. thesis, SLAC report SLAC-R-255, Stanford University, Stanford, CA, U.S.A. (1982) [[INSPIRE](#)].
- [38] M. Oreglia, *A study of the reactions $\psi' \rightarrow \gamma\gamma\psi$* , Ph.D. thesis, SLAC report SLAC-R-236, Stanford University, Stanford, CA, U.S.A. (1980) [[INSPIRE](#)].
- [39] S. Bernstein, *Démonstration du théorème de Weierstrass fondée sur le calcul des probabilités* (in French), *Comm. Soc. Math. Kharkov* **13** (1912) 1.
- [40] R.A. Fisher, *On the interpretation of χ^2 from contingency tables, and the calculation of P* , *J. Roy. Statist. Soc. A* **85** (1922) 87.
- [41] P.D. Dauncey, M. Kenzie, N. Wardle and G.J. Davies, *Handling uncertainties in background shapes: the discrete profiling method*, 2015 *JINST* **10** P04015 [[arXiv:1408.6865](#)] [[INSPIRE](#)].

- [42] CMS collaboration, *CMS luminosity measurements for the 2016 data taking period*, CMS-PAS-LUM-17-001, CERN, Geneva, Switzerland (2017).
- [43] CMS collaboration, *CMS luminosity measurement for the 2018 data-taking period at $\sqrt{s} = 13 \text{ TeV}$* , CMS-PAS-LUM-18-002, CERN, Geneva, Switzerland (2019).
- [44] CMS collaboration, *Precision luminosity measurement in proton-proton collisions at $\sqrt{s} = 13 \text{ TeV}$ in 2015 and 2016 at CMS*, *Eur. Phys. J. C* **81** (2021) 800 [[arXiv:2104.01927](#)] [[INSPIRE](#)].
- [45] T. Junk, *Confidence level computation for combining searches with small statistics*, *Nucl. Instrum. Meth. A* **434** (1999) 435 [[hep-ex/9902006](#)] [[INSPIRE](#)].
- [46] A.L. Read, *Presentation of search results: the CL_s technique*, *J. Phys. G* **28** (2002) 2693 [[INSPIRE](#)].
- [47] G. Cowan, K. Cranmer, E. Gross and O. Vitells, *Asymptotic formulae for likelihood-based tests of new physics*, *Eur. Phys. J. C* **71** (2011) 1554 [Erratum *ibid.* **73** (2013) 2501] [[arXiv:1007.1727](#)] [[INSPIRE](#)].
- [48] K.R. Dienes, C.F. Kolda and J. March-Russell, *Kinetic mixing and the supersymmetric gauge hierarchy*, *Nucl. Phys. B* **492** (1997) 104 [[hep-ph/9610479](#)] [[INSPIRE](#)].
- [49] S. Camarda et al., *DYTurbo: fast predictions for Drell-Yan processes*, *Eur. Phys. J. C* **80** (2020) 251 [Erratum *ibid.* **80** (2020) 440] [[arXiv:1910.07049](#)] [[INSPIRE](#)].
- [50] P. Ilten, Y. Soreq, M. Williams and W. Xue, *Serendipity in dark photon searches*, *JHEP* **06** (2018) 004 [[arXiv:1801.04847](#)] [[INSPIRE](#)].

The CMS collaboration

Yerevan Physics Institute, Yerevan, Armenia

A. Hayrapetyan, A. Tumasyan¹

Institut für Hochenergiephysik, Vienna, Austria

W. Adam¹, J.W. Andrejkovic, T. Bergauer¹, S. Chatterjee¹, K. Damanakis¹, M. Dragicevic¹, A. Escalante Del Valle¹, P.S. Hussain¹, M. Jeitler^{1,2}, N. Krammer¹, D. Liko¹, I. Mikulec¹, J. Schieck^{1,2}, R. Schöfbeck¹, D. Schwarz¹, M. Sonawane¹, S. Templ¹, W. Waltenberger¹, C.-E. Wulz^{1,2}

Universiteit Antwerpen, Antwerpen, Belgium

M.R. Darwish³, T. Janssen¹, P. Van Mechelen¹

Vrije Universiteit Brussel, Brussel, Belgium

E.S. Bols¹, J. D'Hondt¹, S. Dansana¹, A. De Moor¹, M. Delcourt¹, H. El Faham¹, S. Lowette¹, I. Makarenko¹, A. Morton¹, D. Müller¹, A.R. Sahasransu¹, S. Tavernier¹, M. Tytgat^{1,4}, S. Van Putte¹, D. Vannerom¹

Université Libre de Bruxelles, Bruxelles, Belgium

B. Clerbaux¹, G. De Lentdecker¹, L. Favart¹, D. Hohov¹, J. Jaramillo¹, A. Khalilzadeh, K. Lee¹, M. Mahdavikhorrami¹, A. Malara¹, S. Paredes¹, L. Pétré¹, N. Postiau, L. Thomas¹, M. Vanden Bemden¹, C. Vander Velde¹, P. Vanlaer¹

Ghent University, Ghent, Belgium

M. De Coen¹, D. Dobur¹, J. Knolle¹, L. Lambrecht¹, G. Mestdach, C. Rendón, A. Samalan, K. Skovpen¹, N. Van Den Bossche¹, L. Wezenbeek¹

Université Catholique de Louvain, Louvain-la-Neuve, Belgium

A. Benecke¹, G. Bruno¹, C. Caputo¹, C. Delaere¹, I.S. Donertas¹, A. Giannanco¹, K. Jaffel¹, Sa. Jain¹, V. Lemaitre, J. Lidrych¹, P. Mastrapasqua¹, K. Mondal¹, T.T. Tran¹, S. Wertz¹

Centro Brasileiro de Pesquisas Fisicas, Rio de Janeiro, Brazil

G.A. Alves¹, E. Coelho¹, C. Hensel¹, T. Menezes De Oliveira, A. Moraes¹, P. Rebello Teles¹, M. Soeiro

Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil

W.L. Aldá Júnior¹, M. Alves Gallo Pereira¹, M. Barroso Ferreira Filho¹, H. Brandao Malbouisson¹, W. Carvalho¹, J. Chinellato⁵, E.M. Da Costa¹, G.G. Da Silveira^{1,6}, D. De Jesus Damiao¹, S. Fonseca De Souza¹, J. Martins^{1,7}, C. Mora Herrera¹, K. Mota Amarilo¹, L. Mundim¹, H. Nogima¹, A. Santoro¹, S.M. Silva Do Amaral¹, A. Sznajder¹, M. Thiel¹, A. Vilela Pereira¹

**Universidade Estadual Paulista, Universidade Federal do ABC, São Paulo,
Brazil**

C.A. Bernardes⁶, L. Calligaris¹⁰, T.R. Fernandez Perez Tomei¹⁰, E.M. Gregores¹⁰,
P.G. Mercadante¹⁰, S.F. Novaes¹⁰, B. Orzari¹⁰, Sandra S. Padula¹⁰

**Institute for Nuclear Research and Nuclear Energy, Bulgarian Academy of
Sciences, Sofia, Bulgaria**

A. Aleksandrov¹⁰, G. Antchev¹⁰, R. Hadjiiska¹⁰, P. Iaydjiev¹⁰, M. Misheva¹⁰, M. Shopova¹⁰,
G. Sultanov¹⁰

University of Sofia, Sofia, Bulgaria

A. Dimitrov¹⁰, T. Ivanov¹⁰, L. Litov¹⁰, B. Pavlov¹⁰, P. Petkov¹⁰, A. Petrov¹⁰, E. Shumka¹⁰

**Instituto De Alta Investigación, Universidad de Tarapacá, Casilla 7 D, Arica,
Chile**

S. Keshri¹⁰, S. Thakur¹⁰

Beihang University, Beijing, China

T. Cheng¹⁰, Q. Guo, T. Javaid¹⁰, M. Mittal¹⁰, L. Yuan¹⁰

Department of Physics, Tsinghua University, Beijing, China

G. Bauer⁸, Z. Hu¹⁰, K. Yi^{10,8,9}

Institute of High Energy Physics, Beijing, China

G.M. Chen¹⁰, H.S. Chen¹⁰, M. Chen¹⁰, F. Iemmi¹⁰, C.H. Jiang, A. Kapoor¹⁰, H. Liao¹⁰,
Z.-A. Liu¹¹, F. Monti¹⁰, R. Sharma¹⁰, J.N. Song¹¹, J. Tao¹⁰, J. Wang¹⁰, H. Zhang¹⁰

**State Key Laboratory of Nuclear Physics and Technology, Peking University,
Beijing, China**

A. Agapitos¹⁰, Y. Ban¹⁰, A. Levin¹⁰, C. Li¹⁰, Q. Li¹⁰, X. Lyu, Y. Mao, S.J. Qian¹⁰, X. Sun¹⁰,
D. Wang¹⁰, H. Yang, C. Zhou¹⁰

Sun Yat-Sen University, Guangzhou, China

Z. You¹⁰

University of Science and Technology of China, Hefei, China

N. Lu¹⁰

**Institute of Modern Physics and Key Laboratory of Nuclear Physics and
Ion-beam Application (MOE) — Fudan University, Shanghai, China**

X. Gao¹², D. Leggat, H. Okawa¹⁰, Y. Zhang¹⁰

Zhejiang University, Hangzhou, Zhejiang, China

Z. Lin¹⁰, C. Lu¹⁰, M. Xiao¹⁰

Universidad de Los Andes, Bogota, Colombia

C. Avila¹⁰, D.A. Barbosa Trujillo, A. Cabrera¹⁰, C. Florez¹⁰, J. Fraga¹⁰, J.A. Reyes Vega

Universidad de Antioquia, Medellin, ColombiaJ. Mejia Guisao , F. Ramirez , M. Rodriguez , J.D. Ruiz Alvarez **University of Split, Faculty of Electrical Engineering, Mechanical Engineering and Naval Architecture, Split, Croatia**D. Giljanovic , N. Godinovic , D. Lelas , A. Sculac **University of Split, Faculty of Science, Split, Croatia**M. Kovac , T. Sculac **Institute Rudjer Boskovic, Zagreb, Croatia**P. Bargassa , V. Brigljevic , B.K. Chitroda , D. Ferencek , S. Mishra , A. Starodumov ¹³, T. Susa **University of Cyprus, Nicosia, Cyprus**A. Attikis , K. Christoforou , S. Konstantinou , J. Mousa , C. Nicolaou, F. Ptochos , P.A. Razis , H. Rykaczewski, H. Saka , A. Stepennov **Charles University, Prague, Czech Republic**M. Finger , M. Finger Jr. , A. Kveton **Escuela Politecnica Nacional, Quito, Ecuador**E. Ayala **Universidad San Francisco de Quito, Quito, Ecuador**E. Carrera Jarrin **Academy of Scientific Research and Technology of the Arab Republic of Egypt, Egyptian Network of High Energy Physics, Cairo, Egypt**Y. Assran^{14,15}, S. Elgammal¹⁵**Center for High Energy Physics (CHEP-FU), Fayoum University, El-Fayoum, Egypt**M. Abdullah Al-Mashad , M.A. Mahmoud **National Institute of Chemical Physics and Biophysics, Tallinn, Estonia**R.K. Dewanjee ¹⁶, K. Ehataht , M. Kadastik, T. Lange , S. Nandan , C. Nielsen , J. Pata , M. Raidal , L. Tani , C. Veelken **Department of Physics, University of Helsinki, Helsinki, Finland**H. Kirschenmann , K. Osterberg , M. Voutilainen **Helsinki Institute of Physics, Helsinki, Finland**S. Bharthuar , E. Brücken , F. Garcia , J. Havukainen , K.T.S. Kallonen , M.S. Kim , R. Kinnunen, T. Lampén , K. Lassila-Perini , S. Lehti , T. Lindén , M. Lotti,

L. Martikainen , M. Myllymäki , M.m. Rantanen , H. Siikonen , E. Tuominen , J. Tuominiemi 

Lappeenranta-Lahti University of Technology, Lappeenranta, Finland

P. Luukka , H. Petrow , T. Tuuva[†]

IRFU, CEA, Université Paris-Saclay, Gif-sur-Yvette, France

M. Besancon , F. Couderc , M. Dejardin , D. Denegri, J.L. Faure, F. Ferri , S. Ganjour , P. Gras , G. Hamel de Monchenault , V. Lohezic , J. Malcles , J. Rander, A. Rosowsky , M.Ö. Sahin , A. Savoy-Navarro ¹⁷, P. Simkina , M. Titov

Laboratoire Leprince-Ringuet, CNRS/IN2P3, Ecole Polytechnique, Institut Polytechnique de Paris, Palaiseau, France

C. Baldenegro Barrera , F. Beaudette , A. Buchot Perraguin , P. Busson , A. Cappati , C. Charlot , F. Damas , O. Davignon , G. Falmagne , B.A. Fontana Santos Alves , S. Ghosh , A. Gilbert , R. Granier de Cassagnac , A. Hakimi , B. Harikrishnan , L. Kalipoliti , G. Liu , J. Motta , M. Nguyen , C. Ochando , L. Portales , R. Salerno , U. Sarkar , J.B. Sauvan , Y. Sirois , A. Tarabini , E. Vernazza , A. Zabi , A. Zghiche

Université de Strasbourg, CNRS, IPHC UMR 7178, Strasbourg, France

J.-L. Agram ¹⁸, J. Andrea , D. Apparu , D. Bloch , J.-M. Brom , E.C. Chabert , C. Collard , S. Falke , U. Goerlach , C. Grimault, R. Haeberle , A.-C. Le Bihan , M.A. Sessini , P. Van Hove

Institut de Physique des 2 Infinis de Lyon (IP2I), Villeurbanne, France

S. Beauceron , B. Blancon , G. Boudoul , N. Chanon , J. Choi , D. Contardo , P. Depasse , C. Dozen ¹⁹, H. El Mamouni, J. Fay , S. Gascon , M. Gouzevitch , C. Greenberg, G. Grenier , B. Ille , I.B. Laktineh, M. Lethuillier , L. Mirabito, S. Perries, M. Vander Donckt , P. Verdier , J. Xiao

Georgian Technical University, Tbilisi, Georgia

A. Khvedelidze ¹³, I. Lomidze , Z. Tsamalaidze ¹³

RWTH Aachen University, I. Physikalisches Institut, Aachen, Germany

V. Botta , L. Feld , K. Klein , M. Lipinski , D. Meuser , A. Pauls , N. Röwert , M. Teroerde 

RWTH Aachen University, III. Physikalisches Institut A, Aachen, Germany

S. Diekmann , A. Dodonova , N. Eich , D. Eliseev , F. Engelke , M. Erdmann , P. Fackeldey , B. Fischer , T. Hebbeker , K. Hoepfner , F. Ivone , A. Jung , M.y. Lee , L. Mastrolorenzo, M. Merschmeyer , A. Meyer , S. Mukherjee , D. Noll , A. Novak , F. Nowotny, A. Pozdnyakov , Y. Rath, W. Redjeb , F. Rehm, H. Reithler , V. Sarkisov , A. Schmidt , S.C. Schuler, A. Sharma , A. Stein , F. Torres Da Silva De Araujo ²⁰, L. Vigilante, S. Wiedenbeck , S. Zaleski

RWTH Aachen University, III. Physikalisches Institut B, Aachen, Germany

C. Dziwok , G. Flügge , W. Haj Ahmad ²¹, T. Kress , A. Nowack , O. Pooth ,
A. Stahl , T. Ziemons , A. Zottz 

Deutsches Elektronen-Synchrotron, Hamburg, Germany

H. Aarup Petersen , M. Aldaya Martin , J. Alimena , S. Amoroso, Y. An , S. Baxter ,
M. Bayatmakou , H. Becerril Gonzalez , O. Behnke , A. Belvedere , S. Bhattacharya ,
F. Blekman ²², K. Borras , D. Brunner , A. Campbell , A. Cardini , C. Cheng,
F. Colombina , S. Consuegra Rodríguez , G. Correia Silva , M. De Silva , G. Eckerlin,
D. Eckstein , L.I. Estevez Banos , O. Filatov , E. Gallo ²², A. Geiser , A. Giraldi ,
G. Greau, V. Guglielmi , M. Guthoff , A. Hinzmann , A. Jafari ²⁴, L. Jeppe ,
N.Z. Jomhari , B. Kaech , M. Kasemann , H. Kaveh , C. Kleinwort , R. Kogler ,
M. Komm , D. Krücker , W. Lange, D. Leyva Pernia , K. Lipka ²⁵, W. Lohmann ²⁶,
R. Mankel , I.-A. Melzer-Pellmann , M. Mendizabal Morentin , J. Metwally, A.B. Meyer ,
G. Milella , M. Mormile , A. Mussgiller , A. Nürnberg , Y. Otarid, D. Pérez Adán ,
E. Ranken , A. Raspereza , B. Ribeiro Lopes , J. Rübenach, A. Saggio , M. Scham ^{27,23},
V. Scheurer, S. Schnake ²³, P. Schütze , C. Schwanenberger ²², M. Shchedrolosiev ,
R.E. Sosa Ricardo , L.P. Sreelatha Pramod , D. Stafford, F. Vazzoler , A. Ventura Barroso ,
R. Walsh , Q. Wang , Y. Wen , K. Wichmann, L. Wiens ²³, C. Wissing , S. Wuchterl ,
Y. Yang , A. Zimmermann Castro Santos 

University of Hamburg, Hamburg, Germany

A. Albrecht , S. Albrecht , M. Antonello , S. Bein , L. Benato , M. Bonanomi ,
P. Connor , M. Eich, K. El Morabit , Y. Fischer , A. Fröhlich, C. Garbers , E. Garutti ,
A. Grohsjean , M. Hajheidari, J. Haller , H.R. Jabusch , G. Kasieczka , P. Keicher,
R. Klanner , W. Korcari , T. Kramer , V. Kutzner , F. Labe , J. Lange , A. Lobanov ,
C. Matthies , A. Mehta , L. Moureaux , M. Mrowietz, A. Nigamova , Y. Nissan,
A. Paasch , K.J. Pena Rodriguez , T. Quadfasel , B. Raciti , M. Rieger , D. Savoiu ,
J. Schindler , P. Schleper , M. Schröder , J. Schwandt , M. Sommerhalder , H. Stadie ,
G. Steinbrück , A. Tews, M. Wolf 

Karlsruher Institut fuer Technologie, Karlsruhe, Germany

S. Brommer , M. Burkart, E. Butz , T. Chwalek , A. Dierlamm , A. Droll,
N. Faltermann , M. Giffels , A. Gottmann , F. Hartmann ²⁸, M. Horzela , U. Husemann ,
M. Klute , R. Koppenhöfer , M. Link, A. Lintuluoto , S. Maier , S. Mitra , Th. Müller ,
M. Neukum, M. Oh , G. Quast , K. Rabbertz , I. Shvetsov , H.J. Simonis , N. Trevisani ,
R. Ulrich , J. van der Linden , R.F. Von Cube , M. Wassmer , S. Wieland , F. Wittig,
R. Wolf , S. Wunsch, X. Zuo 

Institute of Nuclear and Particle Physics (INPP), NCSR Demokritos, Aghia Paraskevi, Greece

G. Anagnostou, P. Assiouras , G. Daskalakis , A. Kyriakis, A. Papadopoulos²⁸, A. Stakia 

National and Kapodistrian University of Athens, Athens, Greece

D. Karasavvas, P. Kontaxakis , G. Melachroinos, A. Panagiotou, I. Papavergou ,
I. Paraskevas , N. Saoulidou , K. Theofilatos , E. Tziaferi , K. Vellidis , I. Zisopoulos

National Technical University of Athens, Athens, Greece

G. Bakas , T. Chatzistavrou, G. Karapostoli , K. Kousouris , I. Papakrivopoulos ,
E. Siamarkou, G. Tsipolitis, A. Zacharopoulou

University of Ioánnina, Ioánnina, Greece

K. Adamidis, I. Bestintzanos, I. Evangelou , C. Foudas, P. Gianneios , C. Kamtsikis,
P. Katsoulis, P. Kokkas , P.G. Kosmoglou Kioseoglou , N. Manthos , I. Papadopoulos ,
J. Strologas

MTA-ELTE Lendület CMS Particle and Nuclear Physics Group, Eötvös Loránd University, Budapest, Hungary

M. Csanad , K. Farkas , M.M.A. Gadallah ²⁹, . Kadlecseik , P. Major , K. Mandal ,
G. Pasztor , A.J. Radl ³⁰, O. Suranyi , G.I. Veres

Wigner Research Centre for Physics, Budapest, Hungary

M. Bartok ³¹, C. Hajdu , D. Horvath ^{32,33}, F. Sikler , V. Veszpremi

Institute of Nuclear Research ATOMKI, Debrecen, Hungary

G. Bencze, S. Czellar, J. Karancsi ³¹, J. Molnar, Z. Szillasi

Institute of Physics, University of Debrecen, Debrecen, Hungary

P. Raics, B. Ujvari ³⁴, G. Zilizi

Karoly Robert Campus, MATE Institute of Technology, Gyongyos, Hungary

T. Csorgo ³⁰, F. Nemes ³⁰, T. Novak

Panjab University, Chandigarh, India

J. Babbar , S. Bansal , S.B. Beri, V. Bhatnagar , G. Chaudhary , S. Chauhan ,
N. Dhingra ³⁵, R. Gupta, A. Kaur , A. Kaur , H. Kaur , M. Kaur , S. Kumar ,
P. Kumari , M. Meena , K. Sandeep , T. Sheokand, J.B. Singh ³⁶, A. Singla

University of Delhi, Delhi, India

A. Ahmed , A. Bhardwaj , A. Chhetri , B.C. Choudhary , A. Kumar , M. Naimuddin ,
K. Ranjan , S. Saumya

Saha Institute of Nuclear Physics, HBNI, Kolkata, India

S. Baradia , S. Barman ³⁷, S. Bhattacharya , D. Bhowmik, S. Dutta , S. Dutta,
B. Gomber ³⁸, P. Palit , G. Saha , B. Sahu ³⁸, S. Sarkar

Indian Institute of Technology Madras, Madras, India

P.K. Behera , S.C. Behera , S. Chatterjee , P. Jana , P. Kalbhor , J.R. Komaragiri ³⁹,
D. Kumar ³⁹, M. Mohammad Mobassir Ameen , L. Panwar ³⁹, R. Pradhan ,
P.R. Pujahari , N.R. Saha , A. Sharma , A.K. Sikdar , S. Verma

Tata Institute of Fundamental Research-A, Mumbai, IndiaT. Aziz, I. Das^{ID}, S. Dugad, M. Kumar^{ID}, G.B. Mohanty^{ID}, P. Suryadevara**Tata Institute of Fundamental Research-B, Mumbai, India**A. Bala^{ID}, S. Banerjee^{ID}, R.M. Chatterjee, M. Guchait^{ID}, S. Karmakar^{ID}, S. Kumar^{ID},
G. Majumder^{ID}, K. Mazumdar^{ID}, S. Mukherjee^{ID}, A. Thachayath^{ID}**National Institute of Science Education and Research, An OCC of Homi Bhabha National Institute, Bhubaneswar, Odisha, India**S. Bahinipati^{ID}⁴⁰, A.K. Das, C. Kar^{ID}, D. Maity^{ID}⁴¹, P. Mal^{ID}, T. Mishra^{ID},
V.K. Muraleedharan Nair Bindhu^{ID}⁴¹, K. Naskar^{ID}⁴¹, A. Nayak^{ID}⁴¹, P. Sadangi, P. Saha^{ID},
S.K. Swain^{ID}, S. Varghese^{ID}⁴¹, D. Vats^{ID}⁴¹**Indian Institute of Science Education and Research (IISER), Pune, India**A. Alpana^{ID}, S. Dube^{ID}, B. Kansal^{ID}, A. Laha^{ID}, A. Rastogi^{ID}, S. Sharma^{ID}**Isfahan University of Technology, Isfahan, Iran**H. Bakhshiansohi^{ID}^{42,43}, E. Khazaie^{ID}⁴³, M. Zeinali^{ID}⁴⁴**Institute for Research in Fundamental Sciences (IPM), Tehran, Iran**S. Chenarani^{ID}⁴⁵, S.M. Etesami^{ID}, M. Khakzad^{ID}, M. Mohammadi Najafabadi^{ID}**University College Dublin, Dublin, Ireland**M. Grunewald^{ID}**INFN Sezione di Bari^a, Università di Bari^b, Politecnico di Bari^c, Bari, Italy**M. Abbrescia^{ID}^{a,b}, R. Aly^{ID}^{a,c,46}, A. Colaleo^{ID}^a, D. Creanza^{ID}^{a,c}, B. D' Anzi^{ID}^{a,b},
N. De Filippis^{ID}^{a,c}, M. De Palma^{ID}^{a,b}, A. Di Florio^{ID}^{a,c}, W. Elmetenawee^{ID}^{a,b}, L. Fiore^{ID}^a,
G. Iaselli^{ID}^{a,c}, G. Maggi^{ID}^{a,c}, M. Maggi^{ID}^a, I. Margjeka^{ID}^{a,b}, V. Mastrapasqua^{ID}^{a,b}, S. My^{ID}^{a,b},
S. Nuzzo^{ID}^{a,b}, A. Pellecchia^{ID}^{a,b}, A. Pompili^{ID}^{a,b}, G. Pugliese^{ID}^{a,c}, R. Radogna^{ID}^a,
G. Ramirez-Sanchez^{ID}^{a,c}, D. Ramos^{ID}^a, A. Ranieri^{ID}^a, L. Silvestris^{ID}^a, F.M. Simone^{ID}^{a,b},
Ü. Sözbilir^{ID}^a, A. Stamerra^{ID}^a, R. Venditti^{ID}^a, P. Verwilligen^{ID}^a, A. Zaza^{ID}^{a,b}**INFN Sezione di Bologna^a, Università di Bologna^b, Bologna, Italy**G. Abbiendi^{ID}^a, C. Battilana^{ID}^{a,b}, D. Bonacorsi^{ID}^{a,b}, L. Borgonovi^{ID}^a, R. Campanini^{ID}^{a,b},
P. Capiluppi^{ID}^{a,b}, A. Castro^{ID}^{a,b}, F.R. Cavallo^{ID}^a, M. Cuffiani^{ID}^{a,b}, G.M. Dallavalle^{ID}^a,
T. Diotalevi^{ID}^{a,b}, A. Fanfani^{ID}^{a,b}, D. Fasanella^{ID}^{a,b}, P. Giacomelli^{ID}^a, L. Giommi^{ID}^{a,b},
C. Grandi^{ID}^a, L. Guiducci^{ID}^{a,b}, S. Lo Meo^{ID}^{a,47}, L. Lunerti^{ID}^{a,b}, S. Marcellini^{ID}^a, G. Masetti^{ID}^a,
F.L. Navarrone^{ID}^{a,b}, A. Perrotta^{ID}^a, F. Primavera^{ID}^{a,b}, A.M. Rossi^{ID}^{a,b}, T. Rovelli^{ID}^{a,b},
G.P. Siroli^{ID}^{a,b}**INFN Sezione di Catania^a, Università di Catania^b, Catania, Italy**S. Costa^{ID}^{a,b,48}, A. Di Mattia^{ID}^a, R. Potenza^{a,b}, A. Tricomi^{ID}^{a,b,48}, C. Tuve^{ID}^{a,b}

INFN Sezione di Firenze^a, Università di Firenze^b, Firenze, Italy

G. Barbagli ^a, G. Bardelli ^{a,b}, B. Camaiani ^{a,b}, A. Cassese ^a, R. Ceccarelli ^a, V. Ciulli ^{a,b}, C. Civinini ^a, R. D'Alessandro ^{a,b}, E. Focardi ^{a,b}, G. Latino ^{a,b}, P. Lenzi ^{a,b}, M. Lizzo ^{a,b}, M. Meschini ^a, S. Paoletti ^a, A. Papanastassiou ^{a,b}, G. Sguazzoni ^a, L. Viliani ^a

INFN Laboratori Nazionali di Frascati, Frascati, Italy

L. Benussi , S. Bianco , S. Meola ⁴⁹, D. Piccolo 

INFN Sezione di Genova^a, Università di Genova^b, Genova, Italy

P. Chatagnon ^a, F. Ferro , E. Robutti ^a, S. Tosi ^{a,b}

INFN Sezione di Milano-Bicocca^a, Università di Milano-Bicocca^b, Milano, Italy

A. Benaglia ^a, G. Boldrini , F. Brivio ^a, F. Cetorelli ^a, F. De Guio ^{a,b}, M.E. Dinardo ^{a,b}, P. Dini ^a, S. Gennai ^a, A. Ghezzi ^{a,b}, P. Govoni ^{a,b}, L. Guzzi ^a, M.T. Lucchini ^{a,b}, M. Malberti ^a, S. Malvezzi ^a, A. Massironi ^a, D. Menasce ^a, L. Moroni ^a, M. Paganoni ^{a,b}, D. Pedrini ^a, B.S. Pinolini^a, S. Ragazzi ^{a,b}, N. Redaelli ^a, T. Tabarelli de Fatis ^{a,b}, D. Zuolo ^a

INFN Sezione di Napoli^a, Università di Napoli ‘Federico II’^b, Napoli, Italy; Università della Basilicata^c, Potenza, Italy; Università G. Marconi^d, Roma, Italy

S. Buontempo ^a, A. Cagnotta ^{a,b}, F. Carnevali^{a,b}, N. Cavallo ^{a,c}, A. De Iorio ^{a,b}, F. Fabozzi ^{a,c}, A.O.M. Iorio ^{a,b}, L. Lista ^{a,b,⁵⁰}, P. Paolucci ^{a,28}, B. Rossi ^a, C. Sciacca ^{a,b}

INFN Sezione di Padova^a, Università di Padova^b, Padova, Italy; Università di Trento^c, Trento, Italy

R. Ardino ^a, P. Azzi ^a, N. Bacchetta ^{a,51}, D. Bisello ^{a,b}, P. Bortignon ^a, A. Bragagnolo ^{a,b}, R. Carlin ^{a,b}, P. Checchia ^a, T. Dorigo ^a, S. Fantinel ^a, F. Gasparini ^{a,b}, U. Gasparini ^{a,b}, G. Grossi ^a, L. Layer^{a,52}, E. Lusiani ^a, M. Margoni ^{a,b}, A.T. Meneguzzo ^{a,b}, M. Migliorini ^{a,b}, J. Pazzini ^{a,b}, P. Ronchese ^{a,b}, R. Rossin ^{a,b}, F. Simonetto ^{a,b}, G. Strong ^a, M. Tosi ^{a,b}, A. Triossi ^{a,b}, S. Ventura ^a, H. Yarar^{a,b}, M. Zanetti ^{a,b}, A. Zucchetta ^{a,b}, G. Zumerle ^{a,b}

INFN Sezione di Pavia^a, Università di Pavia^b, Pavia, Italy

S. Abu Zeid ^{a,53}, C. Aimè ^{a,b}, A. Braghieri ^a, S. Calzaferri ^{a,b}, D. Fiorina ^{a,b}, P. Montagna ^{a,b}, V. Re ^a, C. Riccardi ^{a,b}, P. Salvini ^a, I. Vai ^{a,b}, P. Vitulo ^{a,b}

INFN Sezione di Perugia^a, Università di Perugia^b, Perugia, Italy

S. Ajmal ^{a,b}, P. Asenov ^{a,54}, G.M. Bilei ^a, D. Ciangottini ^{a,b}, L. Fanò ^{a,b}, M. Magherini ^{a,b}, G. Mantovani^{a,b}, V. Mariani ^{a,b}, M. Menichelli ^a, F. Moscatelli ^{a,54}, A. Piccinelli ^{a,b}, M. Presilla ^{a,b}, A. Rossi ^{a,b}, A. Santocchia ^{a,b}, D. Spiga ^a, T. Tedeschi ^{a,b}

INFN Sezione di Pisa^a, Università di Pisa^b, Scuola Normale Superiore di Pisa^c, Pisa, Italy; Università di Siena^d, Siena, Italy

P. Azzurri ^a, G. Bagliesi ^a, R. Bhattacharya ^a, L. Bianchini ^{a,b}, T. Boccali ^a, E. Bossini ^a, D. Bruschini ^{a,c}, R. Castaldi ^a, M.A. Ciocci ^{a,b}, M. Cipriani ^{a,b}, V. D'Amante ^{a,d}, R. Dell'Orso ^a, S. Donato ^a, A. Giassi ^a, F. Ligabue ^{a,c}, D. Matos Figueiredo ^a, A. Messineo ^{a,b}, M. Musich ^{a,b}, F. Palla ^a, S. Parolia ^a, A. Rizzi ^{a,b}, G. Rolandi ^{a,c}, S. Roy Chowdhury ^a, T. Sarkar ^a, A. Scribano ^a, P. Spagnolo ^a, R. Tenchini ^{a,b}, G. Tonelli ^{a,b}, N. Turini ^{a,d}, A. Venturi ^a, P.G. Verdini ^a

INFN Sezione di Roma^a, Sapienza Università di Roma^b, Roma, Italy

P. Barria ^a, M. Campana ^{a,b}, F. Cavallari ^a, L. Cunqueiro Mendez ^{a,b}, D. Del Re ^{a,b}, E. Di Marco ^a, M. Diemoz ^a, F. Errico ^{a,b}, E. Longo ^{a,b}, P. Meridiani ^a, J. Mijuskovic ^{a,b}, G. Organtini ^{a,b}, F. Pandolfi ^a, R. Paramatti ^{a,b}, C. Quaranta ^{a,b}, S. Rahatlou ^{a,b}, C. Rovelli ^a, F. Santanastasio ^{a,b}, L. Soffi ^a, R. Tramontano ^{a,b}

INFN Sezione di Torino^a, Università di Torino^b, Torino, Italy; Università del Piemonte Orientale^c, Novara, Italy

N. Amapane ^{a,b}, R. Arcidiacono ^{a,c}, S. Argiro ^{a,b}, M. Arneodo ^{a,c}, N. Bartosik ^a, R. Bellan ^{a,b}, A. Bellora ^{a,b}, C. Biino ^a, N. Cartiglia ^a, M. Costa ^{a,b}, R. Covarelli ^{a,b}, N. Demaria ^a, L. Finco ^a, M. Grippo ^{a,b}, B. Kiani ^{a,b}, F. Legger ^a, F. Luongo ^{a,b}, C. Mariotti ^a, S. Maselli ^a, A. Mecca ^{a,b}, E. Migliore ^{a,b}, M. Monteno ^a, R. Mulargia ^a, M.M. Obertino ^{a,b}, G. Ortona ^a, L. Pacher ^{a,b}, N. Pastrone ^a, M. Pelliccioni ^a, M. Ruspa ^{a,c}, F. Siviero ^{a,b}, V. Sola ^{a,b}, A. Solano ^{a,b}, D. Soldi ^{a,b}, A. Staiano ^a, C. Tarricone ^{a,b}, M. Tornago ^{a,b}, D. Trocino ^a, G. Umoret ^{a,b}, A. Vagnerini ^{a,b}, E. Vlasov ^{a,b}

INFN Sezione di Trieste^a, Università di Trieste^b, Trieste, Italy

S. Belforte ^a, V. Candelise ^{a,b}, M. Casarsa ^a, F. Cossutti ^a, K. De Leo ^{a,b}, G. Della Ricca ^{a,b}

Kyungpook National University, Daegu, Korea

S. Dogra , J. Hong , C. Huh , B. Kim , D.H. Kim , J. Kim, H. Lee, J. Lee , S.W. Lee , C.S. Moon , Y.D. Oh , S.I. Pak , M.S. Ryu , S. Sekmen , Y.C. Yang 

Chonnam National University, Institute for Universe and Elementary Particles, Kwangju, Korea

G. Bak , P. Gwak , H. Kim , D.H. Moon 

Hanyang University, Seoul, Korea

E. Asilar , D. Kim , T.J. Kim , J.A. Merlin, J. Park 

Korea University, Seoul, Korea

S. Choi , S. Han, B. Hong , K. Lee, K.S. Lee , J. Park, S.K. Park, J. Yoo 

Kyung Hee University, Department of Physics, Seoul, Korea

J. Goh 

Sejong University, Seoul, KoreaH. S. Kim[✉], Y. Kim, S. Lee**Seoul National University, Seoul, Korea**J. Almond, J.H. Bhyun, J. Choi[✉], S. Jeon[✉], W. Jun[✉], J. Kim[✉], J.S. Kim, S. Ko[✉],
H. Kwon[✉], H. Lee[✉], J. Lee[✉], S. Lee, B.H. Oh[✉], S.B. Oh[✉], H. Seo[✉], U.K. Yang, I. Yoon[✉]**University of Seoul, Seoul, Korea**W. Jang[✉], D.Y. Kang, Y. Kang[✉], S. Kim[✉], B. Ko, J.S.H. Lee[✉], Y. Lee[✉], I.C. Park[✉],
Y. Roh, I.J. Watson[✉], S. Yang[✉]**Yonsei University, Department of Physics, Seoul, Korea**S. Ha[✉], H.D. Yoo[✉]**Sungkyunkwan University, Suwon, Korea**M. Choi[✉], M.R. Kim[✉], H. Lee, Y. Lee[✉], I. Yu[✉]**College of Engineering and Technology, American University of the Middle
East (AUM), Dasman, Kuwait**T. Beyrouthy, Y. Maghrbi[✉]**Riga Technical University, Riga, Latvia**K. Dreimanis[✉], A. Gaile[✉], G. Pikurs, A. Potrebko[✉], M. Seidel[✉], V. Veckalns[✉]⁵⁵**University of Latvia (LU), Riga, Latvia**N.R. Strautnieks[✉]**Vilnius University, Vilnius, Lithuania**M. Ambrozas[✉], A. Juodagalvis[✉], A. Rinkevicius[✉], G. Tamulaitis[✉]**National Centre for Particle Physics, Universiti Malaya, Kuala Lumpur,
Malaysia**N. Bin Norjoharuddeen[✉], I. Yusuff[✉]⁵⁶, Z. Zolkapli**Universidad de Sonora (UNISON), Hermosillo, Mexico**J.F. Benitez[✉], A. Castaneda Hernandez[✉], H.A. Encinas Acosta, L.G. Gallegos Maríñez,
M. León Coello[✉], J.A. Murillo Quijada[✉], A. Sehrawat[✉], L. Valencia Palomo[✉]**Centro de Investigacion y de Estudios Avanzados del IPN, Mexico City,
Mexico**G. Ayala[✉], H. Castilla-Valdez[✉], E. De La Cruz-Burelo[✉], I. Heredia-De La Cruz[✉]⁵⁷,
R. Lopez-Fernandez[✉], C.A. Mondragon Herrera, D.A. Perez Navarro[✉], A. Sánchez Hernández[✉]**Universidad Iberoamericana, Mexico City, Mexico**C. Oropeza Barrera[✉], M. Ramírez García[✉]

Benemerita Universidad Autonoma de Puebla, Puebla, MexicoI. Bautista , I. Pedraza , H.A. Salazar Ibarguen , C. Uribe Estrada **University of Montenegro, Podgorica, Montenegro**I. Bubanja, N. Raicevic **University of Canterbury, Christchurch, New Zealand**P.H. Butler **National Centre for Physics, Quaid-I-Azam University, Islamabad, Pakistan**A. Ahmad , M.I. Asghar, A. Awais , M.I.M. Awan, H.R. Hoorani , W.A. Khan **AGH University of Krakow, Faculty of Computer Science, Electronics and Telecommunications, Krakow, Poland**V. Avati, L. Grzanka , M. Malawski **National Centre for Nuclear Research, Swierk, Poland**H. Bialkowska , M. Bluj , B. Boimska , M. Górski , M. Kazana , M. Szleper , P. Zalewski **Institute of Experimental Physics, Faculty of Physics, University of Warsaw, Warsaw, Poland**K. Bunkowski , K. Doroba , A. Kalinowski , M. Konecki , J. Krolikowski , A. Muhammad **Warsaw University of Technology, Warsaw, Poland**K. Pozniak , W. Zabolotny **Laboratório de Instrumentação e Física Experimental de Partículas, Lisboa, Portugal**M. Araujo , D. Bastos , C. Beirão Da Cruz E Silva , A. Boletti , M. Bozzo , P. Faccioli , M. Gallinaro , J. Hollar , N. Leonardo , T. Niknejad , M. Pisano , J. Seixas , J. Varela **Faculty of Physics, University of Belgrade, Belgrade, Serbia**P. Adzic , P. Milenovic **VINCA Institute of Nuclear Sciences, University of Belgrade, Belgrade, Serbia**M. Dordevic , J. Milosevic , V. Rekovic**Centro de Investigaciones Energéticas Medioambientales y Tecnológicas (CIEMAT), Madrid, Spain**M. Aguilar-Benitez, J. Alcaraz Maestre , M. Barrio Luna, Cristina F. Bedoya , M. Cepeda , M. Cerrada , N. Colino , B. De La Cruz , A. Delgado Peris , D. Fernández Del Val , J.P. Fernández Ramos , J. Flix , M.C. Fouz , O. Gonzalez Lopez , S. Goy Lopez , J.M. Hernandez , M.I. Josa , J. León Holgado , D. Moran , C. M. Morcillo Perez 

Á. Navarro Tobar , C. Perez Dengra , A. Pérez-Calero Yzquierdo , J. Puerta Pelayo , I. Redondo , D.D. Redondo Ferrero , L. Romero, S. Sánchez Navas , L. Urda Gómez , J. Vazquez Escobar , C. Willmott

Universidad Autónoma de Madrid, Madrid, Spain

J.F. de Trocóniz 

Universidad de Oviedo, Instituto Universitario de Ciencias y Tecnologías Espaciales de Asturias (ICTEA), Oviedo, Spain

B. Alvarez Gonzalez , J. Cuevas , J. Fernandez Menendez , S. Folgueras , I. Gonzalez Caballero , J.R. González Fernández , E. Palencia Cortezon , C. Ramón Álvarez , V. Rodríguez Bouza , A. Soto Rodriguez , A. Trapote , C. Vico Villalba , P. Vischia

Instituto de Física de Cantabria (IFCA), CSIC-Universidad de Cantabria, Santander, Spain

S. Bhowmik , S. Blanco Fernández , J.A. Brochero Cifuentes , I.J. Cabrillo , A. Calderon , J. Duarte Campderros , M. Fernandez , C. Fernandez Madrazo , G. Gomez , C. Lasaosa García , C. Martinez Rivero , P. Martinez Ruiz del Arbol , F. Matorras , P. Matorras Cuevas , E. Navarrete Ramos , J. Piedra Gomez , C. Prieels, L. Scodellaro , I. Vila , J.M. Vizan Garcia

University of Colombo, Colombo, Sri Lanka

M.K. Jayananda , B. Kailasapathy ⁵⁸, D.U.J. Sonnadara , D.D.C. Wickramarathna 

University of Ruhuna, Department of Physics, Matara, Sri Lanka

W.G.D. Dharmaratna ⁵⁹, K. Liyanage , N. Perera , N. Wickramage 

CERN, European Organization for Nuclear Research, Geneva, Switzerland

D. Abbaneo , C. Amendola , E. Auffray , G. Auzinger , J. Baechler, D. Barney , A. Bermúdez Martínez , M. Bianco , B. Bilin , A.A. Bin Anuar , A. Bocci , E. Brondolin , C. Caillol , T. Camporesi , G. Cerminara , N. Chernyavskaya , D. d'Enterria , A. Dabrowski , A. David , A. De Roeck , M.M. Defranchis , M. Deile , M. Dobson , F. Fallavollita ⁶⁰, L. Forthomme , G. Franzoni , W. Funk , S. Giani, D. Gigi, K. Gill , F. Glege , L. Gouskos , M. Haranko , J. Hegeman , V. Innocente , T. James , P. Janot , J. Kieseler , N. Kratochwil , S. Laurila , P. Lecoq , E. Leutgeb , C. Lourenço , B. Maier , L. Malgeri , M. Mannelli , A.C. Marini , F. Meijers , S. Mersi , E. Meschi , V. Milosevic , F. Moortgat , M. Mulders , S. Orfanelli, F. Pantaleo , M. Peruzzi , A. Petrilli , G. Petrucciani , A. Pfeiffer , M. Pierini , D. Piparo , H. Qu , D. Rabady , G. Reales Gutierrez, M. Rovere , H. Sakulin , S. Scarfi , M. Selvaggi , A. Sharma , K. Shchelina , P. Silva , P. Sphicas ⁶¹, A.G. Stahl Leiton , A. Steen , S. Summers , D. Treille , P. Tropea , A. Tsirou, D. Walter , J. Wanczyk ⁶², K.A. Wozniak ⁶³, P. Zehetner , P. Zejdl , W.D. Zeuner

Paul Scherrer Institut, Villigen, Switzerland

T. Bevilacqua ⁶⁴, L. Caminada ⁶⁴, A. Ebrahimi , W. Erdmann , R. Horisberger , Q. Ingram , H.C. Kaestli , D. Kotlinski , C. Lange , M. Missiroli ⁶⁴, L. Noehte ⁶⁴, T. Rohe

ETH Zurich — Institute for Particle Physics and Astrophysics (IPA), Zurich, Switzerland

T.K. Arrestad , K. Androsov ⁶², M. Backhaus , A. Calandri , C. Cazzaniga , K. Datta , A. De Cosa , G. Dissertori , M. Dittmar, M. Donegà , F. Eble , M. Galli , K. Gedia , F. Glessgen , C. Grab , D. Hits , W. Lustermann , A.-M. Lyon , R.A. Manzoni ⁶², M. Marchegiani , L. Marchese , C. Martin Perez , A. Mascellani ⁶², F. Nessi-Tedaldi , F. Pauss , V. Perovic , S. Pigazzini , M.G. Ratti , M. Reichmann , C. Reissel , T. Reitenspiess , B. Ristic , F. Riti , D. Ruini, D.A. Sanz Becerra , R. Seidita , J. Steggemann ⁶², D. Valsecchi , R. Wallny

Universität Zürich, Zurich, Switzerland

C. Amsler ⁶⁵, P. Bärtschi , C. Botta , D. Brzhechko, M.F. Canelli , K. Cormier , A. De Wit , R. Del Burgo, J.K. Heikkilä , M. Huwiler , W. Jin , A. Jofrehei , B. Kilminster , S. Leontsinis , S.P. Liechti , A. Macchiolo , P. Meiring , V.M. Mikuni , U. Molinatti , I. Neutelings , A. Reimers , P. Robmann, S. Sanchez Cruz , K. Schweiger , M. Senger , Y. Takahashi

National Central University, Chung-Li, Taiwan

C. Adloff ⁶⁶, C.M. Kuo, W. Lin, P.K. Rout , P.C. Tiwari ³⁹, S.S. Yu

National Taiwan University (NTU), Taipei, Taiwan

L. Ceard, Y. Chao , K.F. Chen , P.s. Chen, Z.g. Chen, W.-S. Hou , T.h. Hsu, Y.w. Kao, R. Khurana, G. Kole , Y.y. Li , R.-S. Lu , E. Paganis , A. Psallidas, X.f. Su, J. Thomas-Wilsker , H.y. Wu, E. Yazgan

High Energy Physics Research Unit, Department of Physics, Faculty of Science, Chulalongkorn University, Bangkok, Thailand

C. Asawatangtrakuldee , N. Srimanobhas , V. Wachirapusanand

Çukurova University, Physics Department, Science and Art Faculty, Adana, Turkey

D. Agyel , F. Boran , Z.S. Demiroglu , F. Dolek , I. Dumanoglu ⁶⁷, E. Eskut , Y. Guler ⁶⁸, E. Gurpinar Guler ⁶⁸, C. Isik , O. Kara, A. Kayis Topaksu , U. Kiminsu , G. Onengut , K. Ozdemir ⁶⁹, A. Polatoz , B. Tali ⁷⁰, U.G. Tok , S. Turkcapar , E. Uslan , I.S. Zorbakir

Middle East Technical University, Physics Department, Ankara, Turkey

K. Ocalan ⁷¹, M. Yalvac ⁷²

Bogazici University, Istanbul, Turkey

B. Akgun , I.O. Atakisi , E. Gülmmez , M. Kaya ⁷³, O. Kaya ⁷⁴, S. Tekten ⁷⁵

Istanbul Technical University, Istanbul, TurkeyA. Cakir , K. Cankocak ⁶⁷, Y. Komurcu , S. Sen ⁷⁶**Istanbul University, Istanbul, Turkey**O. Aydilek , S. Cerci ⁷⁰, V. Epshteyn , B. Hacisahinoglu , I. Hos ⁷⁷, B. Isildak ⁷⁸,
B. Kaynak , S. Ozkorucuklu , H. Sert , C. Simsek , D. Sunar Cerci ⁷⁰, C. Zorbilmez **Institute for Scintillation Materials of National Academy of Science of Ukraine, Kharkiv, Ukraine**A. Boyaryntsev , B. Grynyov **National Science Centre, Kharkiv Institute of Physics and Technology, Kharkiv, Ukraine**L. Levchuk **University of Bristol, Bristol, United Kingdom**D. Anthony , J.J. Brooke , A. Bundock , F. Bury , E. Clement , D. Cussans ,
H. Flacher , M. Glowacki, J. Goldstein , H.F. Heath , L. Kreczko , B. Krikler ,
S. Paramesvaran , S. Seif El Nasr-Storey, V.J. Smith , N. Stylianou ⁷⁹, K. Walkingshaw Pass,
R. White **Rutherford Appleton Laboratory, Didcot, United Kingdom**A.H. Ball, K.W. Bell , A. Belyaev ⁸⁰, C. Brew , R.M. Brown , D.J.A. Cockerill ,
C. Cooke , K.V. Ellis, K. Harder , S. Harper , M.-L. Holmberg ⁸¹, Sh. Jain , J. Linacre ,
K. Manolopoulos, D.M. Newbold , E. Olaiya, D. Petyt , T. Reis , G. Salvi , T. Schuh,
C.H. Shepherd-Themistocleous , I.R. Tomalin , T. Williams **Imperial College, London, United Kingdom**R. Bainbridge , P. Bloch , C.E. Brown , O. Buchmuller, V. Cacchio,
C.A. Carrillo Montoya , G.S. Chahal ⁸², D. Colling , J.S. Dancu, P. Dauncey , G. Davies ,
J. Davies, M. Della Negra , S. Fayer, G. Fedi , G. Hall , M.H. Hassanshahi , A. Howard,
G. Iles , M. Knight , J. Langford , L. Lyons , A.-M. Magnan , S. Malik, A. Martelli ,
M. Mieskolainen , J. Nash ⁸³, M. Pesaresi, B.C. Radburn-Smith , A. Richards, A. Rose ,
C. Seez , R. Shukla , A. Tapper , K. Uchida , G.P. Uttley , L.H. Vage, T. Virdee ²⁸,
M. Vojinovic , N. Wardle , D. Winterbottom **Brunel University, Uxbridge, United Kingdom**K. Coldham, J.E. Cole , A. Khan, P. Kyberd , I.D. Reid **Baylor University, Waco, Texas, U.S.A.**S. Abdullin , A. Brinkerhoff , B. Caraway , J. Dittmann , K. Hatakeyama ,
J. Hiltbrand , A.R. Kanuganti , B. McMaster , M. Saunders , S. Sawant ,
C. Sutantawibul , M. Toms , J. Wilson 

Catholic University of America, Washington, DC, U.S.A.

R. Bartek^{ID}, A. Dominguez^{ID}, C. Huerta Escamilla, A.E. Simsek^{ID}, R. Uniyal^{ID},
A.M. Vargas Hernandez^{ID}

The University of Alabama, Tuscaloosa, Alabama, U.S.A.

R. Chudasama^{ID}, S.I. Cooper^{ID}, S.V. Gleyzer^{ID}, C.U. Perez^{ID}, P. Rumerio^{ID}⁸⁴, E. Usai^{ID},
C. West^{ID}, R. Yi^{ID}

Boston University, Boston, Massachusetts, U.S.A.

A. Akpinar^{ID}, A. Albert^{ID}, D. Arcaro^{ID}, C. Cosby^{ID}, Z. Demiragli^{ID}, C. Erice^{ID}, E. Fontanesi^{ID},
D. Gastler^{ID}, J. Rohlf^{ID}, K. Salyer^{ID}, D. Sperka^{ID}, D. Spitzbart^{ID}, I. Suarez^{ID}, A. Tsatsos^{ID},
S. Yuan^{ID}

Brown University, Providence, Rhode Island, U.S.A.

G. Benelli^{ID}, X. Coubez²³, D. Cutts^{ID}, M. Hadley^{ID}, U. Heintz^{ID}, J.M. Hogan^{ID}⁸⁵, T. Kwon^{ID},
G. Landsberg^{ID}, K.T. Lau^{ID}, D. Li^{ID}, J. Luo^{ID}, S. Mondal^{ID}, M. Narain^{ID}[†], N. Pervan^{ID},
S. Sagir^{ID}⁸⁶, F. Simpson^{ID}, W.Y. Wong, X. Yan^{ID}, W. Zhang

University of California, Davis, Davis, California, U.S.A.

S. Abbott^{ID}, J. Bonilla^{ID}, C. Brainerd^{ID}, R. Breedon^{ID}, M. Calderon De La Barca Sanchez^{ID},
M. Chertok^{ID}, M. Citron^{ID}, J. Conway^{ID}, P.T. Cox^{ID}, R. Erbacher^{ID}, G. Haza^{ID}, F. Jensen^{ID},
O. Kukral^{ID}, G. Mocellin^{ID}, M. Mulhearn^{ID}, D. Pellett^{ID}, B. Regnery^{ID}, W. Wei^{ID}, Y. Yao^{ID},
F. Zhang^{ID}

University of California, Los Angeles, California, U.S.A.

M. Bachtis^{ID}, R. Cousins^{ID}, A. Datta^{ID}, J. Hauser^{ID}, M. Ignatenko^{ID}, M.A. Iqbal^{ID}, T. Lam^{ID},
E. Manca^{ID}, W.A. Nash^{ID}, D. Saltzberg^{ID}, B. Stone^{ID}, V. Valuev^{ID}

University of California, Riverside, Riverside, California, U.S.A.

R. Clare^{ID}, M. Gordon, G. Hanson^{ID}, W. Si^{ID}, S. Wimpenny^{ID}[†]

University of California, San Diego, La Jolla, California, U.S.A.

J.G. Branson^{ID}, S. Cittolin^{ID}, S. Cooperstein^{ID}, D. Diaz^{ID}, J. Duarte^{ID}, R. Gerosa^{ID},
L. Giannini^{ID}, J. Guiang^{ID}, R. Kansal^{ID}, V. Krutelyov^{ID}, R. Lee^{ID}, J. Letts^{ID},
M. Masciovecchio^{ID}, F. Mokhtar^{ID}, M. Pieri^{ID}, M. Quinnan^{ID}, B.V. Sathia Narayanan^{ID},
V. Sharma^{ID}, M. Tadel^{ID}, E. Vourliotis^{ID}, F. Würthwein^{ID}, Y. Xiang^{ID}, A. Yagil^{ID}

University of California, Santa Barbara — Department of Physics, Santa Barbara, California, U.S.A.

L. Brennan^{ID}, C. Campagnari^{ID}, G. Collura^{ID}, A. Dorsett^{ID}, J. Incandela^{ID}, M. Kilpatrick^{ID},
J. Kim^{ID}, A.J. Li^{ID}, P. Masterson^{ID}, H. Mei^{ID}, M. Oshiro^{ID}, J. Richman^{ID}, U. Sarica^{ID},
R. Schmitz^{ID}, F. Setti^{ID}, J. Sheplock^{ID}, D. Stuart^{ID}, S. Wang^{ID}

California Institute of Technology, Pasadena, California, U.S.A.

A. Bornheim^{ID}, O. Cerri, A. Latorre, J.M. Lawhorn^{ID}, J. Mao^{ID}, H.B. Newman^{ID},
T. Q. Nguyen^{ID}, M. Spiropulu^{ID}, J.R. Vlimant^{ID}, C. Wang^{ID}, S. Xie^{ID}, R.Y. Zhu^{ID}

Carnegie Mellon University, Pittsburgh, Pennsylvania, U.S.A.

J. Alison^{ID}, S. An^{ID}, M.B. Andrews^{ID}, P. Bryant^{ID}, V. Dutta^{ID}, T. Ferguson^{ID}, A. Harilal^{ID}, C. Liu^{ID}, T. Mudholkar^{ID}, S. Murthy^{ID}, M. Paulini^{ID}, A. Roberts^{ID}, A. Sanchez^{ID}, W. Terrill^{ID}

University of Colorado Boulder, Boulder, Colorado, U.S.A.

J.P. Cumalat^{ID}, W.T. Ford^{ID}, A. Hassani^{ID}, G. Karathanasis^{ID}, E. MacDonald, N. Manganelli^{ID}, F. Marini^{ID}, A. Perloff^{ID}, C. Savard^{ID}, N. Schonbeck^{ID}, K. Stenson^{ID}, K.A. Ulmer^{ID}, S.R. Wagner^{ID}, N. Zipper^{ID}

Cornell University, Ithaca, New York, U.S.A.

J. Alexander^{ID}, S. Bright-Thonney^{ID}, X. Chen^{ID}, D.J. Cranshaw^{ID}, J. Fan^{ID}, X. Fan^{ID}, D. Gadkari^{ID}, S. Hogan^{ID}, J. Monroy^{ID}, J.R. Patterson^{ID}, J. Reichert^{ID}, M. Reid^{ID}, A. Ryd^{ID}, J. Thom^{ID}, P. Wittich^{ID}, R. Zou^{ID}

Fermi National Accelerator Laboratory, Batavia, Illinois, U.S.A.

M. Albrow^{ID}, M. Alyari^{ID}, O. Amram^{ID}, G. Apollinari^{ID}, A. Apresyan^{ID}, L.A.T. Bauerick^{ID}, D. Berry^{ID}, J. Berryhill^{ID}, P.C. Bhat^{ID}, K. Burkett^{ID}, J.N. Butler^{ID}, A. Canepa^{ID}, G.B. Cerati^{ID}, H.W.K. Cheung^{ID}, F. Chlebana^{ID}, G. Cummings^{ID}, J. Dickinson^{ID}, I. Dutta^{ID}, V.D. Elvira^{ID}, Y. Feng^{ID}, J. Freeman^{ID}, A. Gandrakota^{ID}, Z. Gecse^{ID}, L. Gray^{ID}, D. Green, S. Grünendahl^{ID}, D. Guerrero^{ID}, O. Gutsche^{ID}, R.M. Harris^{ID}, R. Heller^{ID}, T.C. Herwig^{ID}, J. Hirschauer^{ID}, L. Horyn^{ID}, B. Jayatilaka^{ID}, S. Jindariani^{ID}, M. Johnson^{ID}, U. Joshi^{ID}, T. Klijsma^{ID}, B. Klima^{ID}, K.H.M. Kwok^{ID}, S. Lammel^{ID}, D. Lincoln^{ID}, R. Lipton^{ID}, T. Liu^{ID}, C. Madrid^{ID}, K. Maeshima^{ID}, C. Mantilla^{ID}, D. Mason^{ID}, P. McBride^{ID}, P. Merkel^{ID}, S. Mrenna^{ID}, S. Nahn^{ID}, J. Ngadiuba^{ID}, D. Noonan^{ID}, V. Papadimitriou^{ID}, N. Pastika^{ID}, K. Pedro^{ID}, C. Pena^{ID}⁸⁷, F. Ravera^{ID}, A. Reinsvold Hall^{ID}⁸⁸, L. Ristori^{ID}, E. Sexton-Kennedy^{ID}, N. Smith^{ID}, A. Soha^{ID}, L. Spiegel^{ID}, S. Stoynev^{ID}, J. Strait^{ID}, L. Taylor^{ID}, S. Tkaczyk^{ID}, N.V. Tran^{ID}, L. Uplegger^{ID}, E.W. Vaandering^{ID}, I. Zoi^{ID}

University of Florida, Gainesville, Florida, U.S.A.

C. Aruta^{ID}, P. Avery^{ID}, D. Bourilkov^{ID}, L. Cadamuro^{ID}, P. Chang^{ID}, V. Cherepanov^{ID}, R.D. Field, E. Koenig^{ID}, M. Kolosova^{ID}, J. Konigsberg^{ID}, A. Korytov^{ID}, K.H. Lo, K. Matchev^{ID}, N. Menendez^{ID}, G. Mitselmakher^{ID}, A. Muthirakalayil Madhu^{ID}, N. Rawal^{ID}, D. Rosenzweig^{ID}, S. Rosenzweig^{ID}, K. Shi^{ID}, J. Wang^{ID}

Florida State University, Tallahassee, Florida, U.S.A.

T. Adams^{ID}, A. Al Kadhim^{ID}, A. Askew^{ID}, N. Bower^{ID}, R. Habibullah^{ID}, V. Hagopian^{ID}, R. Hashmi^{ID}, R.S. Kim^{ID}, S. Kim^{ID}, T. Kolberg^{ID}, G. Martinez, H. Prosper^{ID}, P.R. Prova, O. Viazlo^{ID}, M. Wulansatiti^{ID}, R. Yohay^{ID}, J. Zhang

Florida Institute of Technology, Melbourne, Florida, U.S.A.

B. Alsufyani, M.M. Baarmand^{ID}, S. Butalla^{ID}, T. Elkafrawy^{ID}⁵³, M. Hohlmann^{ID}, R. Kumar Verma^{ID}, M. Rahmani, F. Yumiceva^{ID}

University of Illinois Chicago, Chicago, U.S.A., Chicago, U.S.A.

M.R. Adams^{ID}, C. Bennett, R. Cavanaugh^{ID}, S. Dittmer^{ID}, R. Escobar Franco^{ID}, O. Evdokimov^{ID}, C.E. Gerber^{ID}, D.J. Hofman^{ID}, J.h. Lee^{ID}, D. S. Lemos^{ID}, A.H. Merrit^{ID},

C. Mills^{1D}, S. Nanda^{1D}, G. Oh^{1D}, B. Ozek^{1D}, D. Pilipovic^{1D}, T. Roy^{1D}, S. Rudrabhatla^{1D}, M.B. Tonjes^{1D}, N. Varelas^{1D}, X. Wang^{1D}, Z. Ye^{1D}, J. Yoo^{1D}

The University of Iowa, Iowa City, Iowa, U.S.A.

M. Alhusseini^{1D}, D. Blend, K. Dilsiz^{1D}⁸⁹, L. Emediato^{1D}, G. Karaman^{1D}, O.K. Köseyan^{1D}, J.-P. Merlo, A. Mestvirishvili^{1D}⁹⁰, J. Nachtman^{1D}, O. Neogi, H. Ogul^{1D}⁹¹, Y. Onel^{1D}, A. Penzo^{1D}, C. Snyder, E. Tiras^{1D}⁹²

Johns Hopkins University, Baltimore, Maryland, U.S.A.

B. Blumenfeld^{1D}, L. Corcodilos^{1D}, J. Davis^{1D}, A.V. Gritsan^{1D}, L. Kang^{1D}, S. Kyriacou^{1D}, P. Maksimovic^{1D}, M. Roguljic^{1D}, J. Roskes^{1D}, S. Sekhar^{1D}, M. Swartz^{1D}, T.Á. Vámi^{1D}

The University of Kansas, Lawrence, Kansas, U.S.A.

A. Abreu^{1D}, L.F. Alcerro Alcerro^{1D}, J. Anguiano^{1D}, P. Baringer^{1D}, A. Bean^{1D}, Z. Flowers^{1D}, D. Grove, J. King^{1D}, G. Krintiras^{1D}, M. Lazarovits^{1D}, C. Le Mahieu^{1D}, C. Lindsey, J. Marquez^{1D}, N. Minafra^{1D}, M. Murray^{1D}, M. Nickel^{1D}, M. Pitt^{1D}, S. Popescu^{1D}⁹³, C. Rogan^{1D}, C. Royon^{1D}, R. Salvatico^{1D}, S. Sanders^{1D}, C. Smith^{1D}, Q. Wang^{1D}, G. Wilson^{1D}

Kansas State University, Manhattan, Kansas, U.S.A.

B. Allmond^{1D}, A. Ivanov^{1D}, K. Kaadze^{1D}, A. Kalogeropoulos^{1D}, D. Kim, Y. Maravin^{1D}, K. Nam, J. Natoli^{1D}, D. Roy^{1D}, G. Sorrentino^{1D}

Lawrence Livermore National Laboratory, Livermore, California, U.S.A.

F. Rebassoo^{1D}, D. Wright^{1D}

University of Maryland, College Park, Maryland, U.S.A.

E. Adams^{1D}, A. Baden^{1D}, O. Baron, A. Belloni^{1D}, A. Bethani^{1D}, Y.m. Chen^{1D}, S.C. Eno^{1D}, N.J. Hadley^{1D}, S. Jabeen^{1D}, R.G. Kellogg^{1D}, T. Koeth^{1D}, Y. Lai^{1D}, S. Lascio^{1D}, A.C. Mignerey^{1D}, S. Nabili^{1D}, C. Palmer^{1D}, C. Papageorgakis^{1D}, M.M. Paranjpe, L. Wang^{1D}, K. Wong^{1D}

Massachusetts Institute of Technology, Cambridge, Massachusetts, U.S.A.

J. Bendavid^{1D}, W. Busza^{1D}, I.A. Cali^{1D}, Y. Chen^{1D}, M. D'Alfonso^{1D}, J. Eysermans^{1D}, C. Freer^{1D}, G. Gomez-Ceballos^{1D}, M. Goncharov, P. Harris, D. Hoang, D. Kovalskyi^{1D}, J. Krupa^{1D}, L. Lavezzi^{1D}, Y.-J. Lee^{1D}, K. Long^{1D}, C. Mironov^{1D}, C. Paus^{1D}, D. Rankin^{1D}, C. Roland^{1D}, G. Roland^{1D}, S. Rothman^{1D}, Z. Shi^{1D}, G.S.F. Stephans^{1D}, J. Wang, Z. Wang^{1D}, B. Wyslouch^{1D}, T. J. Yang^{1D}

University of Minnesota, Minneapolis, Minnesota, U.S.A.

B. Crossman^{1D}, B.M. Joshi^{1D}, C. Kapsiak^{1D}, M. Krohn^{1D}, D. Mahon^{1D}, J. Mans^{1D}, S. Pandey^{1D}, M. Revering^{1D}, R. Rusack^{1D}, R. Saradhy^{1D}, N. Schroeder^{1D}, N. Strobbe^{1D}, M.A. Wadud^{1D}

University of Mississippi, Oxford, Mississippi, U.S.A.

L.M. Cremaldi^{1D}

University of Nebraska-Lincoln, Lincoln, Nebraska, U.S.A.

K. Bloom , M. Bryson, D.R. Claes , C. Fangmeier , F. Golf , J. Hossain , C. Joo , I. Kravchenko , I. Reed , J.E. Siado , G.R. Snow[†], W. Tabb , A. Wightman , F. Yan , D. Yu , A.G. Zecchinelli 

State University of New York at Buffalo, Buffalo, New York, U.S.A.

G. Agarwal , H. Bandyopadhyay , L. Hay , I. Iashvili , A. Kharchilava , C. McLean , M. Morris , D. Nguyen , J. Pekkanen , S. Rappoccio , H. Rejeb Sfar, A. Williams 

Northeastern University, Boston, Massachusetts, U.S.A.

G. Alverson , E. Barberis , Y. Haddad , Y. Han , A. Krishna , J. Li , M. Lu , G. Madigan , B. Marzocchi , D.M. Morse , V. Nguyen , T. Orimoto , A. Parker , L. Skinnari , A. Tishelman-Charny , B. Wang , D. Wood 

Northwestern University, Evanston, Illinois, U.S.A.

S. Bhattacharya , J. Bueghly, Z. Chen , K.A. Hahn , Y. Liu , Y. Miao , D.G. Monk , M.H. Schmitt , A. Taliercio , M. Velasco

University of Notre Dame, Notre Dame, Indiana, U.S.A.

R. Band , R. Bucci, S. Castells , M. Cremonesi, A. Das , R. Goldouzian , M. Hildreth , K.W. Ho , K. Hurtado Anampa , C. Jessop , K. Lannon , J. Lawrence , N. Loukas , L. Lutton , J. Mariano, N. Marinelli, I. Mcalister, T. McCauley , C. McGrady , K. Mohrman , C. Moore , Y. Musienko ¹³, H. Nelson , M. Osherson , R. Ruchti , A. Townsend , M. Wayne , H. Yockey, M. Zarucki , L. Zygalas 

The Ohio State University, Columbus, Ohio, U.S.A.

A. Basnet , B. Bylsma, M. Carrigan , L.S. Durkin , C. Hill , M. Joyce , A. Lesauvage , M. Nunez Ornelas , K. Wei, B.L. Winer , B. R. Yates 

Princeton University, Princeton, New Jersey, U.S.A.

F.M. Addesa , H. Bouchamaoui , P. Das , G. Dezoort , P. Elmer , A. Frankenthal , B. Greenberg , N. Haubrich , S. Higginbotham , G. Kopp , S. Kwan , D. Lange , A. Loeliger , D. Marlow , I. Ojalvo , J. Olsen , D. Stickland , C. Tully 

University of Puerto Rico, Mayaguez, Puerto Rico, U.S.A.

S. Malik 

Purdue University, West Lafayette, Indiana, U.S.A.

A.S. Bakshi , V.E. Barnes , S. Chandra , R. Chawla , S. Das , A. Gu , L. Gutay, M. Jones , A.W. Jung , D. Kondratyev , A.M. Koshy, M. Liu , G. Negro , N. Neumeister , G. Paspalaki , S. Piperov , A. Purohit , J.F. Schulte , M. Stojanovic ¹⁷, J. Thieman , A. K. Virdi , F. Wang , W. Xie 

Purdue University Northwest, Hammond, Indiana, U.S.A.

J. Dolen , N. Parashar , A. Pathak 

Rice University, Houston, Texas, U.S.A.

- D. Acosta , A. Baty , T. Carnahan , S. Dildick , K.M. Ecklund ,
 P.J. Fernández Manteca , S. Freed, P. Gardner, F.J.M. Geurts , A. Kumar , W. Li ,
 O. Miguel Colin , B.P. Padley , R. Redjimi, J. Rotter , E. Yigitbasi , Y. Zhang 

University of Rochester, Rochester, New York, U.S.A.

- A. Bodek , P. de Barbaro , R. Demina , J.L. Dulemba , C. Fallon, A. Garcia-Bellido ,
 O. Hindrichs , A. Khukhunaishvili , P. Parygin , E. Popova , R. Taus , G.P. Van Onsem 

The Rockefeller University, New York, New York, U.S.A.

- K. Goulianos 

Rutgers, The State University of New Jersey, Piscataway, New Jersey, U.S.A.

- B. Chiarito, J.P. Chou , Y. Gershtein , E. Halkiadakis , A. Hart , M. Heindl ,
 D. Jaroslawski , O. Karacheban ²⁶, I. Laflotte , A. Lath , R. Montalvo, K. Nash,
 H. Routray , S. Salur , S. Schnetzer, S. Somalwar , R. Stone , S.A. Thayil , S. Thomas,
 J. Vora , H. Wang 

University of Tennessee, Knoxville, Tennessee, U.S.A.

- H. Acharya, D. Ally , A.G. Delannoy , S. Fiorendi , T. Holmes , N. Karunarathna ,
 L. Lee , E. Nibigira , S. Spanier 

Texas A&M University, College Station, Texas, U.S.A.

- D. Aebi , M. Ahmad , O. Bouhali ⁹⁴, M. Dalchenko , R. Eusebi , J. Gilmore ,
 T. Huang , T. Kamon ⁹⁵, H. Kim , S. Luo , S. Malhotra, R. Mueller , D. Overton ,
 D. Rathjens , A. Safonov 

Texas Tech University, Lubbock, Texas, U.S.A.

- N. Akchurin , J. Damgov , V. Hegde , A. Hussain , Y. Kazhykarim, K. Lamichhane ,
 S.W. Lee , A. Mankel , T. Mengke, S. Muthumuni , T. Peltola , I. Volobouev ,
 A. Whitbeck 

Vanderbilt University, Nashville, Tennessee, U.S.A.

- E. Appelt , S. Greene, A. Gurrola , W. Johns , R. Kunnavalkam Elayavalli , A. Melo ,
 F. Romeo , P. Sheldon , S. Tuo , J. Velkovska , J. Viinikainen 

University of Virginia, Charlottesville, Virginia, U.S.A.

- B. Cardwell , B. Cox , J. Hakala , R. Hirosky , A. Ledovskoy , A. Li , C. Neu ,
 C.E. Perez Lara 

Wayne State University, Detroit, Michigan, U.S.A.

- P.E. Karchin 

University of Wisconsin — Madison, Madison, Wisconsin, U.S.A.

- A. Aravind, S. Banerjee , K. Black , T. Bose , S. Dasu , I. De Bruyn , P. Everaerts ,
 C. Galloni, H. He , M. Herndon , A. Herve , C.K. Koraka , A. Lanaro, R. Loveless 

J. Madhusudanan Sreekala¹⁰, A. Mallampalli¹⁰, A. Mohammadi¹⁰, S. Mondal, G. Parida¹⁰, D. Pinna, A. Savin, V. Shang¹⁰, V. Sharma¹⁰, W.H. Smith¹⁰, D. Teague, H.F. Tsoi¹⁰, W. Vetens¹⁰, A. Warden¹⁰

Authors affiliated with an institute or an international laboratory covered by a cooperation agreement with CERN

S. Afanasiev¹⁰, V. Andreev¹⁰, Yu. Andreev¹⁰, T. Aushev¹⁰, M. Azarkin¹⁰, A. Babaev¹⁰, A. Belyaev¹⁰, V. Blinov⁹⁶, E. Boos¹⁰, V. Borshch¹⁰, D. Budkouski¹⁰, V. Bunichev¹⁰, M. Chadeeva¹⁰⁹⁶, V. Chekhovsky, R. Chistov¹⁰⁹⁶, A. Dermenev¹⁰, T. Dimova¹⁰⁹⁶, D. Druzhkin¹⁰⁹⁷, M. Dubinin¹⁰⁸⁷, L. Dudko¹⁰, A. Ershov¹⁰, G. Gavrilov¹⁰, V. Gavrilov¹⁰, S. Gninenko¹⁰, V. Golovtcov¹⁰, N. Golubev¹⁰, I. Golutvin¹⁰, I. Gorbunov¹⁰, A. Gribushin¹⁰, Y. Ivanov¹⁰, V. Kachanov¹⁰, L. Kardapoltsev¹⁰⁹⁶, V. Karjavine¹⁰, A. Karneyeu¹⁰, V. Kim¹⁰⁹⁶, M. Kirakosyan, D. Kirpichnikov¹⁰, M. Kirsanov¹⁰, V. Klyukhin¹⁰, O. Kodolova¹⁰⁹⁸, D. Konstantinov¹⁰, V. Korenkov¹⁰, A. Kozyrev¹⁰⁹⁶, N. Krasnikov¹⁰, A. Lanev¹⁰, P. Levchenko¹⁰⁹⁹, N. Lychkovskaya¹⁰, V. Makarenko¹⁰, A. Malakhov¹⁰, V. Matveev¹⁰⁹⁶, V. Murzin¹⁰, A. Nikitenko¹⁰^{100,98}, S. Obraztsov¹⁰, V. Oreshkin¹⁰, V. Palichik¹⁰, V. Perelygin¹⁰, S. Petrushanko¹⁰, S. Polikarpov¹⁰⁹⁶, V. Popov, O. Radchenko¹⁰⁹⁶, M. Savina¹⁰, V. Savrin¹⁰, D. Selivanova¹⁰, V. Shalaev¹⁰, S. Shmatov¹⁰, S. Shulha¹⁰, Y. Skovpen¹⁰⁹⁶, S. Slabospitskii¹⁰, V. Smirnov¹⁰, A. Snigirev¹⁰, D. Sosnov¹⁰, V. Sulimov¹⁰, E. Tcherniaev¹⁰, A. Terkulov¹⁰, O. Teryaev¹⁰, I. Tlisova¹⁰, A. Toropin¹⁰, L. Uvarov¹⁰, A. Uzunian¹⁰, A. Vorobyev[†], N. Voytishin¹⁰, B.S. Yuldashev¹⁰¹, A. Zarubin¹⁰, I. Zhizhin¹⁰, A. Zhokin¹⁰

[†] Deceased

¹ Also at Yerevan State University, Yerevan, Armenia

² Also at TU Wien, Vienna, Austria

³ Also at Institute of Basic and Applied Sciences, Faculty of Engineering, Arab Academy for Science, Technology and Maritime Transport, Alexandria, Egypt

⁴ Also at Ghent University, Ghent, Belgium

⁵ Also at Universidade Estadual de Campinas, Campinas, Brazil

⁶ Also at Federal University of Rio Grande do Sul, Porto Alegre, Brazil

⁷ Also at UFMS, Nova Andradina, Brazil

⁸ Also at Nanjing Normal University, Nanjing, China

⁹ Now at The University of Iowa, Iowa City, Iowa, U.S.A.

¹⁰ Also at University of Chinese Academy of Sciences, Beijing, China

¹¹ Also at University of Chinese Academy of Sciences, Beijing, China

¹² Also at Université Libre de Bruxelles, Bruxelles, Belgium

¹³ Also at an institute or an international laboratory covered by a cooperation agreement with CERN

¹⁴ Also at Suez University, Suez, Egypt

¹⁵ Now at British University in Egypt, Cairo, Egypt

¹⁶ Also at Birla Institute of Technology, Mesra, Mesra, India

¹⁷ Also at Purdue University, West Lafayette, Indiana, U.S.A.

¹⁸ Also at Université de Haute Alsace, Mulhouse, France

¹⁹ Also at Department of Physics, Tsinghua University, Beijing, China

²⁰ Also at The University of the State of Amazonas, Manaus, Brazil

²¹ Also at Erzincan Binali Yildirim University, Erzincan, Turkey

²² Also at University of Hamburg, Hamburg, Germany

²³ Also at RWTH Aachen University, III. Physikalisches Institut A, Aachen, Germany

²⁴ Also at Isfahan University of Technology, Isfahan, Iran

- ²⁵ Also at Bergische University Wuppertal (BUW), Wuppertal, Germany
²⁶ Also at Brandenburg University of Technology, Cottbus, Germany
²⁷ Also at Forschungszentrum Jülich, Juelich, Germany
²⁸ Also at CERN, European Organization for Nuclear Research, Geneva, Switzerland
²⁹ Also at Physics Department, Faculty of Science, Assiut University, Assiut, Egypt
³⁰ Also at Wigner Research Centre for Physics, Budapest, Hungary
³¹ Also at Institute of Physics, University of Debrecen, Debrecen, Hungary
³² Also at Institute of Nuclear Research ATOMKI, Debrecen, Hungary
³³ Now at Universitatea Babes-Bolyai — Facultatea de Fizica, Cluj-Napoca, Romania
³⁴ Also at Faculty of Informatics, University of Debrecen, Debrecen, Hungary
³⁵ Also at Punjab Agricultural University, Ludhiana, India
³⁶ Also at UPES — University of Petroleum and Energy Studies, Dehradun, India
³⁷ Also at University of Visva-Bharati, Santiniketan, India
³⁸ Also at University of Hyderabad, Hyderabad, India
³⁹ Also at Indian Institute of Science (IISc), Bangalore, India
⁴⁰ Also at IIT Bhubaneswar, Bhubaneswar, India
⁴¹ Also at Institute of Physics, Bhubaneswar, India
⁴² Also at Deutsches Elektronen-Synchrotron, Hamburg, Germany
⁴³ Now at Department of Physics, Isfahan University of Technology, Isfahan, Iran
⁴⁴ Also at Sharif University of Technology, Tehran, Iran
⁴⁵ Also at Department of Physics, University of Science and Technology of Mazandaran, Behshahr, Iran
⁴⁶ Also at Helwan University, Cairo, Egypt
⁴⁷ Also at Italian National Agency for New Technologies, Energy and Sustainable Economic Development, Bologna, Italy
⁴⁸ Also at Centro Siciliano di Fisica Nucleare e di Struttura Della Materia, Catania, Italy
⁴⁹ Also at Università degli Studi Guglielmo Marconi, Roma, Italy
⁵⁰ Also at Scuola Superiore Meridionale, Università di Napoli ‘Federico II’, Napoli, Italy
⁵¹ Also at Fermi National Accelerator Laboratory, Batavia, Illinois, U.S.A.
⁵² Also at Università di Napoli ‘Federico II’, Napoli, Italy
⁵³ Also at Ain Shams University, Cairo, Egypt
⁵⁴ Also at Consiglio Nazionale delle Ricerche — Istituto Officina dei Materiali, Perugia, Italy
⁵⁵ Also at Riga Technical University, Riga, Latvia
⁵⁶ Also at Department of Applied Physics, Faculty of Science and Technology, Universiti Kebangsaan Malaysia, Bangi, Malaysia
⁵⁷ Also at Consejo Nacional de Ciencia y Tecnología, Mexico City, Mexico
⁵⁸ Also at Trincomalee Campus, Eastern University, Sri Lanka, Nilaveli, Sri Lanka
⁵⁹ Also at Saegis Campus, Nugegoda, Sri Lanka
⁶⁰ Also at INFN Sezione di Pavia, Università di Pavia, Pavia, Italy
⁶¹ Also at National and Kapodistrian University of Athens, Athens, Greece
⁶² Also at Ecole Polytechnique Fédérale Lausanne, Lausanne, Switzerland
⁶³ Also at University of Vienna Faculty of Computer Science, Vienna, Austria
⁶⁴ Also at Universität Zürich, Zurich, Switzerland
⁶⁵ Also at Stefan Meyer Institute for Subatomic Physics, Vienna, Austria
⁶⁶ Also at Laboratoire d’Annecy-le-Vieux de Physique des Particules, IN2P3-CNRS, Annecy-le-Vieux, France
⁶⁷ Also at Near East University, Research Center of Experimental Health Science, Mersin, Turkey
⁶⁸ Also at Konya Technical University, Konya, Turkey
⁶⁹ Also at Izmir Bakircay University, Izmir, Turkey
⁷⁰ Also at Adiyaman University, Adiyaman, Turkey
⁷¹ Also at Necmettin Erbakan University, Konya, Turkey
⁷² Also at Bozok Üniversitesi Rektörlüğü, Yozgat, Turkey
⁷³ Also at Marmara University, Istanbul, Turkey

- ⁷⁴ Also at Milli Savunma University, Istanbul, Turkey
⁷⁵ Also at Kafkas University, Kars, Turkey
⁷⁶ Also at Hacettepe University, Ankara, Turkey
⁷⁷ Also at Istanbul University — Cerrahpasa, Faculty of Engineering, Istanbul, Turkey
⁷⁸ Also at Ozyegin University, Istanbul, Turkey
⁷⁹ Also at Vrije Universiteit Brussel, Brussel, Belgium
⁸⁰ Also at School of Physics and Astronomy, University of Southampton, Southampton, United Kingdom
⁸¹ Also at University of Bristol, Bristol, United Kingdom
⁸² Also at IPPP Durham University, Durham, United Kingdom
⁸³ Also at Monash University, Faculty of Science, Clayton, Australia
⁸⁴ Also at Università di Torino, Torino, Italy
⁸⁵ Also at Bethel University, St. Paul, Minnesota, U.S.A.
⁸⁶ Also at Karamanoğlu Mehmetbey University, Karaman, Turkey
⁸⁷ Also at California Institute of Technology, Pasadena, California, U.S.A.
⁸⁸ Also at United States Naval Academy, Annapolis, Maryland, U.S.A.
⁸⁹ Also at Bingol University, Bingol, Turkey
⁹⁰ Also at Georgian Technical University, Tbilisi, Georgia
⁹¹ Also at Sinop University, Sinop, Turkey
⁹² Also at Erciyes University, Kayseri, Turkey
⁹³ Also at Horia Hulubei National Institute of Physics and Nuclear Engineering (IFIN-HH), Bucharest, Romania
⁹⁴ Also at Texas A&M University at Qatar, Doha, Qatar
⁹⁵ Also at Kyungpook National University, Daegu, Korea
⁹⁶ Also at another institute or international laboratory covered by a cooperation agreement with CERN
⁹⁷ Also at Universiteit Antwerpen, Antwerpen, Belgium
⁹⁸ Also at Yerevan Physics Institute, Yerevan, Armenia
⁹⁹ Also at Northeastern University, Boston, Massachusetts, U.S.A.
¹⁰⁰ Also at Imperial College, London, United Kingdom
¹⁰¹ Also at Institute of Nuclear Physics of the Uzbekistan Academy of Sciences, Tashkent, Uzbekistan