Table 1. Pre-sedative NPO Guidelines

	Solid foods	Clear liquids
Adults	6-8 hr or none after midnight	2-3 hr
Children > 36 month	6-8 hr	2-3 hr
Children $6-36$ month	6 hr	2-3 hr
Children < 6 month	4-6 hr	2 hr

^{*:} American sociefy of anesthesiologists guidelines for sedation and analgesia by non-anesthesiologists.

: 110-744

28

:

Tel: 02-760-3042, Fax: 02-766-9427 E-mail: dentane @plaza.snu.ac.kr

가 Table 2. Sedation Score 1. Fully awake and orienated 2. Drowsy 3. Eyes closed but rousable to command 3 - 54. Eyes closed but rousable to mild physical stimulation 5. Eyes closed but unrousable to mild physical stimulation Anesth Analg 1992; 75: 708-14. scaling, curettage 가 가 가 가 가 가 nasal mask 가 가 N_2O 30 mmHg가 30 3 mmHg (megaloblastic anemia) 가 (< 25 ppm) 가 (sedation score, BIS (Bis-Table 2) 5-7) 가 pectral index) score A-line 가 nasal (Inhalational sedation) mask 가 가 (titration) nasal mask 가 가 100%

						49
가			가	. 가		
가 가	(non-rebreathing)					
가 가 가		,	·		13)	
, 100 ml/kg/min가	8)	70 kg				
가	7 L/min . 100% 6 L/min	,				
가	가		50 60%			
가 .		가	50 – 60 %		,	
가	20%		,	,	,	9)
10% 가	70% 가 가		,	,	,	
		가	,			
가 scaling currettag	ee .					
	, , 가	·	·	1980	10	·
가 , ,	,		10)			2
,	5% 가	2)				
가 .	3-5	100%				
	100%					
, , , 가	pethidine					
가		가	(CPR) 가	가 가	(Basic Life	Support)
가				•		(suction)

.

(가

· ,

1. Whitcher C, Zimmerman DC, Piziali RL: Control of occupational exposure to nitrous oxide in the oral

surgery office. J Oral Surgery 1978; 36: 431-40.

2. : Nitrous Oxide

2001; 1: 52-7.

- Barr G, Anderson RE, Jakobsson JG: A study of bispectral analysis and auditory evoked potential indices during propofol-induced hypnosis in volunteers: the effect of an episode of wakefulness on explicit and implicit memory. Anaesthesia 2001; 56: 888-93.
- Malamed SF: Sedation. A guide to patient management. 3rd ed. St. Louis, Mosby-Year Book Inc. 1995, pp 181-303.
- Layzer RB, Fishman RA, Schafer JA: Neuropathy following abuse of nitrous oxide. Neurology 1978; 28: 504-6.
- NIOSH Warns. Nitrous oxide continues to threaten health care workers. DHNS (NIOSH) Publication No. 94-118, 1994.
- Rowland AS, Baird DD, Weinberg CR, Shore DL, Shy CM, Wilcox AJ: Reduced fertility among women employed as dental assistants exposed to high levels of nitrous oxide. N Engl J Med 1992; 327: 993-7.
- Bain JA, Spoerel WE: Flow requirements for a modified Mapleson D system during controlled ventilation. Can Anaesth Soc J 1973; 20: 629-36.
- D'Eramo EM. Mortality and morbidity with outpatient anesthesia: the Massachusetts experience. J Oral Maxillofac Surg 1999; 57: 531-6.
- Coplans MP, Curson I: Deaths associated with dentistry and dental disease 1980-1989. Anaesthesia 1993;
 48: 435-8.