Short communication

Effects of water level and soil type on the survival and growth of *Persicaria thunbergii* during early growth stages

Do Hee Kim, Heung-Tae Kim, Jae Geun Kim*

Department of Biology Education, Seoul National University, Seoul 151-748, Republic of Korea

ARTICLE INFO

Article history:

Received 23 February 2013

Received in revised form 18 July 2013

Accepted 20 September 2013

Keywords:

Knotweed management

Persicaria thunbergii

Flooding depth

Soil type

Early growth stages

ABSTRACT

We investigate the effects of water level and soil type on the survival and growth of *Persicaria thunbergii* during the recruitment phase. Plants in four early growth stages (uprooted seed, seedling with cotyledon, seedlings with two and four foliage leaves) were planted under combined conditions of six water levels (dry, wet, flooding) and two soil types (loamy sand, sand). Six weeks later, the survival rates, plant heights and biomass were measured. The uprooted seeds survived under water levels lower than 6 cm. The seedlings in other growth stages could not overcome water levels twice their height. Under dry conditions, the plant height and biomass were higher in sand than in loamy sand, yet under saturated conditions, the opposite result was obtained. The water level relative to seedling height was the most critical factor for early survival, and seedling growth was also affected by soil type. Thus, *P. thunbergii* in constructed wetlands can be removed during the seedling recruitment phase using water level regulation.

© 2013 Elsevier B.V. All rights reserved.

1. Introduction

Persicaria thunbergii (Siebold & Zucc.) H. Gross ex Nakai is an annual Polygonaceae herb. This plant inhabits not only flooded environments with few environmental changes (Yun, 2007; Oh et al., 2010) but also streams and canals, which are unstable due to frequent hydrological disturbances (Park et al., 2009), with high resilience to flooding disturbance (Kim et al., 2012). *P. thunbergii* is occasionally problematic in constructed wetlands (Kim et al., 2011; Choe et al., 2010). Wetlands are especially vulnerable to plant invasion due to frequent disturbance and high nutrient levels (Rejmankova, 2011), and *P. thunbergii* can become the dominant species in constructed wetlands. However, there have been few ecological studies on *P. thunbergii*, and the lack of data makes it difficult to understand its ecological characteristics and manage its prevalence in wetlands.

To develop a technique for managing this plant, we focused on its ecological characteristics during its early life history. Early life-history components are very important for the distribution of wetland plants (Weiher and Keddy, 1995; Nicol and Ganf, 2000), especially annual plants (Noe and Zedler, 2001). This study attempts to determine the effects of water level and soil type on the survival rate and growth of *P. thunbergii* in early growth stages.

2. Materials and methods

2.1. Experiment 1. Survival and growth of seedlings under combined conditions of water level and soil type

Seeds were collected during early March in Boonwon-ri, Namjong-myon, Gwangju, Gyeonggi-do, Korea and incubated in a light glasshouse until they formed seedlings with cotyledons (3 cm high). They were planted in plastic trays (20 cm × 20 cm) under combined conditions of six water levels and two soil types. The water levels included soil moisture of 5% and 10% (dry) and water levels of 0 cm (water-logged), 3 cm, 6 cm and 9 cm above the soil surface (flooding). The water was clear, and the temperature was not regulated during the study. The soil types included two treatments: loamy sand (2.26% organic matter) comprised of 1:1 mixtures of bed-soil (for rice seedbeds, Dongbuhannong, Korea) and sand, and sand (0.89% organic matter) comprised of 1:5 mixtures of bed-soil and sand. All conditions had three replicates. Six weeks later, we measured survival rates and the plant heights and biomass of the surviving individuals. For the statistical analysis, we used multi-way ANOVA and Duncan's post hoc test (SPSS 17.0) at the 5% significance level.

2.2. Experiment 2. Survival rates for different water levels during the early growth phase

To identify the relationship between the survival rate and water level during early life history, plants were divided into four early

* Corresponding author. Tel.: +82 2 880 7896; fax: +82 2 886 2117.
E-mail address: jaegkim@snu.ac.kr (J.G. Kim).

growth stages: uprooted seed, seedlings with cotyledon (3 cm high), seedlings with two foliage leaves (7 cm high) and seedlings with four foliage leaves (13 cm high). The uprooted seeds were subjected to four flooding depths: water levels of 0 cm, 2 cm, 4 cm and 6 cm above the soil surface. The other three stages were subjected to four conditions: water levels of 0 cm, the same level as the plant height, and levels of twice and three times the plant height. Three seedlings or uprooted seeds were planted per pot ($\Phi = 15$ cm) in loamy sand, and three replicates were performed. Experiments were conducted from 5th May to 9th June. For the statistical analysis, we used one-way ANOVA and Duncan's post hoc test or an independently sampled *T*-test (SPSS 17.0) at the 5% significance level.

3. Results

3.1. Experiment 1. Survival rates and growth characteristics of seedlings under combined conditions of water level and soil type

Most seedlings planted at soil moistures of 5% or 10% or water levels of 0 cm and 3 cm survived. All seedlings planted at water levels of 6 cm and 9 cm died (Fig. 1A). Regarding plant height and biomass, there were interaction effects between water level and soil type at the 5% significance level (Fig. 1B). Under dry conditions, plant height and biomass were lower in the loamy sand than in sand. Under water-rich conditions, plant height and biomass were higher in the loamy sand than in sand.

Fig. 1. (A) Survival rates of seedlings under combined conditions of water level and soil type. (B) Plant height and biomass under dry conditions (soil moistures of 5% and 10%) and water-rich conditions (water levels of 0 cm and 3 cm). Loamy sand was a 1:1 mixture of bed-soil:sand; sand was a 1:5 mixture of bed-soil:sand. Vertical bars indicate $\pm$ SE.

Fig. 2. Survival rates (A) and plant height (B) and biomass (C) of the uprooted seeds, seedlings with cotyledon, seedlings with two foliage leaves, and seedlings with four foliage leaves under four flooding conditions relative to seedling heights. Vertical bars indicate $\pm$ SE. Alphabetic characters indicate significant differences at the 5% level based on Duncan's groups. Meaningful T-test results at the 5% level are marked with *.

3.2. Experiment 2. Survival rates for different water levels during the early growth phase

In the case of the uprooted seed, many organisms survived at water levels of 0 cm, 2 cm and 4 cm above the soil surface. However, the survival rate was only 14% at a water level of 6 cm (Fig. 2A). Although the surviving individuals at a water level of 6 cm appeared taller than the others (Fig. 2B), they were etiolated. The biomass of the surviving plants at water levels of 2 cm, 4 cm and 6 cm were 50%, 39% and 2% of that found at a water level of 0 cm, respectively (Fig. 2C). The maximum water level for the survival of uprooted seeds was approximately 4 cm, and growth was inhibited when the water level was more than 4 cm above the soil surface.

The seedlings with cotyledon, two foliage leaves and four foliage leaves survived at water levels of 0 cm and 3 cm, 0 cm and 7 cm,

and 0 cm and 13 cm, respectively (Fig. 2A). This finding indicates that seedlings can survive at water levels that are lower than their height. However, all seedlings with cotyledon, two foliage leaves and four foliage leaves died at water levels of 6 cm and 9 cm, 14 cm and 26 cm, and 26 cm and 39 cm, respectively.

4. Discussion

4.1. Survival and growth characteristics of *P. thunbergii* during the recruitment phase

This study investigated the survival traits of *P. thunbergii* during the early growth phase. In experiment 1, the seedlings survived at water levels of 0 cm and 3 cm but died at water levels of 6 cm and 9 cm (Fig. 1A). In experiment 2, the survival rate of uprooted seeds at a water level of 6 cm was only 14%, and the surviving organisms were on the verge of death (Fig. 2A). Kim et al. (2012) reported that the maximum water level of the *P. thunbergii* habitat was 4.8 cm. Our results are consistent with their results and indicate that a flooding condition of less than 5 cm was the maximum water level for survival of this plant during recruitment phase in the field. The relationship between seedling survival and water level became more obvious in experiment 2 (Fig. 2A). The survival of seedlings was related to both water level and seedling height. Seedlings did not overcome water levels twice their height. This result may seem to be common sense, but the fact that seedlings of all stages have the same resistance to flooding conditions is very important and useful, especially for wetland vegetation management.

This study also found that the growth characteristics at different water levels are clearly affected by soil type. Under dry conditions, the seedling height and biomass are higher in sand than in loamy sand. However, under water-rich conditions, the seedling height and biomass are higher in loamy sand than in sand (Fig. 1B). This difference might be related to the soil texture and organic matter content. The organic matter content of loamy sand and sand were 2.26% and 0.89%, respectively. Although we did not measure the available water content of each substrate, soil researchers have suggested that soil water retention differs by soil texture, with loamy sand retaining more water than sand (Cosby et al., 1984; Rawls et al., 1982). Thus, under dry conditions, the limiting factor for plant growth might be the available water. However, under water-rich conditions, the limiting factor might be available nutrients rather than available water. Thus, different management plans according to soil environment are necessary.

4.2. Critical factor for *P. thunbergii* survival and management practice in constructed wetland

It is well known that water is the most crucial factor for wetland plants. Water depth and fluctuation are clearly of greater importance than other factors, especially during the germination and establishment phases (Weiher and Keddy, 1995). Different species are specialized for different areas of the moisture gradient during the recruitment phase (Keddy and Ellis, 1985; Coops and Van Der Velde, 1995; Nicol and Ganf, 2000). Especially for annual plants, germination and establishment are critical for plant assemblages (Noe and Zedler, 2001). Therefore, understanding the ecological characteristics of the early phases of each plant is very important for wetland management. In this regard, this study provided additional support for the importance of water conditions for wetland annual plants. Additionally, this study contributes the practical information that *P. thunbergii* in constructed wetlands can be easily removed during the seedling recruitment phase using water-level regulation.

5. Conclusions

The water environment is the most critical factor for the survival and growth of *P. thunbergii* seedlings. They cannot overcome water levels that are 4–6 cm above the soil surface. They were able to survive at water levels equal to their height but not water levels twice their height. The growth characteristics at different water levels are clearly affected by soil type. Thus, *P. thunbergii* in constructed wetlands can be easily removed during the early growth phase using water level regulation, and different management plans should be designed according to the soil environment.

Acknowledgments

This research was supported by the Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education, Science and Technology (No. 2012R1A1A2001007), the Center for Aquatic Ecosystem Restoration (CAER) of the Eco-STAR project from the Ministry of Environment (MOE), Republic of Korea (MOE; 08-III-12), and Korea Ministry of Environment as “The Eco-Innovation Project”.

References

- Choe, I.H., Hann, B.H., Ki, K.S., 2010. The change of riverside vegetation by construction of ecological stream in Suwoncheon, Gyeonggi Province. *Korean J. Environ. Ecol.* 24, 723–734 (in Korean, with English abstract).
- Coops, H., Van Der Velde, G., 1995. Seed dispersal, germination and seedling growth of six helophyte species in relation to water-level zonation. *Freshwater Biol.* 34, 13–20.
- Cosby, B.J., Hornberger, G.M., Clapp, R.B., Ginn, T.R., 1984. A statistical exploration of the relationships of soil moisture characteristics to the physical properties of soils. *Water Resour. Res.* 20, 682–690.
- Keddy, P.A., Ellis, T.H., 1985. Seedling recruitment of 11 wetland plant species along a water level gradient: shared or distinct responses? *Can. J. Bot.* 63, 1876–1879.
- Kim, C.H., Choi, Y.E., Kim, J.W., Myung, H., Lee, S.I., 2011. 3-Year change of vegetation and life form at the man-made wetland in Sinpyeoncheon city. *Korean J. Environ. Ecol.* 25, 57–64 (in Korean, with English abstract).
- Kim, D.H., Choi, H., Kim, J.G., 2012. Occupational strategy of *Persicaria thunbergii* in riparian area: rapid recovery after harsh flooding disturbance. *J. Plant Biol.* 55, 226–232.
- Nicol, J.M., Ganf, G.G., 2000. Water regimes, seedling recruitment and establishment in three wetland plant species. *Mar. Freshwater Res.* 51, 305–309.
- Noe, G.B., Zedler, J.B., 2001. Spatio-temporal variation of salt marsh seedling establishment in relation to the abiotic and biotic environment. *J. Veg. Sci.* 12, 61–74.
- Oh, Y.J., Lee, B.M., Sohn, S.I., Lee, Y.K., Nam, H.S., Lee, S.B., Kang, C.K., Jee, H.J., 2010. Vegetation types and soil environment as affected by fallow paddy. *Korean J. Weed Sci.* 30, 405–411 (in Korean, with English abstract).
- Park, I.H., Jang, G.S., Cho, K.J., Ra, J.H., 2009. Analysis on the flora and vegetation for ecological restoration of local stream – in case of local streams in Namwon. *J. Korean Environ. Res. Technol.* 12, 153–163 (in Korean, with English abstract).
- Rawls, W.J., Brakensiek, D.L., Saxton, K.E., 1982. Estimation of soil water properties. *Trans. ASAE* 25, 1316–1320.
- Rejmankova, E., 2011. The role of macrophytes in wetland ecosystems. *J. Ecol. Field Biol.* 34, 333–345.
- Weiher, E., Keddy, P.A., 1995. The assembly of experimental wetland plant communities. *Oikos* 73, 323–335.
- Yun, K.S., 2007. Soil and Vegetation characteristics of abandoned paddy field. *J. Korean Assoc. Reg. Geogr.* 13, 129–142 (in Korean, with English abstract).