

Seasonal Distribution of Ticks in Four Habitats near the Demilitarized Zone, Gyeonggi-do (Province), Republic of Korea

Sung Tae Chong¹, Heung Chul Kim¹, In-Yong Lee², Thomas M. Kollars Jr.³, Alfredo R. Sancho⁴, William J. Sames⁵, Joon-Seok Chae⁶ and Terry A. Klein^{7,*}

¹5th Medical Detachment, 168th Multifunctional Medical Battalion, 65th Medical Brigade, Unit 15247, APO AP 96205-5247, USA; ²Department of Environmental Medical Biology, Yonsei University College of Medicine, Seoul 120-752, Korea; ³Medical Section, 415th Chemical Brigade, 814 Perimeter Rd., Greenville, SC 29605, USA; ⁴U.S. Public Health Service Commissioned Corps, 8904 Darnestown Road, Rockville, MD 20850, USA; ⁵Armed Forces Pest Management Board, WRAMC-Forest Glen Annex, Bldg. 172, 6900 Georgia Ave., NW, Washington, DC 20307-5001, USA; ⁶Veterinary Internal Medicine, College of Veterinary Medicine and Research Institute for Veterinary Science, Seoul National University, Seoul 151-742, Korea; ⁷U.S. Army Public Health Command Region-Pacific, Camp Zama, Japan; Address: 65th Medical Brigade, Unit 15281, APO AP 96205-5281, USA

Abstract: This study describes the seasonal distribution of larvae, nymph, and adult life stages for 3 species of ixodid ticks collected by tick drag and sweep methods from various habitats in the Republic of Korea (ROK). Grasses less than 0.5 m in height, including herbaceous and crawling vegetation, and deciduous, conifer, and mixed forests with abundant leaf/needle litter were surveyed at United States (US) and ROK operated military training sites and privately owned lands near the demilitarized zone from April-October, 2004 and 2005. *Haemaphysalis longicornis* Neumann adults and nymphs were more frequently collected from April-August, while those of *Haemaphysalis flava* Neumann and *Ixodes nipponensis* Kitaoka and Saito were collected more frequently from April-July and again during October. *H. longicornis* was the most frequently collected tick in grass habitats (98.9%), while *H. flava* was more frequently collected in deciduous (60.2%) and conifer (57.4%) forest habitats. While more *H. flava* (54.1%) were collected in mixed forest habitats than *H. longicornis* (35.2%), the differences were not significant. *I. nipponensis* was more frequently collected from conifer (mean 8.8) compared to deciduous (3.2) and mixed (2.4) forests.

Key words: *Haemaphysalis*, *Ixodes*, tick, seasonal distribution, habitats, Korea

INTRODUCTION

Tick-borne disease surveillance is becoming increasingly important as zoonotic tick-borne pathogens are recognized that affect man and wild and domestic animals worldwide [1-3]. In the Republic of Korea (ROK), zoonotic pathogens affecting human health, i.e., spotted fever group (SFG) rickettsia [4], *Ehrlichia* and *Anaplasma* spp. [5], *Bartonella* spp. [6,7], *Borrelia* spp. [7-9], and tick-borne encephalitis virus [10-12], have been reported from ticks. While ticks collected from birds (migratory bird surveillance) [13], mammals (rodent-borne disease surveillance and large animal surveys) [14,15], and reptiles

(wild-life studies) [16] infer tick-host and host-pathogen reservoir relationships, they are often impractical since the live capture of wild animals and birds in their natural habitat is manpower intensive and in some cases requires Biosafety level 3+ (BSL-3+) facilities (i.e., for reservoirs of Hantaan virus) [17-19]. Therefore, other methods, i.e., dragging and sweeping vegetation for questing ticks, are employed for specific environments where associated animal and bird hosts are present. Additionally, unfed larval ticks collected by these methods support evidence for potential transovarial transmission [20].

Previous studies have shown disparities in tick collection methods, as some species are frequently collected from animal hosts, while few are collected when dragging or sweeping vegetation [14]. However, dragging or sweeping vegetation for questing ticks among various habitats frequented by tick hosts provides an estimate of their geographical, habitat, seasonal, and life stage distributions and health-related risks associated with military, agriculture, and recreational activities. The pur-

•Received 20 March 2013, revised 21 April 2013, accepted 30 April 2013.

*Corresponding author (terry.a.klein2.civ@mail.mil)

© 2013, Korean Society for Parasitology and Tropical Medicine

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

pose of the present study was to provide estimates of the seasonal abundance and distribution of life stages during periods when ixodid ticks are most active. These data serve to provide information that is necessary for the development of tick-borne disease threat assessments for selected military training sites and privately owned lands as part of the 65th Medical Brigade (MED BDE) tick-borne disease surveillance program.

MATERIALS AND METHODS

Habitats

Tick-borne disease surveillance was conducted at US-ROK operated training sites and privately owned properties in northern Gyeonggi Province near the demilitarized zone (DMZ) that separates North and South Korea, from April-October 2004-2005, as described by Chong et al. [20] (Fig. 1). Habitats surveyed included grasses (<0.5 m), with various proportions of herbaceous and crawling vegetation, and deciduous, conifer (planted groves of pine, fir, larch and cedar), and mixed forests of planted and volunteer deciduous and coniferous trees with abundant leaf and/or needle litter as ground cover and limited understory of patches of grasses, herbaceous vegetation, and shrubs.

Tick collections by dragging and sweeping methods

Tick drags consisted of a 1.5 m long×1.0 m wide flannel cloth attached to a wooden dowel (1.1 m long, 2.0 cm diameter) and a small diameter of 4 m long nylon rope attached to

each end of the wooden dowel, as previously described by Daniels et al. [21] and Scoles et al. [22]. The tick sweep was similarly constructed as described by Carroll and Schmidtman [23], except that it was attached to an aluminum pole bent at a 45° angle where the 1.5 m long×1.0 m wide flannel cloth was attached along the leading edge.

Collection sites included 2 or 4 linear transects that alternated sweep and drag collections conducted concurrently and with collectors/transects separated by approximately 10 m as described by Chong et al. [20]. To reduce "border effect", collection lanes were at least 10 m distant from other habitat types, e.g., adjacent forest habitats and grass habitats. Collections were made by slowly walking and dragging or sweeping the flannel cloth over the vegetation (grasses, herbaceous vegetation, shrubs) or ground cover (i.e., leaf litter), stopping at 5 m intervals, laying the cloth on the ground, and removing the attached ticks using a fine forceps from both sides of the cloth. Nymphs and adults (n=20) were placed in 2 ml cryovials, while larvae (n=100) were placed in 2 ml cryovials containing 100% ethanol. This was repeated until each collector traveled a distance of approximately 100 m.

Ticks were transported to the Entomology Section, 5th Medical Detachment, 168th Multifunctional Medical Battalion (MMB), 65th MED BDE, where they were identified to species and developmental stage under a dissecting microscope using standard keys and current nomenclature [24,25]. The identified specimens were sent to Seoul National University for assay of selected zoonotic pathogens and these data were reported elsewhere [6,26,27].

RESULTS

A total of 19,821 ticks (larvae, nymphs, and adults) belonging to 2 genera and 3 species, *H. longicornis* (15,020), *H. flava* (3,889), and *I. nipponensis* (912), were collected from 4 habitats in northern Gyeonggi-do (Province), ROK, from April through October of 2004 and 2005 (Table 1). Overall, for all stages, *H. longicornis* [adults 865 (5.7%), nymphs 5,282 (75.8%), and larvae 8,873 (59.1%)] accounted for 75.8% of the 3 species collected from all habitats, followed by *H. flava* 19.6% [adults 133 (3.4%), nymphs 993 (25.5%), and larvae 2,763 (71.1%)] and *I. nipponensis* 4.6% [adults 35 (3.8%), nymphs 119 (13.1%), and larvae 758 (83.1%)]. Significantly more *H. longicornis* adults (n=639) and nymphs (n=4,788) were collected from grass habitats and accounted for 61.9% and 74.9%,




Fig. 1. Geographical location of tick collection sites at US installations and US and ROK operated training sites conducted near the demilitarized zone, Gyeonggi Province, April-October 2004-2005.

Table 1. Monthly mean number (total) larvae, nymph, and adult ixodid ticks collected using tick drag and sweep methods for four habitats at US-ROK operated military training sites and private lands located near the DMZ, Republic of Korea, 2004 and 2005

Habitats (No. repli- cates*)	Species	Months										Mean (Total) Numbers	Percent by habi- tats (%)						
		Apr		May		Jun		Jul		Aug				Sep		Oct			
Grass (n=98)	<i>Haemaphysalis longicornis</i>	Larvae	4.6	(37)	4.4	(88)	1.6	(26)	25.3	(557)	137.3	(1,648)	428.2	(5,138)	24.6	(197)	78.5	(7,691)	58.6
		Nymphs	184.4	(1,475)	29.2	(592)	123.1	(1,970)	29.3	(645)	7.6	(84)	1.8	(22)	0.0	(0)	48.9	(4,788)	36.5
		Adults	0.4	(3)	1.5	(30)	10.6	(170)	15.4	(339)	7.8	(94)	0.3	(3)	0.0	(0)	6.5	(639)	4.9
		Subtotal	189.4	(1,515)	35.5	(710)	135.4	(2,166)	70.0	(1,541)	152.2	(1,826)	430.3	(5,163)	24.6	(197)	133.9	(13,118)	98.9
	<i>Haemaphysalis flava</i>	Larvae	0.0	(0)	0.0	(0)	0.0	(0)	0.0	(0)	0.0	(0)	0.4	(5)	0.4	(3)	0.1	(9)	16.1
		Nymphs	0.0	(0)	1.1	(22)	0.2	(3)	0.2	(5)	0.0	(0)	0.0	(0)	1.1	(9)	0.4	(39)	69.6
		Adults	0.0	(0)	0.2	(4)	0.1	(1)	0.0	(0)	0.0	(0)	0.1	(1)	0.3	(2)	0.1	(8)	14.3
		Subtotal	0.0	(0)	1.3	(26)	0.3	(4)	0.3	(6)	0.0	(0)	0.5	(6)	1.8	(14)	0.6	(56)	0.4
	<i>Ixodes nipponensis</i>	Larvae	0.0	(0)	0.0	(0)	0.0	(0)	0.6	(13)	2.6	(31)	0.0	(0)	0.0	(0)	0.4	(44)	46.8
		Nymphs	0.6	(5)	0.8	(15)	0.2	(3)	0.0	(1)	0.1	(1)	0.1	(1)	0.1	(1)	0.3	(27)	28.7
		Adults	0.4	(3)	0.5	(9)	0.6	(10)	0.0	(0)	0.0	(0)	0.0	(0)	0.1	(1)	0.2	(23)	24.5
		Subtotal	1.0	(8)	1.2	(24)	0.8	(13)	0.6	(14)	2.7	(32)	0.1	(1)	0.3	(2)	1.0	(94)	0.7
	Total grasses	Larvae	4.6	(37)	4.4	(88)	1.6	(26)	26.0	(571)	139.9	(1,679)	428.6	(5,143)	25.0	(200)	79.0	(7,744)	58.4
		Nymphs	185.0	(1,480)	31.5	(629)	123.5	(1,976)	29.6	(651)	7.1	(85)	1.9	(23)	1.3	(10)	49.5	(4,854)	36.6
		Adults	0.8	(6)	2.2	(43)	11.3	(181)	15.4	(339)	7.8	(94)	0.3	(4)	0.4	(3)	2.5	(670)	5.0
		Total	190.4	(1,523)	38.0	(760)	136.4	(2,183)	71.0	(1,561)	154.8	(1,858)	430.8	(5,170)	26.6	(213)	135.4	(13,268)	100.0
Deciduous forests (n=68)	<i>Haemaphysalis longicornis</i>	Larvae	0.0	(0)	8.3	(33)	0.3	(4)	2.9	(23)	27.6	(442)	56.5	(226)	2.9	(58)	11.6	(786)	57.5
		Nymphs	19.3	(77)	18.0	(72)	17.1	(205)	2.0	(16)	1.1	(17)	0.8	(3)	0.0	(0)	5.7	(390)	28.6
		Adults	0.3	(1)	2.0	(8)	6.3	(75)	10.5	(75)	1.4	(22)	0.0	(0)	0.0	(0)	2.8	(190)	13.9
		Subtotal	19.5	(78)	28.3	(113)	23.7	(284)	15.4	(123)	30.1	(481)	57.3	(229)	2.9	(58)	20.1	(1,366)	34.3
	<i>Haemaphysalis flava</i>	Larvae	0.0	(0)	0.0	(0)	0.0	(0)	0.0	(0)	116.5	(1,864)	3.3	(13)	12.9	(258)	31.4	(2,138)	89.2
		Nymphs	4.3	(17)	3.3	(13)	4.8	(57)	0.6	(5)	0.6	(9)	1.0	(4)	6.4	(127)	3.4	(232)	9.7
		Adults	0.0	(0)	0.8	(3)	0.2	(2)	0.0	(0)	0.4	(6)	0.0	(0)	0.9	(17)	0.4	(28)	1.1
		Subtotal	4.3	(17)	4.0	(16)	4.9	(59)	0.6	(5)	117.4	(1,879)	4.3	(17)	20.1	(402)	35.2	(2,395)	60.2
	<i>Ixodes nipponensis</i>	Larvae	0.0	(0)	0.0	(0)	0.0	(0)	3.3	(26)	9.2	(147)	1.5	(6)	0.3	(5)	2.7	(184)	83.6
		Nymphs	0.5	(2)	1.5	(6)	1.8	(21)	0.1	(1)	0.1	(2)	0.0	(0)	0.0	(0)	0.5	(32)	14.6
		Adults	0.0	(0)	0.5	(2)	0.1	(1)	0.0	(0)	0.0	(0)	0.0	(0)	0.1	(1)	0.1	(4)	1.8
		Subtotal	0.5	(2)	2.0	(8)	1.8	(22)	3.4	(27)	9.3	(149)	1.5	(6)	0.3	(6)	3.2	(220)	5.5
	Total deciduous forests	Larvae	0.0	(0)	8.3	(33)	0.3	(4)	6.1	(49)	153.3	(2,453)	61.3	(245)	16.1	(321)	45.8	(3,105)	78.0
		Nymphs	24.0	(96)	22.8	(91)	23.6	(283)	2.8	(22)	1.8	(28)	1.8	(7)	6.4	(127)	9.6	(654)	16.4
		Adults	0.3	(1)	3.3	(13)	6.5	(78)	10.5	(84)	1.8	(28)	0.0	(0)	0.9	(18)	3.3	(222)	5.6
		Total	24.3	(97)	34.3	(137)	30.4	(365)	19.4	(155)	156.8	(2,509)	63.0	(252)	23.3	(466)	58.5	(3,981)	100.0
Conifer forests (n=27)	<i>Haemaphysalis longicornis</i>	Larvae	0.0	(0)	0.0	(0)	0.0	(0)	0.2	(2)	5.9	(47)	0.8	(3)	0.4	(3)	1.0	(55)	42.0
		Nymphs	0.5	(1)	1.5	(34)	1.1	(13)	0.5	(6)	0.6	(5)	0.0	(0)	0.0	(0)	1.1	(59)	45.0
		Adults	0.0	(0)	0.0	(0)	0.4	(5)	0.9	(11)	0.1	(1)	0.0	(0)	0.0	(0)	0.3	(17)	13.0
		Subtotal	0.5	(1)	1.5	(34)	1.5	(18)	1.6	(19)	6.6	(53)	0.8	(3)	0.4	(3)	2.4	(131)	9.2
	<i>Haemaphysalis flava</i>	Larvae	0.0	(0)	0.0	(0)	0.0	(0)	4.3	(52)	38.8	(310)	3.3	(13)	0.9	(7)	7.1	(382)	46.9
		Nymphs	4.5	(9)	11.5	(167)	2.3	(28)	2.7	(32)	1.9	(15)	0.5	(2)	16.3	(130)	7.1	(383)	47.0
		Adults	1.0	(2)	1.8	(16)	0.6	(7)	0.7	(8)	1.4	(11)	0.5	(2)	0.5	(4)	0.9	(50)	6.1
		Subtotal	5.5	(11)	13.3	(183)	2.9	(35)	7.7	(92)	42.0	(336)	4.3	(17)	17.6	(141)	15.1	(815)	57.4
	<i>Ixodes nipponensis</i>	Larvae	0.0	(0)	0.0	(0)	0.0	(0)	35.8	(430)	2.4	(19)	0.3	(1)	0.3	(2)	8.4	(452)	95.4
		Nymphs	0.0	(0)	0.3	(6)	0.3	(4)	0.2	(2)	0.8	(6)	0.0	(0)	0.0	(0)	0.3	(18)	3.8
		Adults	0.0	(0)	0.1	(1)	0.0	(0)	0.2	(2)	0.0	(0)	0.0	(0)	0.1	(1)	0.1	(4)	0.8
		Subtotal	0.0	(0)	0.3	(7)	0.3	(4)	36.2	(434)	3.1	(25)	0.3	(1)	0.4	(3)	8.8	(474)	33.4
	Total conifer forests	Larvae	0.0	(0)	0.0	(0)	0.0	(0)	40.3	(484)	47.0	(376)	4.3	(17)	1.5	(12)	16.5	(889)	62.6
		Nymphs	5.0	(10)	13.3	(207)	3.8	(45)	3.3	(40)	3.3	(26)	0.5	(2)	16.3	(130)	8.5	(460)	32.4
		Adults	1.0	(3)	1.8	(17)	1.0	(12)	1.8	(21)	1.5	(12)	0.5	(2)	0.6	(5)	1.3	(72)	5.0
		Total	6.0	(12)	15.2	(224)	4.8	(57)	45.4	(545)	51.8	(414)	5.3	(21)	18.4	(147)	26.3	(1,420)	100.0

(Continued to the next page)

respectively, of all adults and nymphs collected from that habitat. *H. flava* adults and nymphs were more frequently collected from conifer forests (433/532; 81.4%) and mixed forests (386/496; 77.8%). Mean numbers of *I. nipponensis* adults (0.1) and nymphs (0.3-0.8) were similarly collected from all habitats. Significantly more *H. longicornis* adults and nymphs (ANOVA, F value= 10.88, df= 3, $P < 0.001$) (mean no. 55.4) were collect-

ed from grass habitats, compared to deciduous (8.5), mixed (1.4), and conifer (1.2) forests. Significantly more *H. flava* ($F = 15.98$, $df = 3$, $P < 0.001$) were collected from conifer (8.0) and mixed (7.4) forest habitats, while few were collected from deciduous forest (3.8) and grass (0.5) habitats. While *I. nipponensis* nymphs were much less commonly collected from all habitats, significantly more were collected from mixed forests (mean

Table 1. (Continued from the previous page) Monthly mean number (total) larvae, nymph, and adult ixodid ticks collected using tick drag and sweep methods for four habitats at US-ROK operated military training sites and private lands located near the DMZ, Republic of Korea, 2004 and 2005

Habitats (No. replicates*)	Species		Months												Mean (Total) Numbers	Percent by habi- tats (%)			
			Apr		May		Jun		Jul		Aug		Sep				Oct		
Mixed forests (n=52)	<i>Haemaphysalis longicornis</i>	Larvae	0.0	(0)	0.0	(0)	0.0	(0)	0.1	(1)	85.0	(340)	0.0	(0)	0.0	(0)	6.6	(341)	84.2
		Nymphs	1.1	(18)	1.3	(10)	0.8	(3)	0.1	(2)	3.0	(12)	0.0	(0)	0.0	(0)	0.9	(45)	11.1
		Adults	0.0	(0)	0.0	(0)	0.3	(1)	0.5	(9)	2.3	(9)	0.0	(0)	0.0	(0)	0.4	(19)	4.7
		Subtotal	1.1	(18)	1.3	(10)	1.0	(4)	0.6	(12)	90.3	(361)	0.0	(0)	0.0	(0)	7.8	(405)	35.2
	<i>Haemaphysalis flava</i>	Larvae	0.0	(0)	1.5	(12)	0.0	(0)	0.3	(6)	54.8	(219)	0.0	(0)	0.0	(0)	4.6	(237)	38.1
		Nymphs	2.8	(44)	18.3	(146)	11.5	(46)	4.7	(93)	2.5	(10)	0.0	(0)	0.0	(0)	6.5	(339)	54.4
		Adults	0.5	(8)	3.6	(29)	0.8	(3)	0.3	(6)	0.3	(1)	0.0	(0)	0.0	(0)	0.9	(47)	7.5
		Subtotal	3.3	(52)	23.4	(187)	12.3	(49)	5.3	(105)	57.5	(230)	0.0	(0)	0.0	(0)	12.0	(623)	54.1
	<i>Ixodes nipponensis</i>	Larvae	0.0	(0)	0.0	(0)	0.0	(0)	0.9	(18)	15.0	(18)	0.0	(0)	0.0	(0)	1.5	(78)	62.9
		Nymphs	0.0	(0)	2.3	(18)	3.8	(15)	0.4	(8)	0.3	(1)	0.0	(0)	0.0	(0)	0.8	(42)	33.9
		Adults	0.2	(3)	0.1	(1)	0.0	(0)	0.0	(0)	0.0	(0)	0.0	(0)	0.0	(0)	0.1	(4)	3.2
		Subtotal	0.2	(3)	2.4	(19)	3.8	(15)	1.3	(26)	15.3	(61)	0.0	(0)	0.0	(0)	2.4	(124)	10.7
	Total mixed forests	Larvae	0.0	(0)	1.5	(12)	0.0	(0)	1.3	(25)	154.8	(619)	0.0	(0)	0.0	(0)	12.6	(656)	56.9
		Nymphs	3.9	(62)	21.8	(174)	16.0	(64)	5.2	(103)	5.8	(23)	0.0	(0)	0.0	(0)	8.2	(426)	37.0
		Adults	0.7	(11)	3.8	(30)	1.0	(4)	0.8	(15)	2.5	(10)	0.0	(0)	0.0	(0)	1.3	(70)	6.1
		Total	4.6	(73)	27.0	(216)	17.0	(68)	7.2	(143)	163.0	(652)	0.0	(0)	0.0	(0)	22.2	(1,152)	100.0
Total (n=272)	<i>Haemaphysalis longicornis</i>	Larvae	1.2	(37)	3.0	(121)	0.7	(30)	9.4	(583)	61.9	(2,477)	268.4	(5,367)	7.2	(258)	32.6	(8,873)	59.1
		Nymphs	52.4	(1,571)	17.7	(708)	49.8	(2,191)	10.8	(669)	3.0	(118)	1.3	(25)	0.0	(0)	19.4	(5,282)	35.2
		Adults	0.1	(4)	1.0	(38)	5.7	(251)	7.1	(443)	3.2	(126)	0.2	(3)	0.0	(0)	3.2	(865)	5.7
		Subtotal	53.7	(1,612)	21.7	(867)	56.2	(2,472)	27.3	(1,695)	68.0	(2,721)	269.8	(5,395)	7.2	(258)	55.2	(15,020)	75.8
	<i>Haemaphysalis flava</i>	Larvae	0.0	(0)	0.3	(12)	0.0	(0)	1.0	(59)	59.8	(2,393)	1.6	(31)	7.4	(268)	10.2	(2,763)	71.1
		Nymphs	2.3	(70)	8.7	(348)	3.0	(134)	2.2	(135)	0.9	(34)	0.3	(6)	7.4	(266)	3.7	(993)	25.5
		Adults	0.3	(10)	1.3	(52)	0.3	(13)	0.2	(14)	0.5	(18)	0.2	(3)	0.6	(23)	0.5	(133)	3.4
		Subtotal	2.7	(80)	10.3	(412)	3.3	(147)	3.4	(208)	61.1	(2,445)	2.0	(40)	15.5	(557)	14.3	(3,889)	19.6
	<i>Ixodes nipponensis</i>	Larvae	0.0	(0)	0.0	(0)	0.0	(0)	7.9	(487)	6.4	(257)	0.4	(7)	0.2	(7)	2.8	(758)	83.1
		Nymphs	0.2	(7)	1.1	(45)	1.0	(43)	0.2	(12)	0.3	(10)	0.1	(1)	0.1	(1)	0.4	(119)	13.1
		Adults	0.2	(6)	0.3	(13)	0.3	(11)	0.1	(2)	0.0	(0)	0.0	(0)	0.1	(3)	0.1	(35)	3.8
		Subtotal	0.4	(13)	1.5	(58)	1.2	(54)	8.1	(501)	6.7	(267)	0.4	(8)	0.3	(11)	3.4	(912)	4.6
	Grand total	Larvae	1.2	(37)	3.3	(133)	0.7	(30)	18.2	(1,129)	128.2	(5,127)	270.3	(5,405)	14.8	(533)	45.6	(12,394)	62.5
		Nymphs	54.9	(1,648)	27.5	(1,101)	53.8	(2,368)	13.2	(816)	4.1	(162)	1.6	(32)	7.4	(267)	23.5	(6,394)	32.3
		Adults	0.7	(20)	2.6	(103)	6.3	(275)	7.4	(456)	3.6	(144)	0.3	(6)	0.7	(26)	3.8	(1,033)	5.2
		Total	56.8	(1,705)	33.4	(1,337)	60.8	(2,673)	38.8	(2,404)	135.8	(5,433)	272.2	(5,443)	22.9	(826)	72.9	(19,821)	100.0

*Total number of replicates by 100 m transect for each habitat.

0.8), while overall, there were no significant differences for adults and nymphs combined from all habitats sampled ($F=1.44$, $df=3$, $P=0.23$).

The seasonal distributions of *H. longicornis*, *H. flava*, and *I. nipponensis* collected by both drag and sweep are shown in Table 1 and Fig. 2. In general, for all habitats, *H. longicornis* adults were collected more frequently from June-August, while nymphs were most prevalent from April-June. Adult *H. flava* populations peaked in May, with low numbers collected during the remaining months. Mean numbers of *H. flava* nymphs peaked in May, declined through September, and then peaked again in October. Mean numbers of *I. nipponensis* adult and nymph populations peaked during May-June, with fewer numbers collected during April and July-September, with increases in adult numbers during October.

Large numbers of *H. longicornis* larvae first appeared in August, peaked in September, and few were collected during Oc-

tober. Large numbers of *H. flava* larvae were collected only during August that may have resulted in increased numbers of nymphs collected during October. *I. nipponensis* larval populations peaked in July-August with few collected during the remaining months. While variable patterns were observed for the different habitats, these differences were likely due to host habitat preferences and, in some cases, low numbers collected.

DISCUSSION

The ecology of the ROK following the Korean War consisted of grass and herbaceous covered hills and mountains with occasional scattered trees that provided limited habitat and refuge for medium to large mammals and forest dwelling bird populations. As a result of a national tree planting policy instituted in the 1960s, the hillsides and mountains were populated with planted groves of conifer, deciduous, and later mixed




Fig. 2. Seasonal distribution of 3 tick species collected from 4 different habitats at US and ROK operated military training sites and installations and private lands located near the DMZ, Republic of Korea, from April through October 2004 and 2005 (GR, grasses and portions of herbaceous and crawling vegetation; DF, deciduous forest; CF, conifer forest; MF, mixed forest).

planted and volunteer forests with an understory of associated shrubs and small open areas of grasses and herbaceous vegetation. These ecological changes provided increased habitat, harborage, and an enrichment of flora and fauna provided for the expansion, diversity, and increased numbers of mammals and birds. Concurrently, the geographical dispersion of mammals and birds resulted in the potential for increased tick populations and diversity and prevalence of associated zoonotic tick-borne pathogens that heightens the medical threat for soldiers conducting military exercises and local populations performing agricultural and recreational activities. To evaluate the significance of these changes, a tick-borne disease surveillance program was established to identify the seasonal, geographical, and habitat distributions, relative abundance, and prevalence of tick-associated zoonotic pathogens.

Host and preferred habitat associations are principal for the

distribution of ticks. *H. longicornis* is associated with the large wild and domestic mammals and was collected primarily in habitats consisting of grasses and other herbaceous vegetation frequented by larger herbivores. Hosts of *H. flava* are associated with small/medium-sized mammals and resident/migratory birds and were collected primarily in forest habitats. *I. nipponensis* is a 2-host tick with larvae and nymphs feeding on small/medium-sized mammals, while nymphs and adults blood feed on medium/large mammals. Neither sweeping nor dragging methods were highly effective at collecting *I. nipponensis* [14,20]. However, more than 98.9% of all ticks collected from live captured small mammals (rodents, shrews, and squirrels) in tall grass and herbaceous vegetation in northern Gyeonggi Province was *I. nipponensis* (mostly larvae, a few nymphs, and no adults), while only one (<0.1%) *H. flava*, no *H. longicornis*, and 1.1% *Ixodes pomerantzevi* Serdyukova were

collected [14]. *I. pomerantzevi* is a nest species and has only been reported from small mammals (squirrels and rodents). Adult and nymph *I. nipponensis* were more frequently collected in forest habitats, indicating their association with hosts that utilize forest habitats for refuge but perhaps feed or seek prey in open areas with grasses and other herbaceous vegetation.

Unmanaged lands that consist of open areas of tall grasses and herbaceous and crawling vegetation and forested hillsides/mountains harbor large populations of rodents and other small-large domestic (e.g., cats, dogs, cattle, and horses) and wild mammals (e.g., feral cats, raccoon dogs, deer, and wild pigs) and indigenous and migratory birds that serve as hosts for ticks and other ectoparasites [13,14]. Human activities in these areas, whether military or civilian, increases exposure to questing ticks that are potentially infected with tick-borne pathogens. While human activities occur year round, e.g., hiking, construction, agriculture, and military training, they are more common during the months when ticks are active from early spring through late fall.

Tick drag and sweep methods provide estimates of population densities for selected species of ticks and associated disease risks. These data can be used to estimate population densities, identify and determine the prevalence of tick-borne pathogens, determine the potential for transovarial transmission and pathogen maintenance in host and vector populations, and also serve to estimate the potential for transmission of tick-borne pathogens to military and civilian populations as a necessary step for developing disease threat analyses. Tick drag and sweep methods, in addition to collections of ectoparasites from small/large mammals and birds, form the basis of a comprehensive tick-borne disease surveillance program that identifies tick-borne health threats and provides for the development of diseases risk reduction strategies to reduce tick-borne infections among US personnel deployed to the ROK. Finally, in collaboration with US counterparts, local universities, and the Korean Ministry of Health professionals for the identification of ticks and the identification and detection of tick-borne pathogens, these data provides the US military and ROK Ministry of Health with a comprehensive knowledge for endemic tick-borne pathogens and epidemiology of tick-associated diseases in the ROK.

ACKNOWLEDGMENTS

We thank Col. Hee-Choon Sam Lee, Chief, Force Health

Protection and Preventive Medicine (FHP&PM), 65th MED BDE, for his dedicated support. We sincerely thank Ms. Suk-Hee Yi, FHP&PM, 65th MED BDE, for providing GIS maps of collection sites and data statistical analysis. We thank Maj. Lisa L. O'Brian and Cpt. Brett W Collier, 5th Medical Detachment Commanders and their staff for assistance with field collections of ticks throughout the study. We also thank Dr. Joel Gaydos, Armed Forces Health Surveillance Center, Global Emerging Infections Surveillance and Response System (AFHSC-GEIS), Silver Spring, MD, for his support throughout this study.

Funding for this work was provided through the joint partnership between the AFHSC-GEIS, Silver Spring, MD, the 65th MED BDE, and the National Center for Medical Intelligence, Ft Detrick, MD.

The opinions and assertions contained herein are those of the authors and are not to be construed as official or reflecting the views of the Department of the Army or the Department of Defense.

REFERENCES

1. Murphy GL, Ewing SA, Whitworth LC, Fox JC, Kocan AA. A molecular and serologic survey of *Ehrlichia canis*, *E. chaffeensis*, and *E. ewingii* in dogs and ticks from Oklahoma. *Vet Parasitol* 1998; 79: 325-339.
2. Chiba N, Osada M, Komoro K, Mizutani T, Kariwa H, Takashima I. Protection against tick-borne encephalitis virus isolated in Japan by active and passive immunization. *Vaccine* 1999; 17: 1532-1539.
3. Alekseev AN, Dubinina HV, Van De Pol I, Schouls LM. Identification of *Ehrlichia* species and *Borrelia burgdorferi* in *Ixodes* ticks in the Baltic Regions of Russia. *J Clin Microbiol* 2001; 39: 2237-2242.
4. Lee JH, Park HS, Jung KD, Jang WJ, Koh SE, Kang SS, Lee IY, Lee WJ, Kim BJ, Kook YH, Park KH, Lee SH. Identification of the spotted fever group rickettsiae detected from *Haemaphysalis longicornis* in Korea. *Microbiol Immunol* 2003; 47: 301-304.
5. Chae JS, Kim CM, Kim EH, Hur EJ, Klein TA, Kang TK, Lee HC, Song JW. Molecular epidemiological study for tick-borne disease (*Ehrlichia* and *Anaplasma* species) surveillance at selected U.S. military training sites/installations in Korea. *Ann NY Acad Sci* 2003; 990: 118-125.
6. Kim CM, Kim JY, Yi YH, Lee MJ, Cho MR, Shah DH, Klein TA, Kim HC, Song JW, Chong ST, O'Guinn ML, Lee JS, Lee IY, Park JH, Chae JS. Detection of *Bartonella* species from ticks, mites and small mammals in Korea. *J Vet Sci* 2005; 6: 327-334.
7. Kang JG, Kim HC, Choi CY, Nam HY, Chae HY, Klein TA, Ko SJ, Chae JS. Molecular detection of *Anaplasma*, *Bartonella* and *Borrelia* species in ticks collected from migratory birds from Hong-do

- Island, Republic of Korea. Vector-Borne Zoonotic Dis 2013; 13: 215-225.
8. Park KH, Lee SH, Won WJ, Jang WJ, Chang WH. Isolation of *Borrelia burgdorferi*, the causative agent of Lyme disease, from *Ixodes* ticks in Korea. J Korean Soc Microbiol 1992; 27: 307-312.
 9. Kee S, Hwang KJ, Oh HB, Kim MB, Shim JC, Ree HI, Park KS. Isolation and identification of *Borrelia burgdorferi* in Korea. J Korean Soc Microbiol 1994; 29: 301-310.
 10. Kim SY, Yun SM, Han MG, Lee IY, Lee NY, Jeong YE, Lee BC, Ju YR. Isolation of tick-borne encephalitis viruses from wild rodents, South Korea. Vector-borne Zoonotic Dis 2008; 8: 7-13.
 11. Kim SY, Jeong YE, Yun SM, Lee IY, Han MG, Ju YR. Molecular evidence for tick-borne encephalitis virus in ticks in South Korea. Med Vet Entomol 2009; 23: 15-20.
 12. Ko S, Kang JG, Kim SY, Klein TA, Kim HC, Chong ST, Sames WJ, Yun SM, Ju YR, Chae JS. Prevalence of tick-borne encephalitis virus in ticks from southern Korea. J Vet Sci 2010; 11: 197-203.
 13. Kim HC, Ko S, Choi CY, Nam HY, Chae HY, Chong ST, Klein TA, Sames WJ, Robbins RG, Chae JS. Migratory bird tick surveillance, including a new record of *Haemaphysalis ornithophila* Hoogstraal and Kohls 1959 (Acari: Ixodidae) from Hong-do (Hong Island), Republic of Korea. Syst Appl Acarol 2009; 14: 3-10.
 14. Kim HC, Chong ST, Sames WJ, Nunn PV, Wolf SP, Robbins RG, Klein TA. Tick surveillance of small mammals captured in Gyeonggi and Gangwon Provinces, Republic of Korea, 2004-2008. Syst Appl Acarol 2010; 15: 100-108.
 15. Kim HC, Han SH, Chong ST, Klein TA, Choi CY, Nam HY, Chae HY, Lee H, Ko SJ, Kang JG, Chae JS. Tick collected from selected mammalian hosts surveyed in the Republic of Korea during 2008-2009. Korean J Parasitol 2011; 49: 331-335.
 16. Fajfer M. Acari (Chelicerata)-Parasites of Reptiles, Acarina 2012; 20: 108-129.
 17. Song JW, Moon SS, Gu SH, Song KJ, Baek LJ, Kim HC, Kijek T, O'Guinn ML, Lee JS, Turell MJ, Klein TA. Hemorrhagic fever with renal syndrome in 4 US soldiers, South Korea, 2005. Emerg Infect Dis 2009; 15: 1833-1836.
 18. Sames WJ, Klein TA, Kim HC, Gu SH, Kang HJ, Shim SH, Ha SJ, Chong ST, Lee IY, Richards AL, Yi SH, Song JW. Serological surveillance of scrub typhus, murine typhus, and leptospirosis in small mammals captured at Twin Bridges training area, Gyeonggi Province, Republic of Korea, 2005-2007. Mil Med 2010; 175: 48-54.
 19. Klein TA, Kang HJ, Gu SH, Moon SS, Shim SH, Park YM, Lee SY, Kim HC, Chong ST, O'Guinn ML, Lee JS, Turell MJ, Song JW. Hantaan virus surveillance targeting small mammals at Dagmar North training area, Gyeonggi Province, Republic of Korea, 2001-2005. J Vector Ecol 2011; 36: 373-381.
 20. Chong ST, Kim HC, Lee IY, Kollars TM, Jr., Sancho AR, Sames WJ, Klein TA. Comparison of dragging and sweeping methods for collecting ticks and determining their seasonal distributions for various habitats, Gyeonggi Province, Republic of Korea. J Med Entomol 2013; 50: 611-618.
 21. Daniels TJ, Boccia TM, Varde S, Marcus J, Le J, Bucher DJ, Falco RC, Schwartz I. Geographic risk for Lyme disease and human granulocytic ehrlichiosis in southern New York State. Appl Environ Microbiol 1998; 64: 4663-4669.
 22. Scoles GA, Papero M, Beati L, Fish D. A relapsing fever group spirochete transmitted by *Ixodes scapularis* ticks. Vectorborne Zoonotic Dis 2001; 1: 21-34.
 23. Carroll JF, Schmidtmann ET. Tick sweep: Modification of the tick drag-flag method for sampling nymphs of the deer tick (Acari: Ixodidae). J Med Entomol 1992; 29: 352-355.
 24. Yamaguti N, Tipton VJ, Keegan HL, Toshioka S. Ticks of Japan, Korea, and the Ryukyu Islands. Brigham Young University Science Bulletin 1971; 15: 1-226.
 25. Robbins RG, Keirans JK. Systematics and Ecology of the Subgenus *Ixodiopsis* (Acari: Ixodidae: Ixodes). Thomas Say Foundation Monograph XIV. Lanham, Maryland. Entomological Society of America. 1992, viii + 159 pp.
 26. Chae JS, Yu DH, Shringi S, Klein TA, Kim HC, Chong ST, Lee IY, Foley J. Microbial pathogens in ticks, rodents, and a shrew in northern Gyeonggi-do near the DMZ, Korea. J Vet Sci 2008; 9: 285-293.
 27. Kim CM, Yi YH, Yu DH, Lee MJ, Cho MR, Desai AR, Shringi S, Klein TA, Kim HC, Song JW, Baek LJ, Chong ST, O'Guinn ML, Lee JS, Lee IY, Park JH, Foley J, Chae JS. Tick-borne rickettsial pathogens in ticks and small mammals in Korea. Applied Environ Microbiol 2006; 72: 5766-5776.

